

m. G. Zwick
A-27-44

VIII.—*A Synonymic List of the Ants of the Ethiopian Region.* BY WM. M. WHEELER

IX.—*A Synonymic List of the Ants of the Malagasy Region.* BY WM. M. WHEELER

BULLETIN

OF

THE AMERICAN MUSEUM OF NATURAL HISTORY

VOL. XLV, pp. 711-1055.

New York

Issued October 25, 1922

Extracted from 'Ants of the American Congo Expedition.
A Contribution to the Myrmecology of Africa.' By William Morton Wheeler, with the collaboration of J. Bequaert, I. W. Bailey, F. Santachi, and W. M. Mann. Parts I-IX.

VIII.—A SYNONYMIC LIST OF THE ANTS OF THE ETHIOPIAN REGION

By WM. M. WHEELER

The following catalogue of the known Ethiopian and Malagasy Formicidae had its inception in an attempt to master the abundant and widely scattered literature as a basis for the study of the Congo ants collected by Messrs. Lang, Chapin, and Bequaert. The attempt was the more urgent because no general list had been published since the appearance of the seventh volume of Dalla Torre's '*Catalogus Hymenopterorum*,' which included the forms known in 1890. He cites only 228 Ethiopian and 119 Malagasy species. It will be seen that the numbers have risen during the past thirty years to 920 and 237 respectively, almost exclusively as the result of the untiring labors of five myrmecologists: Mayr, Emery, Forel, Santschi, and Arnold. Had the varieties and subspecies been included in the foregoing estimates, the numbers would be still more impressive. Those who may believe that little entomological exploration has been carried out in the dark continent will also be astonished when they scan the lists of localities and the names of the collectors who have secured the materials for the taxonomists. It almost seems as if ant-collectors had been more numerous and more diligent in Africa than in North America!

I have endeavored to include all the literature on Ethiopian and Malagasy ants down to January 1st, 1920, together with some of the papers that have since appeared, but no claim for completeness is made for this year. Except in a very few instances the references have been checked up with the original papers by Dr. Bequaert, who has also gone to much pains in looking up species not contained in my original list, in arranging the species and genera and in many other ways relieving me of the meticulous and often exasperating labor of giving the catalogue its present appearance.

The localities recorded for the various forms have been grouped geographically according to the political divisions in use at the beginning of 1914. So far as possible the spelling of the geographical names has been made uniform throughout, though this has been very difficult owing to the typographical and other errors so numerous in taxonomic papers.

With Emery and other zoologists I have called the category below the species "subspecies," not "race," or "stirps," as is the custom with Forel, Santschi and Arnold. I have, moreover, given both the subspecific and varietal names the same gender as the corresponding specific names, departing in this respect from the procedure of other myrmecologists, who treat the subspecific or at least the varietal name as feminine,

irrespective of the gender of the specific name. My reasons for this departure are two-fold: First, ornithologists, mammalogists, herpetologists and others who employ trinomials make the subspecific as well as the specific names agree with the generic name in gender, and I am unable



MAP 47. Showing localities in the Belgian Congo where ants have been collected.

to see why quadrinomials should be treated differently. Second, the status of the specific, subspecific and varietal categories in myrmecography are still in a state of flux, and since different writers and even the same writer on different occasions are in the habit of exalting varie-

ties or subspecies to specific rank and degrading species to varietal or subspecific rank—a process which must continue till knowledge of a particular species-complex sufficient to establish the precise dignity of its various categories is attained—it is obviously preferable to keep the names all of one gender.

The confusion introduced into the literature by this at present unavoidable inconsistency and instability in the employment of specific, subspecific and varietal names makes it necessary in the compilation of such a catalogue as the following to choose between different appreciations or even to rely altogether on one's own interpretation. This is particularly the case in dealing with large, complex genera such as *Pheidole*, *Crematogaster*, *Monomorium*, *Tetramorium*, *Camponotus*, and *Polyrhachis*. I gladly admit that my judgment may have been at fault in several of my appraisals of taxonomic status, especially of forms unknown to me in nature. A catalogue, however, is not a place for disputation or the weighing of evidence but merely a condensed expression of a small fragment of our present taxonomic knowledge, compiled under rigid limitations of space and conventions of arrangement. That my work will be subjected to criticism by those who have never attempted to deal with a similar body of taxonomic literature, is to be expected, but I am certain that students will find it very useful, at least till it is absorbed at some future time in a much ampler and more perfect conspectus of the Ethiopian and Malagasy ant-fauna.

APPROXIMATE LOCATION OF AFRICAN LOCALITIES, RIVERS, MOUNTAINS,
LAKES, ETC., MENTIONED IN THE CATALOGUE OF ETHIOPIAN ANTS
AND ELSEWHERE IN THE PAPER¹

Ababis.—22° 10' S., 15° 45' E.	Ambelokudi.—Near Pawa, Belgian Congo.
Abaja (L.).—6° 30' N., 38° E.	Amboni (R.).—0° 20' S., 36° 55' E.
Abba (I.).—13° 20' N., 32° 40' E.	Ambriz.—7° 45' S., 13° 5' E.
Aberdare (Mts.).—0° 30' S., 37° E.	Amu (R.).—7° 20' N., 1° 10' E.
Aberdeen.—32° 25' S., 24° 5' E.	Angra Pequena.—26° 40' S., 15° 10' E.; same as Lüderitz Bay.
Aberio.—10° N., 44° 35' E.	Angu.—3° 30' N., 24° 20' E.
Abo.—5° 30' N., 6° 25' E.	Annobon (I.).—1° 15' S., 6° E.
Abo (R.).—4° 15' N., 9° 45' E.	Anseba (R.).—15° to 17° 10' N., 38° 45' to 39° E.
Abou.—8° 50' N., 38° 35' E.	Antongil Bay.—15° 30' S., 49° 50' E.
*Abuker.—Near Dire Dawa, Abyssinia.	Aouache (R.), same as Hauash (R.).
Aburi.—5° 45' N., 0° 10' W.	Api.—3° 40' N., 25° 25' E.
Addah.—5° 55' N., 0° 35' E.	Arasab (R.).—26° 55' to 27° 10' S., 16° 15' to 16° 35' E.
Aden.—12° 50' N., 45° E.	Arigalgalu.—4° 25' N., 39° 55' E.
Adi Caie.—14° 50' N., 39° 20' E.	Artesia.—24° 5' S., 26° 20' E.
Adis-Abeba.—9° N., 38° 45' E.	Arusha-chini.—3° 35' S., 37° 25' E.
Adi Ugri.—14° 55' N., 38° 50' E.	Arussi Galla, Canale Gudda.—7° 30' N., 40° 15' E.
Adua.—14° 10' N., 38° 55' E.	Aruwimi (R.).—1° 20' N., 27° 40' E.
Agege.—6° 40' N., 3° 20' E.	Asmara.—15° 20' N., 39° E.
Agouagon.—8° N., 2° 20' E.	Assab.—13° 5' N., 42° 50' E.
*Aguagua.—Lower Dahomey. Probably misspelling for Agouagon.	Assinie.—5° N., 3° 20' W.
Aikota.—15° 10' N., 37° 5' E.	Assuan.—24° 5' N., 32° 50' E.
Ain (Mt.).—12° 45' N., 33° 5' E.	Atbara (R.).—12° 30' to 17° 40' N., 34° to 37° E.
Akaki (R.).—8° 50' N., 38° 40' E.	Athi Plain.—1° 20' S., 37° 10' E.
Akenge.—2° 55' N., 26° 50' E.	Auata (R.).—5° 15' to 6° N., 38° 50' to 39° 10' E.
Akono Linga.—3° 55' N., 12° 45' E.	Avakubi.—1° 20' N., 27° 40' E.
Akra.—5° 40' N., 0° 15' W.	Babeyru.—1° 55' N., 27° 40' E.
Akropong.—6° N., 0° 5' W.	Bachi-Shombe.—4° 55' S., 20° 35' E.
Akwapim (Mts.).—5° 50' N., 0° 20' W.	Bafuka.—4° 20' N., 27° 50' E.
Albany.—33° 20' S., 26° 25' E.	Bafwabaka.—2° 10' N., 27° 50' E.
Albert Edward (L.).—0° to 0° 30' S., 29° 30' E.	Bafwalipa.—1° 30' N., 27° 45' E.
Alen.—2° 5' N., 11° E.	Bafwasende.—1° 10' N., 27° 15' E.
Algoa Bay.—33° 45' S., 25° 45' E.; same as Port Elizabeth.	Bagamoyo.—6° 25' S., 38° 55' E.
Algota.—15° 45' N., 38° 55' E.	Baguirmi.—12° 20' N., 15° to 17° E.
Alto Queta.—9° 15' S., 14° 55' E.	
Amani.—5° 5' S. 38° 40' E.	
Amanzimtoti.—30° 5' S., 30° 50' E.	
Amarr Burgi.—5° 25' N., 37° 55' E.	
Amatongas Forest.—19° S., 33° 40' E.	
Ambás Bay.—4° N., 9° 10' E.	

¹Localities marked with an asterisk have not been found on any map.

- Bahr-el-Salaam.—12° 55' to 13° 50' N., 36° 10' to 37° 30' E.
- *Bakaia.—Between Nyangwe and Stanleyville, Belgian Congo.
- Bakusu.—3° 35' S., 25° 30' E.
- Balgowan.—29° 20' S., 30° 5' E.
- Balla Balla.—20° 25' S., 29° 20' E.
- *Balli Neck Pass.—Ex-German Southwest Africa.
- Bamako.—12° 40' N., 7° 55' W.
- *Bamayanga.—Belgian Congo.
- Bamba.—7° S., 13° 40' E.
- Bambaya.—10° 45' N., 13° 35' W.
- Bamu (I.).—4° 25' S., 15° 30' E.
- Banalia.—1° 30' N., 25° 40' E.
- Banana (Lower Congo).—6° S., 12° 20' E.
- Banana (Ituri Forest).—1° 20' N., 29° 15' E.
- Banas.—3° 55' N., 40° 15' E.
- Bangala, see Nouvelle Anvers.
- Bangweolo (L.).—11° S., 30° E.
- Banza Masola.—5° 10' S., 13° E.
- Banzville.—4° 15' N., 21° 10' E.
- Barawa.—1° 5' N., 44° 5' E.
- Barberton.—25° 50' S., 31° E.
- Bargunett (R.), Mt. Kenia.—0°, 37° E.
- Barikiwa.—9° 30' S., 37° 55' E.
- *Barko.—Abyssinia.
- Barombi (L.).—4° 40' N., 0° 25' E.
- Barumbu.—1° 10' N., 23° 20' E.
- Basoko.—1° 20' N., 23° 35' E.
- Bassam, see Grand Bassam.
- Bass Marle, see Stephanie (L.).
- Bass Narok, see Rudolf (L.).
- Basutoland.—28° 35' to 30° 35' S., 27° 5' to 29° 20' E.
- Bata.—1° 50' N., 9° 45' E.
- Batama.—1° N., 26° 40' E.
- Batanga.—2° 50' N., 9° 55' E.
- Batempa.—5° S., 23° 45' E.
- Bathurst.—13° 30' N., 16° 45' W.
- *Batiamponde.—Near Stanleyville, Belgian Congo.
- Bawi (I.).—6° 10' S., 39° 5' E.
- *Bazen.—Abyssinia.
- *Beach Bush.—Near Durban, Natal.
- *Bedza.—Matoppo Hills, Southern Rhodesia.
- Beira.—20° S., 35° E.
- Bela.—4° N., 41° 30' E.
- Belingwe.—20° 25' S., 30° E.
- Bembesi (R.).—19° to 20° S., 28° to 29° 30' E.
- Bena Dibebe.—4° S., 22° 45' E.
- Bena Makima.—4° 45' S., 20° 45' E.
- *Benda.—French Congo.
- Bendo.—6° 55' N., 11° 15' W.
- Bengamisa.—1° N., 25° 10' E.
- Benguela.—12° 30' S., 13° 20' E.
- Beni.—0° 20' N., 29° 40' E.
- Benin.—6° 20' N., 5° 40' E.
- Benito (R.).—1° 35' N., 9° 35' E.
- Benue (R.).—8° N., 7° to 10° E.
- Berbera.—10° 25' N., 45° 5' E.
- *Bergvliet.—Cape Flats, Cape Colony.
- Bethanien.—26° 30' S., 17° 10' E.
- *Betutu.—Maringa-Lopori District, Belgian Congo.
- Bianza.—1° 25' S., 20° 10' E.
- Bibundi.—4° 15' N., 9° E.
- Bihunga.—0° 20' N., 30° 5' E.
- Bindura.—17° 25' S., 31° 25' E.
- Bipindi.—3° 5' N., 10° 25' E.
- *Bisa Timo.—Near Harar, Abyssinia.
- Bismarekburg.—8° 5' N., 1° 20' E.
- Bismarekhügel.—3° 5' S., 37° 30' E.
- Bissao.—11° 45' N., 15° 40' W.
- Bissis (I.).—11° 45' N., 16° 5' W.
- Bizen (Mt.).—15° 20' N., 39° 10' E.
- Blantyre.—15° 45' S., 35° 5' E.
- Bloemfontein.—29° 5' S., 26° 10' E.
- Blue Cliff.—33° 30' S., 25° 25' E.
- Blue Nile (R.).—11° 15' to 15° 40' N., 32° 25' to 35° E.
- Blue Post.—1° 5' S., 37° 10' E.
- *Bobi.—Near Gali, Belgian Congo.
- *Boda.—French Congo.
- Boga.—1° N., 30° E.
- Bogos.—15° 50' N., 38° E.
- Bokakata.—1° 10' N., 19° 25' E.
- Bokala.—3° 15' S., 17° 5' E.
- Bolengi.—0° 5' S., 18° 10' E.
- Bolobo.—2° 15' S., 16° 15' E.

- Bolombo.—1° 25' N., 18° 55' E.
 Boma.—5° 50' S., 13° 10' E.
 *Boma Gombe.—German East Africa.
 Boma Sundi.—5° 20' S., 12° 50' E.
 Bombaie.—4° 50' S., 23° 35' E.
 Bomili.—1° 30' N., 27° 20' E.
 Bondei.—5° 20' S., 38° 50' E.
 Bonjongo.—4° 5' N., 9° 10' E.
 Bonny (R.).—4° 25' to 4° 45' N., 7° 10' E.
 Boran Galla, Upper Ganale.—4° 30' N., 39° 30' E.
 *Borda.—French Congo.
 *Boro (R.).—Darbanda, French Congo.
 Bothaville.—27° 20' S., 26° 35' E.
 Botuma.—0° 30' N., 19° 30' E.
 Bourka.—9° 25' N., 41° 15' E.
 Boyeka.—1° 10' N., 19° 5' E.
 Boyenge.—0° 25' N., 18° 45' E.
 Boyulu.—1° N., 27° E.
 Brazzaville.—4° 25' S., 15° 20' E.
 Buarsangueli.—10° 50' N., 48° E.
 Bububu.—6° 5' S., 39° 20' E.
 Buddu Forest.—1° S., 31° 40' E.
 Buditu.—5° 25' N., 38° 30' E.
 Bugalla (I.).—0° 30' S., 32° 15' E.
 Buiko.—4° 42' S., 38° E.
 Bujongolo.—0° 20' N., 29° 55' E.
 Bukama.—9° 15' S., 25° 40' E.
 Bukoba.—1° 30' S., 32° E.
 Bularli.—7° 55' N., 43° 30' E.
 Bulawayo.—20° 10' S., 28° 50' E.
 Bumba.—2° 10' N., 22° 30' E.
 *Bunthorne Mine.—Near Bulawayo, Rhodesia.
 Bura.—3° 30' S., 38° 18' E.
 Butiaba.—1° 50' N., 31° 30' E.
 Butiti.—0° 45' N., 30° 20' E.
 Buzubizi.—0° 20' N., 32° 5' E.
 Caconda.—13° 45' S., 15° E.
 Cairo.—30° 12' N., 31° 10' E.
 Caledon.—34° 10' S., 19° 25' E.
 Camayenne.—9° 25' N., 13° 40' W.
 Cameroon (Mt.).—4° 15' N., 9° 10' E.
 Cameroon (R.).—3° 55' N., 9° 35' E.
 Campo.—2° 20' N., 9° 50' E.
 Campo Tembo.—3° 5' S., 38° 10' E.
 *Candolo.—Belgian Congo.
 Cape Cross.—21° 45' S., 13° 55' E.
 Cape Flats.—Near Cape Town, Cape Colony.
 Cape Lopez.—0° 40' S., 8° 45' E.
 Cape Mount.—6° 45' N., 11° 25' W.
 Cape of Good Hope.—34° 15' S., 18° 30' E.
 Cape Palmas.—4° 30' N., 7° 35' W.
 Cape Town.—33° 55' S., 18° 25' E.
 Cape Verde.—14° 35' N., 17° 50' W.
 Casamance (R.).—12° 40' N., 14° to 16° 45' W.
 *Cawston Farm.—On Umgusa R., Southern Rhodesia.
 Ceres.—33° 25' S., 19° 20' E.
 Chacansengula.—0° 20' N., 31° 55' E.
 Chakamakue.—23° 45' S., 22° 5' E.
 Chake Chake, Pemba Island.—5° 15' S., 39° 45' E.
 Chama.—5° N., 1° 40' W.
 Changamwe.—4° 2' S., 39° 35' E.
 *Changmane, probably Changamwe.
 Chania (R.).—0° 40' to 1° 10' S., 36° 50' E.
 Charlestown.—27° 25' S., 29° 55' E.
 Cheik Osman.—12° 55' N., 45° E.
 Cheteni.—4° 5' S., 39° 40' E.
 Chikai.—5° 50' S., 12° 25' E.
 Chinchoxo.—5° 15' S., 12° 15' E.
 Chirinda (Mt.).—21° S., 32° 45' E.
 Chumbiri.—2° 40' S., 16° 15' E.
 *Ciama.—Southern Rhodesia.
 Coffee (Mt.).—6° 30' N., 10° 35' W.
 *Colba (R.).—Abyssinia.
 Colenso.—28° 45' S., 29° 50' E.
 *Comba Ibre.—French Congo.
 *Combra Tora.—French Congo.
 Conde.—5° 5' S., 12° 15' E.
 Congo da Lemba.—5° 40' S., 13° 40' E.
 Constantia, near Cape Town.—34° 5' S., 18° 25' E.
 Conway.—31° 45' S., 25° 15' E.
 Coquilhatville.—0° 1' N., 18° 20' E.
 Coromma.—5° 30' N., 38° E.
 Cubango (R.).—12° 30' to 20° S., 16° 15' to 22° 30' E.

*Cucala.—Portuguese West Africa.

Cuito (R.).—12° 35' to 18° S., 18° to 20° 30' E.

Dakar.—14° 40' N., 17° 35' W.

Damba (I.).—0°, 32° 50' E.

Danakil.—12° N., 42° E.

Daouele.—8° 45' N., 44° 5' E.

Darbanda.—7° 40' N., 21° 35' E.

Darassalaam.—6° 50' S., 39° 15' E.

Daua (R.).—4° to 5° 20' N., 39° 10' to 42° E.

De Aar.—30° 40' S., 24° 5' E.

Deep (R.).—34° 5' S., 18° 25' E.

*Degabolla.—Abyssinia.

Delagoa Bay.—26° S., 32° 40' E.

*Denge.—Near Niangara, Belgian Congo.

Diaghe (R.).—4° 30' N., 28° 25' E.

Dibeile.—4° S., 22° 45' E.; same as Bena Dibeile.

Dima.—3° 20' S., 17° 20' E.

Dimbroko.—6° 40' N., 4° 55' W.

Dimé.—5° 35' N., 36° 50' E.

Diré Daua.—9° 40' N., 41° 50' E.

Djebel Akhmed Aga.—11° N., 33° E.

Djebel Hakim.—9° 25' N., 42° 25' E.

Djipe (L.).—3° 35' S., 37° 45' E.

Djoungou.—9° 45' N., 1° 50' E.

Djur (R.).—8° N., 28° E.

Dolo.—4° 25' S., 15° 25' E.

*Dongola.—Eritrea.

Duala.—4° N., 9° 40' E.

*Dukudu.—Zululand.

Duma.—3° 50' N., 18° 35' E.

Dunbrody.—33° 30' S., 25° 30' E.

Dundusana.—2° 45' N., 22° 20' E.

Dungu, Mayombe.—4° 45' S., 12° 55' E.

Dungu, Uele.—3° 30' N., 28° 30' E.

Durban.—29° 50' S., 31° E.

Eala.—0° 1' N., 18° 25' E.

East London.—33° S., 27° 55' E.

Ebolowa.—2° 55' N., 11° 5' E.

*Edeloud.—Kordofan Desert, Anglo-Egyptian Sudan.

Efulen.—2° 40' N., 10° 45' E.

*Ekeneli.—Near Metit, Cameroon.

Eket.—4° 35' N., 7° 55' E.

*Elat.—Near Metit, Cameroon.

El Burgon.—0° 15' S., 35° 50' E

*El Hefera.—On the Settiti, Anglo-Egyptian Sudan.

Elisabethville.—11° 45' S., 27° 40' E.

Ellahelaj.—7° N., 49° 20' E.

Eloby (I.).—1° N., 9° 30' E.

Endessa.—8° 40' N., 40° E.

Entebbe.—0° 5' N., 32° 30' E.

*Entendweni.—Zululand.

Epulu (R.).—1° 20' N., 28° 40' E.

Erdal.—5° 30' N., 48° 45' E.

Error-es-Saghir.—9° 30' N., 44° 5' E.; same as Hargeisa.

Estcourt.—29° S., 29° 50' E.

*Esuk Ekkpo Abassi.—Eket District, Nigeria.

*Etombe.—Cameroon.

Ettoko.—4° 40' N., 39° E.

Faf Plain.—6° 30' N., 44° 10' E.

Falaba.—9° 45' N., 11° 20' W.

Faradje.—3° 40' N., 29° 40' E.

Fariala.—1° 25' N., 28° E.

Fashoda.—10° N., 32° E.

*Fello.—Senegambia.

Fernando Po.—3° 30' N., 8° 30' E.

Fikilini, near Stanleyville.

Forcados (R.).—5° 20' N., 5° 35' to 6° 20' E.

Fort Archambault.—9° 5' N., 18° 35' E.

Fort Crampel.—7° 10' N., 9° 20' E.

Fort de Possel.—5° N., 19° 15' E.

Fort Hall.—0° 50' S., 37° 15' E.

Fort Johnston.—14° 30' S., 35° 15' E.

Fort Portal.—0° 45' N., 30° 15' E.

Fort Sibut.—5° 55' N., 19° E.; same as Krebedje.

Freretown.—4° 3' S., 39° 40' E.

Fundu (I.).—5° 5' S., 39° 40' E.

Gaboon (R.).—0° 10' to 0° 30' N., 9° 20' to 10° 10' E.

*Galago (L.).—Northern Ruanda, German East Africa.

Gali.—2° 25' N., 21° 35' E.

Gamangui.—2° 10' N., 27° 20' E.

- Ganale Gudda (R.).—5° 25' to 6° N., 39° to 40° 45' E.
 Ganana (R.), see Ganale (R.).
 Ganda Sundi.—4° 50' S., 12° 50' E.
 Garamba.—4° 10' N., 29° 40' E.
 Gawieb.—22° 45' S., 15° 10' E.
 Gazi.—4° 25' S., 39° 30' E.
 Gebelein, White Nile.—12° 35' N., 32° 45' E.
 George.—33° 55' S., 22° 30' E.
 Ghinda.—15° 35' N., 39° E.
 *Ghrab el Aish.—Anglo-Egyptian Sudan.
 Giari Bule.—5° 55' N., 38° 50' E.
 Gilgil.—0° 30' S., 36° 20' E.
 *Gischin.—Southern Arabia.
 Giuba (R.).—0° 10' S. to 4° 5' N., 42° 30' E.
 *Glatkop. — Little Namaland, Cape Colony.
 Gobabis.—22° 25' S., 18° 55' E.
 *Goda.—French Congo.
 Godo Burka.—9° 5' N., 39° 25' E.
 Gogfale.—9° 50' N., 41° 20' E.
 Golah.—7° 5' N., 8° 35' W.
 Golungo Alto.—9° 5' S., 14° 55' E.
 Gomba.—4° 10' S., 14° 20' E.
 Gomod.—15° 35' N., 39° 10' E.
 Gondokoro.—4° 50' N., 31° 45' E.
 Gordon Bay.—34° 10' S., 18° 50' E.
 Gorée (I.).—14° 40' N., 17° 30' W.
 Gorongoza.—18° 30' S., 34° E.
 Gotta.—9° 35' N., 41° 20' E.
 Grahamstown.—33° 15' S., 26° 35' E.
 Grand Bassa.—5° 45' N., 10° W.
 Grand Bassam.—5° 10' N., 3° 50' W.
 Grand Batanga.—2° 50' N., 9° 55' E.
 Grand Lahou.—5° 10' N., 5° W.
 *Greymine.—Natal.
 Grootfontein.—19° 35' S., 18° 55' E.
 Gubbet.—11° N., 47° 50' E.
 *Guengera.—Pungwe Valley, Portuguese East Africa.
 Gwaai.—19° 20' S., 27° 40' E.
 Haitajwa Cave.—6° 15' S., 39° 15' E.
 Hamman's Kraal.—25° 25' S., 28° 15' E.
 Harar.—9° 25' N., 42° 25' E.
 Hargeisa.—9° 30' N., 44° 5' E.
 Hartley.—18° 10' S., 30° 30' E.
 Hauacio.—5° N., 39° E.
 Hauash (R.).—8° 25' to 11° 40' N., 38° 25' to 41° 30' E.
 Hebron.—25° 30' S., 27° 55' E.
 Herrer.—9° 40' N., 41° 20' E.
 Herschel.—30° 40' S., 27° 15' E.
 *Hička Bourka.—Abyssinia.
 Hillside, near Bulawayo.—20° 10' S., 28° 50' E.
 Himo (R.)., Mt. Kilimanjaro.—3° 10' to 3° 35' S., 37° 35' E.
 Hoima.—1° 35' N., 31° 30' E.
 Homran.—14° 25' N., 36° 10' E.
 Hope Fountain.—20° 20' S., 28° 55' E.
 Hopetown.—29° 35' S., 24° 5' E.
 Ibadan.—7° 25' N., 3° 55' E.
 Ibaka.—1° 35' S., 16° 40' E.
 Ibali.—2° S., 18° 10' E.
 Ibanda.—0° 20' N., 30° 5' E.
 Ibo.—12° 25' S., 40° 35' E.
 Ihulu (R.), same as Epulu (R.).
 Ikela.—1° S., 23° 10' E.
 Ikelemba (R.).—0° 20' N., 18° 15' to 20° 20' E.
 Ikenge.—0° 5' S., 18° 35' E.
 *Imbokro.—Ivory Coast. Probably misspelling of Dimbroko.
 *Ingfal.—Abyssinia.
 Inhambane.—23° 55' S., 35° 35' E.
 *Injolo. — Equator District, Belgian Congo.
 Inkisi (R.).—4° 40' to 6° 15' S., 15° to 15° 35' E.
 Inongo.—1° 55' S., 18° 20' E.
 Irebu.—0° 35' S., 17° 50' E.
 *Iringui.—Belgian Congo.
 Irumu.—1° 20' N., 30° E.
 *Isalinio. — Mpororo, German East Africa.
 Isangi.—0° 50' N., 24° 15' E.
 Isipingo.—30° S., 30° 50' E.
 Issawe.—2° 35' S., 29° 45' E.
 Itigi.—5° 45' S., 34° 30' E.
 Ituri (R.).—1° 30' N., 26° to 30° E.
 Jabassi.—4° 30' N., 9° 55' E.
 Jacquenville.—5° 15' N., 4° 25' W.

- Johann-Albrechtshöhe.—4° 40' N., 9° 25' E.
 Johannesburg.—26° 10' S., 28° E.
 Junk (R.).—6° 5' to 6° 30' N., 9° 35' to 10° 25' W.
 Kabambare.—4° 40' S., 27° 45' E.
 Kabanza, Lovoi R.—8° 15' S., 26° 30' E.
 Kabare.—0° 35' S., 29° 30' E.
 Kabinda.—6° 10' S., 24° 20' E.
 Kabwe (L.).—9° 20' S., 25° 50' E.
 Kadjura, Unyoro.—1° 40' N., 31° 20' E.
 *Kafogo.—Limba, Sierra Leone.
 Kagera (R.).—1° to 2° 30' S., 30° to 31° 50' E.
 Kahe, Mt. Kilimanjaro.—3° 30' S., 37° 25' E.
 *Kaibo.—Uganda.
 Kairouan.—35° 40' N., 10° 5' E.
 *Kaka.—Schioa, Abyssinia.
 Kaka, White Nile.—10° 40' N., 32° 15' E.
 *Kakir.—Kalahari.
 Kakoulina (Mt.).—9° 40' N., 13° 25' W.
 *Kalanga.—On the Upper Lukuga, Belgian Congo.
 *Kalumba.—Belgian Congo. Probably Kalumbi.
 Kalumbi.—5° 50' S., 28° 35' E.
 Kamaggas.—29° 45' S., 17° 25' E.
 Kambove.—10° 50' S., 26° 35' E.
 Kamiti (R.).—1° 5' S., 36° 50' E.
 Kampala.—0° 20' N., 32° 20' E.
 Kana.—8° 25' S., 26° 20' E.
 Kandahar (I.), Zambesi R.—17° 55' S., 25° 35' E.
 Kang.—23° 40' S., 22° 45' E.
 Kapema.—10° 40' S., 28° 20' E.
 *Kapinga.—Belgian Congo.
 Kapiri.—10° 15' S., 26° 20' E.
 Karagwe.—1° to 2° S., 31° E.
 Karemi.—0° 5' N., 29° 40' E.
 Karisimbi (Mt.).—1° 30' S., 29° 25' E.
 Karoli (Mt.).—3° 55' N., 37° 30' E.
 Karssa.—9° 25' N., 41° 35' E.
 Kasai (R.).—3° to 10° S., 16° to 22° E.
 Kasenga.—10° 15' S., 28° 45' E.
 Kasengui.—0° 20' N., 31° 40' E.
 Kasindi.—0°, 29° 40' E.
 Kasongo.—4° 20' S., 26° 25' E.
 Kasonsero.—1° N., 30° 10' E.
 Kassala.—15° 25' N., 36° 25' E.
 Kassenje.—1° 25' N., 30° 30' E.
 Katakai, Katanga.—5° 45' S., 29° E.
 Katala.—6° S., 12° 45' E.
 Katende.—0° 15' N., 32° 25' E.
 *Katumba.—Belgian Congo.
 Kavirondo Bay.—0° 20' S., 34° 30' E.
 Kawa, White Nile.—13° 45' N., 32° 30' E.
 Kazungula.—17° 50' S., 25° 5' E.
 Kenia (Mt.).—0° 20' S., 37° 25' E.
 Kentani.—32° 30' S., 28° 20' E.
 Keren.—15° 45' N., 38° 30' E.
 Kerreri.—15° 45' N., 32° 30' E.
 Kgokong.—24° 15' S., 23° E.
 Khakhea.—24° 40' S., 23° 25' E.
 Khami (R.).—20° 15' S., 28° E.
 Khartum.—15° 40' N., 32° 35' E.
 Khutu Steppe.—7° 30' S., 38° E.
 Kiambi.—7° 20' S., 27° 55' E.
 Kibombo.—4° S., 26° E.
 Kibonoto, Mt. Kilimanjaro.—3° 15' S., 37° 10' E.
 Kibosho.—3° 15' S., 37° 20' E.
 Kibwezi.—2° 25' S., 37° 55' E.
 Kifumbiro.—1° 15' S., 31° 25' E.
 Kigali.—2° S., 30° E.
 *Kigerama.—Near Kagera R., German East Africa.
 Kihenga.—5° 45' S., 37° 35' E.
 Kijabe.—0° 55' S., 36° 35' E.
 Kika.—9° 15' N., 2° 45' E.
 Kikondja.—8° 10' S., 26° 25' E.
 Kikuyu.—1° 15' S., 36° 45' E.
 Kikwit.—5° 25' S., 18° 50' E.
 Kilema.—3° 20' S., 37° 30' E.
 Kilimanjaro (Mt.).—3° S., 37° 20' E.
 Kilindini.—4° S., 39° 45' E.
 Kilo.—1° 55' N., 30° E.
 *Kilongalunga.—Near St. Gabriel, Belgian Congo.
 *Kimaha.—Gold Coast.
 Kimberley.—28° 45' S., 24° 45' E.
 Kimpoko.—4° 10' S., 15° 40' E.
 Kimuenza.—4° 25' S., 15° 20' E.

- Kinangop.—0° 35' S., 36° 30' E.
 Kindia.—10° N., 12° 45' W.
 Kindu.—3° S., 26° E.
 King William's Town.—32° 50' S., 27° 25' E.
 Kiniati.—5° 20' S., 13° E.
 Kinshasa.—4° 20' S., 15° 20' E.
 Kipaila.—10° 45' S., 28° 35' E.
 Kirstenbosch.—34° S., 18° 30' E.
 Kirundu.—1° 15' S., 25° 30' E.
 Kisantu.—5° 10' S., 15° 10' E.
 Kisumu.—0° 5' S., 34° 45' E.
 *Kitagweta.—Uganda. Probably misspelling for Kitagwenda.
 Kitagwenda.—0° 10' N., 30° 45' E.
 Kitempuka.—5° 25' S., 28° 45' E.
 Kitobola.—5° 20' S., 14° 40' E.
 Kitta.—5° 55' N., 0° 55' E.
 Kitui.—1° 15' S., 38° E.
 Kivu (L.).—2° S., 29° E.
 Kynsna.—34° S., 23° 5' E.
 Koloka, near Angu.—3° 5' N., 24° 35' E.
 Konakry.—9° 25' N., 13° 45' W.
 Kondué.—4° 55' S., 23° 15' E.
 Kooa.—24° 50' S., 24° 25' E.
 Kor Attar.—0° 55' to 10° 25' N., 32° 15' to 34° E.
 Kordofan.—12° 25' N., 31° 15' E.
 Kor Gasch.—15° 5' to 17° N., 35° to 36° 25' E.
 *Kor Guillo.—Homran, Anglo-Egyptian Sudan.
 Kor Langhebb.—17° 25' N., 37° E.
 Kor Lebka.—16° 5' to 16° 20' N., 38° 30' to 39° 15' E.
 Korogwe.—5° 10' S., 38° 30' E.
 *Kortright Hill.—Sierra Leone.
 Kotonou.—6° 20' N., 2° 25' E.
 Kouandé.—10° 20' N., 1° 45' E.
 Kounhi.—9° 15' N., 41° 5' E.
 Krebedje.—5° 55' N., 19° E.
 Kribi.—3° N., 10° E.
 Kroonstad.—27° 40' S., 27° 20' E.
 Kubub.—26° 45' S., 16° 15' E.
 Kuilu (R.).—3° 30' to 4° 30' S., 11° 45' to 12° 30' E.
 Kulumuzi (R.).—5° 5' S., 38° 50' to 39° 5' E.
 Kunga, near Malela.—5° 55' S., 12° 35' E.
 Kwamouth.—3° 20' S., 16° 10' E.
 Kwango (R.).—3° to 10° S., 17° to 19° E.
 Kwesi.—1° 5' N., 30° E.
 Kwidjwi (I.), Lake Kivu.—2° 10' S., 29° 20' E.
 *Kyaka Fort.—On Kagera R., German East Africa.
 Ladismith.—33° 30' S., 21° 20' E.
 Lado.—5° 5' N., 31° 45' E.
 Ladysmith.—28° 30' S., 29° 45' E.
 Laffarugh.—10° 5' N., 44° 50' E.
 Lagos.—6° 30' N., 3° 25' E.
 Lambarene.—0° 40' S., 10° 15' E.
 Lamu (I.).—2° 15' S., 40° 55' E.
 Landana.—5° 15' S., 12° 15' E.
 Landjoro.—3° 25' S., 37° 50' E.
 Langenburg.—9° 35' S., 34° 10' E.
 *Las Ej.—Somaliland.
 *Leboi.—Abyssinia.
 Lehututu.—23° 55' S., 21° 55' E.
 Leitokitok.—2° 55' S., 37° 45' E.
 Lemba (R.), Mayombe.—5° 20' to 5° 40' S., 12° 30' to 12° 45' E.
 Leopold II (L.).—2° S., 18° 15' E.
 Leopoldville.—4° 25' S., 15° 20' E.
 Leribe.—28° 45' S., 28° 15' E.
 Lesse.—0° 40' N., 29° 40' E.
 Lessouto, see Basutoland.
 Letlake.—23° 40' S., 22° 20' E.
 Let Marefia.—9° 45' N., 40° E.
 Lettema (Mt.).—3° 40' S., 37° 20' E.
 *Lewa Mambaa.—German East Africa.
 Libenge.—3° 35' N., 18° 30' E.
 Libreville.—0° 25' N., 9° 25' E.
 Lie.—2° N., 21° 20' E.
 Lifungula, Uele.—3° 45' N., 27° 20' E.
 Likimi.—2° 40' N., 20° 40' E.
 Likoni.—4° 5' S., 39° 40' E.
 Limba.—9° 25' N., 12° W.
 Lindi.—10° S., 39° 45' E.
 Lindi (R.).—1° 25' N., to 0° 25' S., 25° 5' to 29° E.
 Lingunda.—1° N., 20° 40' E.
 *Linkanda.—Kasai, Belgian Congo.
 Lisala.—2° 10' N., 21° 30' E.

- Lisasa.—0° 15' N., 29° 30' E.
 Livingstone.—18° S., 25° 45' E.
 Loango.—4° 35' S., 11° 45' E.
 Loangwa (R.).—11° to 16° S., 30° to 33° E.
 Lobatsi.—25° 10' S., 25° 40' E.
 Lobay (R.).—4° 15' S., 18° E.
 *Lobombo Borges.—Portuguese East Africa.
 Lokelenge.—1° 20' N., 22° 45' E.
 Lokundje (R.).—3° 5' to 3° 25' N., 10° to 11° 15' E.
 Lolodorf.—3° 15' N., 10° 40' E.
 Lomami (R.).—0° 45' N. to 8° 45' S., 24° to 26° E.
 Lomie.—3° 10' N., 13° 40' E.
 Lonely Mine.—19° 25' S., 29° 25' E.
 *Longa.—Senegambia. Probably misspelling for Louga.
 Longji.—3° 5' N., 10° E.
 Longonot Crater.—0° 55' S., 36° 25' E.
 Lonkala.—4° 40' S., 23° 10' E.
 Los (Is.).—9° 30' N., 13° 50' E.
 Louga.—15° 35' N., 16° 25' W.
 Lovoi (R.).—8° 10' S., 26° 20' E.
 Luahi.—5° S., 12° 25' E.
 Luapula (R.).—9° to 12° S., 29° E.
 Lubi (R.).—5° to 6° 40' S., 23° 20' E.
 Lubila.—1° N., 27° 10' E.
 Lubumbashi (R.).—11° 45' S., 27° 40' E.
 Lubutu.—0° 40' S., 26° 40' E.
 Lüderitz Bay.—26° 40' S., 15° 10' E.; same as Angra Pequena.
 Lugh.—3° 50' N., 42° 35' E.
 *Lugny.—French Congo.
 Lugombe.—5° 40' S., 28° 50' E.
 Lukaya (R.).—4° 30' S., 15° 20' E.
 Luki.—5° 35' S., 13° 10' E.
 Lukolela.—1° 10' S., 17° 10' E.
 Lukonzolwa.—8° 50' S., 28° 40' E.
 Lukuga (R.).—6° S., 27° to 29° E.
 Lukula.—5° 25' S., 13° E.
 Lukungu (R.).—5° to 5° 30' S., 14° 15' to 14° 45' E.
 *Lumaliza.—Belgian Congo.
 *Lumbulumbu.—Belgian Congo.
 Lumbwa.—0° 40' S., 35° 20' E.
 Lungube.—Katanga. Probably misspelling for Lugombe.
 Lusambo.—4° 55' S., 23° 15' E.
 Lusinga (I.), see Rusinga (I.).
 Lydenburg.—25° 5' S., 30° 25' E.
 Maddo Wells.—3° 55' N., 41° E.
 Madibi.—4° 25' S., 18° 35' E.
 Madingu.—4° 10' S., 13° 30' E.
 *Madona.—Nyamyas, Nyasaland.
 Mafeking.—25° 50' S., 25° 40' E.
 Mafia (I.).—7° 45' S., 39° 30' E.
 *Mafungu.—Belgian Congo.
 Magalapye.—23° 5' S., 26° 55' E.
 Magala Re Umberto.—4° 45' N., 41° 25' E.
 *Magnarra (R.). — Portuguese East Africa.
 Maji-ya-chumvi.—3° 50' S., 39° 25' E.
 *Majuba. — Herschel District, Cape Colony.
 *Majuba Neck.—Cape Colony.
 *Majunga.—Belgian Congo.
 Makalla.—14° 30' N., 49° 10' E.
 Makalle.—13° 30' N., 39° 25' E.
 Makanga.—3° 40' S., 25° 50' E.
 Makapan.—25° 15' S., 28° 5' E.
 Makdischu.—2° N., 45° 30' E.
 Malange.—9° 35' S., 16° 25' E.
 Malela.—6° S., 12° 40' E.
 Malema.—2° N., 21° 30' E.
 Malende.—4° 35' N., 9° 30' E.
 *Malindi.—German East Africa.
 Malindi (Southern Rhodesia).—19° S., 27° 10' E.
 *Malome.—On the Okiavo R., Belgian Congo.
 Malvern.—29° 50' S., 31° E.
 Mamou.—10° 20' N., 12° 15' W.
 Manamama.—2° N., 28° E.
 Manda (I.).—2° 20' S., 40° 55' E.
 Mandimbo.—0° 55' S., 27° 20' E.
 *Mandouga.—French Congo.
 Mandungu.—2° 30' N., 23° 15' E.
 Mangapwani.—6° S., 39° 10' E.
 Mangbetu Country, see Niangara.
 Mange.—4° S., 19° 30' E.

- *Maniou.—French Guinea. Probably Mamou.
- Manow.—9° 15' S., 33° 45' E.
- Marangu.—3° 20' S., 37° 30' E.
- Masai Steppe.—1° S., 35° E.
- Masaki.—1° 5' S., 28° 10' E.
- Mascot.—23° 20' N., 58° 40' E.
- Maseru.—29° 20' S., 27° 25' E.
- Masisi.—1° S., 28° 30' E.
- Masongo.—1° S., 27° 35' E.
- Massaua.—15° 40' N., 39° 25' E.
- Matadi.—5° 50' S., 13° 35' E.
- Matagoi.—2° 30' N., 43° 25' E.
- *Matchacha.—Between Ponthierville and Nyangwe, Belgian Congo.
- Matetsi.—18° 15' S., 26° 5' E.
- Matjesfontein.—23° 15' S., 20° 35' E.
- Matoppo Hills.—20° 45' S., 28° 50' E.
- Matroosberg.—33° 20' S., 19° 40' E.
- Mau Escarpment.—0° to 1° S., 35° 30' to 36° 20' E.
- *Maveli (Mts.).—Near the Munda R., French Congo.
- Mawambi.—1° 10' N., 28° 45' E.
- *Mayabal.—Eritrea.
- Mayombe.—5° S., 13° E.
- M' Bale.—1° 5' N., 34° 10' E.
- Mbalmajo.—3° 30' N., 11° 35' E.
- Mbamu.—4° 15' S., 14° 45' E.
- Mbaramu.—4° 25' S., 38° 20' E.
- *M'Bounion.—French Congo.
- Mbusini.—6° 15' S., 38° E.
- Mbuyuni.—3° 30' S., 37° 55' E.
- Medje.—2° 25' N., 27° 30' E.
- *Melelia.—Eritrea.
- Merca.—1° 45' N., 44° 55' E.
- Meru (Mt.).—3° 15' S., 36° 45' E.
- Metemma.—16° 45' N., 33° 20' E.
- Metit.—3° 45' N., 11° 35' E.
- *Mfongosi.—Zululand.
- Mful Aja.—Near Metit, Cameroon.
- *Mgaturu.—On Kwidjiwi Island.
- Mhonda.—6° 5' S., 37° 35' E.
- Milmil.—8° 20' N., 43° 50' E.
- *Mindouga.—French Congo.
- Mindouli.—4° 15' S., 14° 15' E.
- Misahöhe.—6° 55' N., 0° 35' E.
- Mitiana.—0° 25' N., 32° 5' E.
- *Mkosi.—Zululand.
- Mobuku (R.).—0° 20' N., 30° 15' E.
- Mobwasa.—2° 40' N., 23° 5' E.
- Moera.—0° 35' N., 29° 30' E.
- Moero (L.).—9° S., 29° E.
- Mohasi (L.).—1° 50' S., 30° 20' E.
- *Mokbe.—Near Dume, Cameroon.
- *Mokundange.—Cameroon.
- Moliwe.—4° 5' N., 9° 15' E.
- Molo.—0° 15' S., 35° 45' E.
- Molundu.—2° 5' N., 15° 15' E.
- Mombasa.—4° S., 39° 50' E.
- Mombo.—4° 55' S., 38° 15' E.
- Mondombe.—0° 35' S., 22° 30' E.
- *Mondu (R.).—Southern Rhodesia.
- Monga.—Near Amani, German East Africa.
- Mongalla.—5° 10' N., 31° 45' E.
- Mongumba.—3° 35' N., 18° 30' E.
- Monrovia.—6° 30' N., 10° 50' W.
- Montagu Pass.—33° 45' S., 22° 30' E.
- *Moor Plantation. — Near Ibadan, Nigeria.
- Morogoro.—6° 50' S., 37° 50' E.
- *Morumballe (Mts.). — Zambesi R., Portuguese East Africa.
- Moschi.—3° 20' S., 37° 25' E.
- Mosekowa.—0° 45' S., 26° 55' E.
- Mossamedes.—15° 15' S., 12° 15' E.
- *Motombe.—Okavango R., Belgian Congo.
- *Mountain Rise.—Near Pietermaritzburg, Natal.
- *Mowange.—Cameroon.
- Mozambique (I.).—15° S., 40° 40' E.
- *M'Piaka.—French Congo.
- M'Pila.—4° 10' S., 15° 20' E.
- Mpongwe District.—0° 15' N., 9° 20' E. and 0° 40' S., 9° 20' E.
- Mpororo.—0° 40' to 1° 25' S., 29° 20' to 30° 30' E.
- Msozi.—0° 55' S., 31° 50' E.
- *Mto-ya-Kifaru.—Arusha-chini, German East Africa.
- Mubende Region.—1° 15' N., 31° 15' E.
- Mubuku, eastern Ruwenzori Mt., see Mobuku.
- Muculla.—7° S., 12° 50' E.
- Muamba.—4° 10' S., 37° 50' E.

- Mukenge.—6° S., 22° 30' E.
 Mukonje (Farm).—4° 35' N., 9° 30' E.
 Munda (R.).—0° 25' N., 9° 35' to 10° E.
 Mundame.—4° 35' N., 9° 35' E.
 Mungo (R.).—4° to 4° 50' N., 9° 30' E.
 Munza.—3° 20' N., 28° E.
 Mussengue, see Sobato.
 Mwatate.—3° 30' S., 38° 20' E.
 Mwengwa.—13° S., 27° 40' E.
 Mwera (R.), Zanzibar.—6° 10' S., 39° 15' E.
- Nabambisso (R.).—4° 45' N., 28° 40' E.
- *Nabena.—Liberia.
 Nahoon (R.).—32° 55' S., 27° 50' E.
 Nairobi.—1° 5' S., 36° 50' E.
 Naivasha.—0° 40' S., 36° 35' E.
 Nakitawa.—0° 25' N., 30° E.
 Nakuru.—0° 15' S., 36° 10' E.
 Nala.—2° 50' N., 27° 50' E.
 Namaland, Little.—28° 45' to 30° 15' S., 16° 45' to 19° E.
 Nanga (Mts.).—4° 5' N., 13° 5' E.
 Nana (R.).—6° 35' N., 19° E.
 Naremuru (R.).—0° 10' S., 37° 15' E.
 Natron Lakes, foot of Mt. Kilimanjaro.—3° 10' S., 36° 55' E.
 N'daika.—0° 10' S., 37° 10' E.
 Ndara.—3° 30' S., 38° 30' E.
 Ndjoro.—0° 20' S., 36° E.
- *N'Douna.—French Congo.
 Nduye.—1° 20' N., 28° 30' E.
 Nefasit.—15° 25' N., 39° E.
 Newcastle.—27° 45' S., 29° 50' E.
 New Hanover.—29° 20' S., 30° 35' E.
 New Moschi.—3° 20' S., 37° 20' E.
 Ngamba (I.).—0° 5' S., 32° 40' E.
 Ngami (L.).—20° 50' S., 23° E.
 Ngare-na-Nyuki (R.), Mt. Meru.—3° to 3° 15' S., 36° 55' E.
 Ngare Nyuki (R.), Mt. Kenia.—0° 30' N. to 0° 10' S., 37° to 37° 25' E.
 Ngare Rongai.—2° 50' S., 37° 30' E.
 Ngayu.—1° 40' N., 27° 40' E.
 Ngazi.—0° 55' N., 24° 50' E.
 Nguelo.—5° 5' S., 38° 40' E.
 Ngumba Country.—3° 15' N., 10° 53' E.
- Niam Niam Country.—4° 30' to 7° N., 27° 30' to 29° 30' E.
 Niangara.—3° 40' N., 27° 50' E.
 Niapu.—2° 15' N., 26° 50' E.
 Niembo, see Pale.
 Nieuwdorp.—12° 5' S., 27° 50' E.
 Njussi.—5° 12' S., 38° 35' E.
 Nkolebunde.—3° 5' N., 10° 35' E.
 Nkolentanga.—1° 50' N., 10° 50' E.
 No (L.).—0° 30' N., 30° 35' E.
 Nord Hook.—34° 5' S., 18° 20' E.
 Nouvelle Anvers.—1° 40' N., 19° 10' E.
 Nssanakang.—5° 50' N., 8° 55' E.
 Nun (R.).—4° 20' N., 6° 5' E.
 Nyamandhloru.—19° 55' S., 28° 5' E.
- *Nyamyas.—Nyasaland.
 Nyangnori.—0°, 35° E.
 Nyangwe.—4° 15' S., 26° 15' E.
- *Nyanza (L.).—Katanga, Belgian Congo.
 Nyere, see Nyeri.
 Nyeri.—0° 30' S., 37° E.
 Nyiro (Mt.).—2° 40' N., 36° 25' E.
- Oban.—5° 20' N., 8° 35' E.
 Obbia.—5° 20' N., 48° 25' E.
 Obock.—12° N., 43° 15' E.
 Ogaden.—7° 10' N., 44° 15' E.
 Ogowe (R.).—1° S., 10° E.
 Okahandja.—22° S., 16° 55' E.
 Okiavo (R.), same as Lindi (R.).
 Old Calabar.—5° N., 8° E.
 Old Kasongo.—4° 30' S., 26° 35' E.
 Olukemeji.—7° 25' N., 3° 30' E.
- *Olombo.—Belgian Congo.
 Omaheke.—21° to 21° 30' S., 17° 30' to 20° E.
 Omdurman.—15° 50' N., 32° 5' E.
 Ondoa.—2° 55' N., 10° 15' E.
 Oni Camp.—6° 35' N., 4° 15' E.
- *Ontys.—Ex-German Southwest Africa.
 Oso (R.).—1° S., 27° 20' E.
 Ossidinge.—5° 45' N., 9° 15' E.
 Ouossou.—6° 20' N., 4° 55' W.
- *Paku.—Uele, Belgian Congo.
 Pale (Niembo).—1° S., 27° 25' E.
 Panga.—1° 45' N., 26° 15' E.
 Pangani.—5° 25' S., 39° E.

- Park Rynie.—30° 20' S., 30° 45' E.
- *Patalonga.—Near Yambuya, Belgian Congo.
- Patta (I.).—2° 5' S., 41° 5' E.
- Pawa.—2° 25' N., 27° 50' E.
- Pemba (I.).—4° 50' to 5° 30' S., 39° 40' to 39° 50' E.
- Penge.—1° 25' N., 28° 15' E.
- Pietermaritzburg.—29° 35' S., 30° 25' E.
- Pinetown.—29° 45' S., 30° 50' E.
- Pirie Forest.—32° 45' S., 27° 15' E.
- Plumtree.—20° 30' S., 27° 35' E.
- Poko.—3° 10' N., 26° 50' E.
- Ponthierville.—0° 25' S., 25° 30' E.
- Popokabaka.—5° 40' S., 17° E.
- Pori.—3° 35' S., 38° 10' E.
- Port Alfred.—33° 35' S., 26° 55' E.
- Port Elizabeth.—34° S., 25° 35' E.
- Port Florence.—0° 5' S., 34° 45' E.
- Port Natal, see Durban.
- Port Nolloth.—29° 15' S., 16° 55' E.
- Port Sudan.—19° 35' N., 37° 5' E.
- Possession (I.).—27° S., 15° 10' E.
- Pretoria.—25° 40' S., 28° 15' E.
- Prince of Wales Bay.—27° 5' S., 15° 15' E.
- Pungwe Valley.—18° 45' to 19° 50' S., 33° to 34° 55' E.
- *Quibanga.—Golungo Alto, Portuguese West Africa.
- Quitangondo.—8° 45' S., 13° 20' E.
- Quilimane.—17° 45' S., 37° E.
- Ramisi (R.).—4° 20' to 4° 30' S., 39° 5' to 39° 25' E.
- Ras Doumeira.—12° 40' N., 43° 5' E.
- Ras Fartak.—15° 50' N., 52° 30' E.
- Ras Shoab.—12° 35' N., 54° E.
- Rau (R.).—3° 10' to 3° 35' S., 37° 20' to 37° 25' E.
- *Redbank.—Rhodesia.
- Reddersburg.—29° 35' S., 26° 10' E.
- Redjaf.—4° 45' N., 31° 35' E.
- Rendile.—2° 20' N., 38° E.
- Renk.—11° 45' N., 32° 45' E.
- Revoue (R.).—18° 50' to 19° 50' S., 32° 40' to 33° 50' E.
- Richmond.—29° 50' S., 30° 15' E.
- Rikatla.—25° 45' S., 32° 35' E.
- Rio Muni.—1° 40' N., 10° E.
- Risimu.—1° N., 26° 45' E.
- Romée.—0° 35' N., 24° 50' E.
- Rooibank.—23° 10' S., 14° 40' E.
- Rosako.—6° 25' S., 38° 40' E.
- Ruanda.—2° S., 29° 50' E.
- Rudolf (L.).—2° 20' to 5° N., 36° E.
- Rungu.—3° N., 28° E.
- Rurunga.—0° 40' S., 37° 5' E.
- Rusinga (I.).—0° 25' S., 34° 10' E.
- Rutshuru.—1° 15' S., 29° 30' E.
- Ruwenzori (Mt.).—0° 30' N., 29° 50' E.
- Saati.—15° 35' N., 39° 15' E.
- Sabderat.—15° 30' N., 36° 40' E.
- Sabuka.—4° 30' S., 15° 10' E.
- Sachsenwald, see Daressalam.
- Saganeiti.—15° 5' N., 39° 10' E.
- St. Gabriel, near Stanleyville.
- Saint Louis.—16° N., 16° 35' W.
- St. Paul de Loanda.—8° 55' S., 13° 10' E.
- Saldanha Bay.—33° S., 18° E.
- Salem.—22° 40' S., 15° 25' E.
- Salisbury.—17° 45' S., 31° E.
- *Salmoreto.—Somaliland.
- Salole.—4° 15' N., 39° 25' E.
- Samburu.—3° 40' S., 38° 55' E.
- Same.—4° 5' S., 37° 45' E.
- Samkita.—0° 25' S., 10° 25' E.
- *Samlia Falls.—On the N'Gamie R., Sierra Leone?
- Sampwe.—9° 30' S., 27° 25' E.
- San Antonio.—6° 10' S., 12° 20' E.
- Sancurar.—4° N., 40° E.
- Sanda, Lower Congo.—4° 35' S., 15° 35' E.
- *Sanda.—Uganda.
- Sankisia.—9° 30' S., 25° 55' E.
- Sankuru (R.).—4° to 10° S., 21° to 22° E.
- San Pedro.—4° 45' N., 6° 40' W.
- Santa Isabel.—3° 45' N., 8° 45' E.
- San Thomé (I.).—0° 20' N., 6° 43' E.
- *Sawmills.—Umgusa R., Southern Rhodesia.

- Schoa.—10° N., 39° 30' E.
 Sciotel.—15° 35' N., 38° 20' E.
 Sebakwe.—19° 10' S., 30° E.
 Sekgoma.—24° 30' S., 23° 50' E.
 Sele (R.).—4° 20' to 5° 40' S.; 15° 35' to 15° 55' E.
 Semliki (R.).—0° to 1° N., 29° 30' to 30° E.
 *Senge (Mts.).—Cameroon.
 Sennar.—13° 30' N., 33° 35' E.
 Serengeti, British East Africa.—3° 15' S., 38° 10' E.
 Serui (R.).—22° 20' S., 26° 45' to 28° E.
 Sette-Cama.—2° 30' S., 9° 45' E.
 Settiti (R.).—14° 20' N., 35° 50' to 37° 20' E.
 Severelela.—24° 55' S., 24° 55' E.
 *Shanks Station.—Cape Colony.
 Sherbro (I.).—7° 25' N., 12° 45' W.
 Shiloango (R.).—5° S., 12° to 13° E.
 Shiloh.—19° 45' S., 28° 40' E.
 Shilouvane.—24° 10' S., 30° 15' E.
 Shimoni.—4° 35' S., 39° 15' E.
 Shinsenda.—12° 25' S., 28° E.
 *Shivyre.—Natal.
 Shoshong Road Station.—23° 35' S., 26° 35' E.
 Sibange, Maveli Mts., near the Munda R.—0° 25' N., 9° 35' E.
 Sibayi (L.).—27° 20' S., 32° 45' E.
 Sikasso.—11° 15' N., 5° 35' W.
 Simon's Town.—34° 10' S., 18° 25' E.
 Sinadogo.—5° 30' N., 46° 10' E.
 Sinai Peninsula.—28° to 30° N., 32° 30' to 35° E.
 *Sipapoma.—On the Umgusa R., Southern Rhodesia.
 Sir Lowry Pass.—34° 10' S., 18° 55' E.
 Sitaweza.—1° S., 27° 20' E.
 Smithwinkle Bay.—34° 15' S., 18° 25' E.
 *Sobato de Mussengue.—Golungo Alto, Portuguese West Africa.
 Sogodas.—14° 55' N., 36° 50' E.
 Sokode Basan.—9° N., 1° 10' E.
 Somabula.—19° 35' S., 30° 40' E.
 Songalong.—6° 25' N., 11° 15' E.
 Soppo.—4° 10' N., 9° 15' E.
 *Springvale.—Rhodesia.
 *Stamford Hill.—Near Durban, Natal.
 Stanleyville.—0° 30' N., 25° 15' E.
 *Steekstown.—Cape Colony.
 Steinkopf.—29° 15' S., 17° 45' E.
 *Stella Bush.—Durban, Natal.
 Stellenbosch.—33° 55' S., 18° 50' E.
 Stephanie (L.).—4° 40' N., 36° 50' E.
 Steynsburg.—31° 20' S., 25° 50' E.
 Stormberg Junction.—31° 20' S., 26° 20' E.
 Stormsvlei.—34° 5' S., 20° 5' E.
 Suakin.—19° 10' N., 37° 22' E.
 Sunday River Mts., near Port Elizabeth.—33° 40' S., 25° 45' E.
 *Sweetwaters.—Near Pietermaritzburg, Natal.
 Swellendam.—34° 5' S., 20° 30' E.
 *Sydenham.—Durban, Natal.
 Table (Mt.).—34° S., 18° 25' E.
 Tabora.—5° 10' S., 32° 50' E.
 Talavanje.—5° 40' S., 12° 35' E.
 *Tanda.—Usambara, German East Africa.
 Tanga.—5° 5' S., 39° 5' E.
 Tanganyika (L.).—3° to 9° S., 29° to 31° E.
 Taufkia.—9° 25' N., 31° 45' E.
 Taveta.—3° 25' S., 37° 45' E.
 Tchafianani.—8° 40' N., 40° 25' E.
 Tchercher (L.).—8° 50' N., 40° 35' E.
 Techeckna.—11° 25' N., 16° 10' E.
 Teneriffe.—28° 35' to 28° 5' N., 23° 20' to 24° 20' E.
 *Terwidja.—Northeast Africa.
 Tes.—13° 35' N., 44° E.
 Tete, Belgian Congo.—1° 20' N., 28° 50' E.
 Tete, Portuguese East Africa.—16° 10' S., 33° 30' E.
 The Bluff, Durban.—29° 55' S., 31° 5' E.
 Thiès.—14° 45' N., 16° 15' W.
 Thopane.—24° 45' S., 24° 15' E.
 Thysville.—5° 30' S., 15° E.
 Tibati.—6° 35' N., 12° 35' E.
 Tiko.—4° 5' N., 9° 25' E.
 Tiwi.—4° 5' S., 39° 35' E.

- Toro, same as Fort Portal.
- *Toukola.—Sudan.
- Toulo.—9° 20' N., 41° 10' E.
- Tsana (L.).—12° N., 37° 40' E.
- Tsavo.—3° S., 38° 25' E.
- *Tsessebe Station.—Rhodesia.
- Tshoa.—5° 35' S., 12° 45' E.
- Tshopo (R.).—0° 55' N. to 0° 5' S., 25° 5' to 28° E.
- Tua.—3° 30' S., 16° 40' E.
- Tulbagh.—33° 20' S., 19° E.
- Tumba.—5° 25' S., 14° 35' E.
- Ubangi (R.).—0° to 5° N., 18° to 23° E.
- Uchibango.—0° 40' S., 26° 20' E.
- Uebi, see Webi.
- Uele (R.).—3° 30' N., 23° to 30° E.
- Uelleburg.—1° 45' N., 10° 35' E.
- Ufumbiro (Mts.).—1° 15' S., 29° 30' E.
- *Ugono (Mt.).—German East Africa.
- Uitenhage.—33° 45' S., 25° 25' E.
- Ulenge (I.).—5° 1' S., 39° 10' E.
- Uluguru (Mts.).—7° 5' S., 37° 40' E.
- Umangi.—2° 10' N., 21° 25' E.
- *Umbilo.—Near Durban, Natal.
- Umfolosi.—28° 25' S., 32° 15' E.
- Umgeni (R.).—29° 25' to 29° 50' S., 29° 50' to 31° E.
- Umgusa (R.).—19° 25' to 20° 5' S., 27° 35' to 28° 25' E.
- Unyoro Province.—1° 50' N., 31° 35' E.
- *Uomber.—Abyssinia.
- Uorandi.—5° 45' N., 47° 15' E.
- Usambara.—5° 50' S., 38° 40' E.
- *Usegua.—6° S., 38° E.
- *Uzaga.—Region of the Great Lakes, Belgian Congo.
- Valdezia.—23° 10' S., 30° 15' E.
- Vankerckhovenville.—3° 20' N., 29° 20' E.
- Van Reenen Pass.—28° 25' S., 29° 30' E.
- Verulam.—29° 35' S., 31° E.
- Victoria.—4° N., 9° 15' E.
- Victoria Falls.—18° S., 25° 50' E.
- Vivi.—5° 45' S., 13° 35' E.
- Voi.—3° 30' S., 38° 30' E.
- Vrijburg.—26° 55' S., 24° 35' E.
- Waboniland.—2° 10' S., 40° 50' E.
- Wadi Halfa.—21° 55' N., 31° 25' E.
- Waki (R.).—1° 30' to 1° 50' N., 31° 25' E.
- Walfish Bay.—22° 55' S., 14° 30' E.
- Walikale.—1° 25' S., 28° E.
- Wambugu.—0° 40' S., 37° 5' E.
- Wanga.—4° 40' S., 39° 15' E.
- Wari, Forcados R.—5° 30' N., 5° 45' E.
- Waterval Onder.—25° 40' S., 30° 25' E.
- *Watikaia.—Belgian Congo.
- Webi (R.).—4° 10' to 7° 15' N., 39° 35' to 42° E.
- Weenen District.—28° 50' S., 30° 5' E.
- Welgelegen, Katanga.—12° S., 27° 45' E.
- Weranjanje.—1° 40' S., 31° E.
- *Wessels Neck.—Natal.
- *Willbrook.—Near Ladysmith, Natal.
- Willowmore.—33° 20' S., 23° 30' E.
- Windhoek.—22° 35' S., 17° 10' E.
- Winnebah.—5° 25' N., 0° 40' W.
- Wombali.—3° 20' S., 17° 10' E.
- *Xalosi.—Portuguese East Africa.
- *Xoce (R.).—Southern Rhodesia.
- Yabena Mabote.—1° 15' S., 24° 40' E.
- Yakuluku.—4° 20' N., 28° 50' E.
- Yakusu.—0° 35' N., 25° E.
- *Yalutcha.—Belgian Congo.
- Yambata.—2° 20' N., 22° 5' E.
- Yaminga.—2° 5' N., 22° 35' E.
- Yambuya.—1° 20' N., 24° 50' E.
- Yandumba.—0° 55' N., 23° 20' E.
- Yanga.—5° 15' S., 13° 40' E.
- *Yangandi.—Belgian Congo.
- Yanonge.—0° 40' N., 24° 30' E.
- Yaunde.—3° 50' N., 11° 35' E.
- Yukaduma.—3° 25' N., 15° E.
- Yumbi.—1° 50' S., 16° 40' E.
- Zambesi (R.).—11° 5' to 18° 45' S., 22° 35' to 36° 15' E.
- Zambi.—6° S., 12° 50' E.
- Zanzibar (I.).—6° S., 39° 30' E.
- Zomba.—15° 20' S., 35° 20' E.
- Zoutpansberg.—23° 20' S., 30° 30' E.
- Zyoul (Mt.).—8° 35' N., 38° 55' E.

FORMICIDÆ**DORYLINÆ** Leach**Dorylini** Forel**Dorylus** FABRICIUS

Dorylus FABRICIUS, 1793, 'Ent. Syst.,' II, p. 365. SHUCKARD, 1840, Ann. Nat. Hist., V, p. 268. EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, p. 697; 1910, 'Gen. Insect., Dorylinæ,' p. 5.

Typhlopone WESTWOOD, 1840, 'Introduct. Class. Insects,' II, p. 219. SHUCKARD, 1840, Ann. Nat. Hist., V, p. 262. F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 110.

Dichthadia GERSTÄCKER, 1863, Stettin. Ent. Zeitg., XXIV, p. 85.

Alaopone EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 274.

Vespa LINNÆUS. *Mutilla* LINNÆUS, LAMARCK.

Genotype: *Vespa helvola* Linnæus, 1764.

Subgenus 1. *Dorylus* FABRICIUS, sensu stricto

Dorylus subg. *Dorylus* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 701 and 714; 1910, 'Gen. Insect., Dorylinæ,' p. 9.

Subgenotype: same as genotype.

1. ***Dorylus affinis*** SHUCKARD, 1840, Ann. Nat. Hist., V, p. 316 (♂). WESTWOOD, 1842, 'Arcana Ent.,' I, p. 79 (♂). F. SMITH, 1859, 'Cat. Hym. Brit. Mus.,' VII, p. 2, Pl. I, fig. 6. ROGER, 1863, 'Verzeich. Formicid.,' p. 41. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 408. GERSTÄCKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 355; 1873, in 'v. d. Decken's Reisen in Ost Afrika, Gliederthiere,' p. 357 (♂). EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 110 (♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 9. MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 200 (♂). ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 5 (♂). EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 708 and 719, figs. H, J. (♂); 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 596, fig. (♂). H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 42. EMERY, 1899, Ann. Mus. Civ. Genova, XXXIX, p. 499 (♂); 1901, Mem. Accad. Sc. Bologna, (5) IX, pp. 416, 417, 421, 428, and 429 (♀, larva, pupa), Pl. I, figs. 1-8, 10-11; Pl. II, figs. 1-9. FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 47 (♂). MAYR, 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. 1, p. 387; 1907, in Sjöstedt, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 8 (♂). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, p. 352 (1909), (♂). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 53. ZAVATTARI, 1909, 'Il Ruwenzori, Parte Scientific.,' I, p. 212 (♂). SANTSCHI, 1910, Rev. Suisse Zool., XVIII, pp. 737 and 744, fig. 1d (♂). EMERY, 1910, 'Gen. Insect., Dorylinæ,' p. 9 (♂, ♀). FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 362 (♀). STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 375 (♂). FOREL, 1911, Rev. Zool. Afr., I, p. 274 (♀, ♂). SILVESTRI, 1913, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 145; 'Report Exped. to Africa in Search Nat. Enem. Fruit Flies,' 1914, p. 128. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 58 (♀, ♂); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 332 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 120 (2, ♀, ♂), Pl. IV, fig. 35. SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, p. 18 (♂).

Dorylus planiceps HALDEMAN, 1849, Proc. Acad. Nat. Sci. Philadelphia, IV, p. 204 (♂).

Typhlopone oraniensis var. *brevinodosa* MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 737 (♀); 1863, *ibid.*, XIII, p. 457.

Typhlopone brevinodosa ROGER, 1863, 'Verzeich. Formicid.,' p. 20.

Dorylus brevinodosa EMERY, 1887, Bull. Soc. Ent. Italiana, XIX, p. 349. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 10. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 5 (♀). EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, p. 707 and 717, figs. F1-3 (♀); 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 596 (♀). ZAVATTARI, 1907, Boll. Mus. Zool. Anat. Comp. Torino, XXII, No. 550, p. 2 (♀).

Type locality: Gambia River, GAMBIA.

SENEGAMBIA: Dakar (F. Silvestri). PORTUGUESE GUINEA: Bambaya (Lucas). FRENCH GUINEA: Konakry; Camayenne (F. Silvestri). LIBERIA: Cape Palmas (Savage). GOLD COAST: Kitta; Akra (H. Brauns). CAMEROON: (Conradt). FERNANDO PO: (Conradt). FRENCH CONGO: Gaboon (Dinklage); Ogowé (Moequers); Brazzaville (A. Weiss); southern Darbanda, Krebedje; Cape Lopez. BELGIAN CONGO: Leopoldville (de Pauw); Kitobola (Rovere). PORTUGUESE CONGO: Landana. NORTHERN RHODESIA: Kazungula (Jallá). GERMAN EAST AFRICA: Amani (H. Prell); Kibonoto, Kilimanjaro (Sjöstedt); Lake Galago in northern Ruanda (Schubotz). ZANZIBAR: (Stuhlmann). BRITISH EAST AFRICA: Mombasa (v. d. Decken; C. Alluaud); Kavirondo Bay, Victoria Nyanza; Mau Escarpment between El Burgon and Ndjoro, 2100 m. (Alluaud and Jeannel). UGANDA: Unyoro Province near Hoima; east of Lake Albert; M'Bale (C. Alluaud); Ibanda; Kaibo (Duke of Abruzzi); Gondokoto (F. Werner). SOMALILAND: Errer-es-Saghir (Brichetti-Robecchi). ABYSSINIA: Budito to Dimé (V. Bottego); Webi; Magala Re Umberto; Ogaden; Ganale (Ruspoli); ANGLO-EGYPTIAN SUDAN: Kor Attar south of Fashoda (F. Werner); Blue Nile (C. Alluaud).

1. Var. *ægyptiacus* (MAYR) EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 708 and 720 (♂); 1896, Ann. Mus. Civ. Genova, XXXVII, p. 153 (♂); 1897, *ibid.*, XXXVIII, p. 596, fig. (♂). MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 1 (♂). SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 744 (♂). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 10 (♂). FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 248 (♀, ♂). KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 3 (♂). EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, pp. 3 and 5 (♀, ♂). WASMANN, 1916, Ent. Mitt. Deutsch. Ent. Mus. Berlin, V, p. 109; 1917, Zeitschr. Wiss. Zool., CXVII, p. 309.

Dorylus ægyptiacus MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidae, p. 76, footnote (♂). ERN. ANDRÉ, 1882, 'Spec. Hym. Europ. Algérie,' II, p. 255 (♂). MAGRETTI, 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 244 (♂); 1884, Ann. Mus. Civ. Genova, XXI, p. 538 (♂). EMERY, 1892, Ann. Soc. Ent. France, XLI, Bull., p. liv.

Dorylus affinis subsp. *ægyptiacus* SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, p. 19 (♂).

Dorylus affinis var. *abyssinicus* EMERY, 1901, Mem. Accad. Sc. Bologna, (5) IX, p. 424 (♀), Pl. I, fig. 9; 1910, 'Gen. Insect., Dorylinae,' p. 10 (♀).

Typhlopone brevinodosa EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 381, fig. (♀); 1881, *ibid.*, XVI, p. 271 (♀).

Dorylus brevinodosa FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 78.

Dorylus brevinodosus var. *abyssinicus* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, p. 717 (♂).

Type locality: EGYPT.

ERITREA: Nefasit (K. Escherich; F. Silvestri); Keren (Beccari; F. Silvestri); Ghinda (F. Silvestri); Aikota (Magretti). ANGLO-EGYPTIAN SUDAN: Sennar (C. Alluaud); Kawa, White Nile (I. Trägårdh); Khartum (Karawaiew); El Hefera on Settit; Kor Guillo, Homran (Magretti). BRITISH EAST AFRICA: Maddo Wells (V. Bottego). ABYSSINIA: Let Marefia, Schoa (Antinori); Salole (Ruspoli); southern Abyssinia (Ilg).

12. Var. *exilis* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 59 (♂); 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, p. 18 (♂).

Dorylus affinis subsp. *lowyi* var. *exilis* SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 744 (♂).

Type locality: Kibosho, Mt. Kilimanjaro, GERMAN EAST AFRICA (C. Alluaud).

GERMAN EAST AFRICA: New Moschi, 800 m.; Himo River, 1000 m.; Kilema, 1400 m. (Alluaud and Jeannel). BRITISH EAST AFRICA: Nairobi; Rurunga, Kikuyu; Bura, Wa-Taita; Taveta; Mombasa; Ngare Nyuki, Kenia (Alluaud and Jeannel).

13. Var. *hirsutus* (SANTSCHI).

Dorylus affinis subsp. *egyptiacus* var. *hirsutus* SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, pp. 19 and 20 (♂).

Type locality: ABYSSINIA (Reichensperger).

14. Var. *parapsidalis* SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, pp. 19 and 20 (♂).

Type locality: Blantyre, NYASALAND (G. Arnold).

15. Var. *pulliceps* SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, pp. 18 and 19 (♂).

Type locality: IVORY COAST (Lohier).

16. Var. *sudanicus* SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, pp. 18 and 19 (♂).

Type locality: Fort Sibut, southern Darbanda, FRENCH CONGO (Decorse).

1a. Subsp. *denudatus* SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, p. 19 (♂).

Dorylus denudatus SANTSCHI, 1910, Rev. Suisse Zool., XVIII, pp. 737 and 742, figs. 1a and 4 (♂).

Type locality: Upper Niger (Claveau).

1b. Subsp. *lowyi* FOREL, 1907, Ann. Soc. Ent. Belgique, LI, p. 202 (♂). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 10 (♂). SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 744 (♂). FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 254 (♂).

Dorylus affinis subsp. *lowyi* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 60 (♂).

Dorylus affinis subsp. *lowyi* SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, pp. 19 and 20 (♂).

Type locality: ZANZIBAR (Löwy).

ERITREA. BRITISH EAST AFRICA: Mombasa; Athi Plain (Feringue); Rurunga, Kikuyu; Wambogo; Nairobi, Masai Plain (Alluaud and Jeannel). RHODESIA: Bulawayo (G. Arnold).

2. *Dorylus alluaudi* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 60, fig. 4 (♂).

Type locality: Eastern slopes of Mt. Ruwenzori, 1600 m., UGANDA (C. Alluaud).

3. **Dorylus atratus** F. SMITH, 1859, 'Cat. Hym. Brit. Mus.,' VII, p. 4 (♂), Pl. I, fig. 15. ROGER, 1863, 'Verzeich. Formicid.,' p. 41. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 408. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 10. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 5 (♂). EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 708 and 722 (♂). W. A. SCHULZ, 1906, 'Spolia Hymenopterologica,' p. 297 (♂). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 127 (♂). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 10 (♂). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 374 (♂). See p. 43.

Type locality: Old Calabar, NIGERIA.

FERNANDO PO: (Conradt); Santa Isabel (Schultze). CAMEROON: Bibundi (Tessmann). FRENCH CONGO: Ogowé (Mocquerys). BELGIAN CONGO: Stanleyville (Lang and Chapin).

4. **Dorylus bequaerti** FOREL, 1913, Rev. Zool. Afr., II, p. 311 (♀). BEQUAERT, *ibid.*, p. 423. See p. 43.

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

BELGIAN CONGO: Pasacorde near Zambi (J. Bequaert); Banana (Lang and Chapin).

5. **Dorylus braunsi** EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 707 and 718, fig. G (♀); 1901, Mem. Accad. Sc. Bologna, (5) IX, pp. 416 and 428 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 352 (♀). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 10 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 120 (♀, ♂).

Type locality: LIBERIA (H. Brauns).

FRENCH CONGO: Mandouga; Mbamu (A. Weiss). RHODESIA: Bulawayo (G. Arnold).

5a. Subsp. **anceps** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 215 (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

6. **Dorylus brevipennis** EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 708 and 721, figs. K and L (♂); 1901, Mem. Accad. Sc. Bologna, (5) IX, pp. 424 and 428 (♀), Pl. I, fig. 12. MAYR, 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. 1, p. 387. ZAVATTARI, 1907, Boll. Mus. Zool. Anat. Comp. Torino, XXII, No. 550, p. 2 (♂). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 10 (♀, ♂). SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 741 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 61 (♀); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 14 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 122 (♀, ♂).

Type locality: Tabora, GERMAN EAST AFRICA.

ZULULAND: Umfolosi (I. Trägårdh). NORTHERN RHODESIA: Kazungula (Jallé). BRITISH EAST AFRICA: Gilgil, Rift Valley, 1980 m. (Alluaud and Jeannel). UGANDA: Bujongolo Region (C. Alluaud). ANGLO-EGYPTIAN SUDAN: Mongalla opposite Lado (F. Werner).

6i. Var. **marshalli** EMERY, 1901, Mem. Accad. Sc. Bologna, (5) IX, p. 425 (♀, ♂), Pl. I, fig. 13. SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 741 (♀). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 10 (♀, ♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 122 (♀, ♂). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 402 (♀). See p. 43.

Type locality: Salisbury, SOUTHERN RHODESIA (H. Brauns).

MOZAMBIQUE: Valley of the Revoue (G. Vasse). BELGIAN CONGO: (Kohl); Medje (Lang and Chapin).

6a. Subsp. *zimmermanni* SANTSCHI, 1910, Rev. Suisse Zool., XVIII, pp. 737 and 738, figs. 1c and 2a-c (♀, ♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 123 (2, ♀, ♂).

Type locality: Madingu, FRENCH CONGO (P. Zimmermann).

SOUTHERN RHODESIA: Hillside, Bulawayo (G. Arnold).

6a1. Var. *bulawayensis* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 364.

Dorylus affinis subsp. *zimmermanni* var. *bulawayensis* FOREL, 1914, *ibid.*, (5) L, p. 215 (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

7. *Dorylus depilis* (EMERY) FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 71 (♂). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 10 (♂). SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 737, fig. 1e (♂). FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 254 (♀); 1911, Rev. Zool. Afr., I, p. 274 (♂); 1913, *ibid.*, II, p. 312 (♂); 1913, Ann. Soc. Ent. Belgique, LVII, p. 348 (♂); 1913, Rev. Suisse Zool., XXI, p. 668 (♂). See p. 43.

Dorylus affinis var. *depilis* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 708 and 720 (♂). MAYR, 1896, Ent. Tidskr., XVII, p. 230 (♂).

Dorylus affinis subsp. *depilis* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 53 (♂); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 4 (♂); 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 362 (♂). SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, p. 19 (♂).

Type locality: GOLD COAST.

WEST AFRICA: (Fülleborn). TOGO. CAMEROON: (Sjöstedt; Conradt). NIGERIA: Old Calabar (Bates). BELGIAN CONGO: Sankuru (Laja); Valley of the Lubumbashi (Buttgenbach); Kitobola (Rovere); Lukula (Daniel); Kabambare (Flamand); Kapiri (Leplae); Api (Laplume); Sankisia (J. Bequaert); Dima (A. Koller); Faradje; Medje; Stanleyville (Lang and Chapin). GERMAN EAST AFRICA: Amani (Zimmer). BRITISH EAST AFRICA: Nairobi (H. Prell). UGANDA: (Benoit).

71. Var. *clarior* (SANTSCHI).

Dorylus affinis subsp. *depilis* var. *clarior* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 246 (♂); 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, p. 19 (♂).

Type locality: Zambézie (Durand).

BELGIAN CONGO: Ubangi (Augustin). FERNANDO PO.

72. Var. *ugandensis* (SANTSCHI).

Dorylus affinis var. *ugandensis* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicide,' p. 59 (♂).

Dorylus affinis subsp. *depilis* var. *ugandensis* SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, p. 19 (♂).

Type locality: Unyoro Province, east of Lake Albert, UGANDA (C. Alluaud).

8. *Dorylus furcatus* (GERSTÄCKER) DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 10. EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, p. 718 (♀); 1910, 'Gen. Insect., Dorylinae,' p. 10 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 124 (♀).

Dichthadia furcata GERSTÄCKER, 1872, Stettin. Ent. Zeitg., XXXIII, p. 267 (♀). EMERY, 1887, Bull. Soc. Ent. Italiana, XIX, p. 350 (♀), Pl. xi, figs. 4 and 5.

Type locality: SOUTH AFRICA.

According to Arnold this is perhaps only an individual variation of *D. helcolus* ♀ or a variety of that species.

9. **Dorylus gaudens** SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 244 (♂).

Type locality: BOGA, BELGIAN CONGO (A. Pilette).

10. **Dorylus gribodoi** EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 570 (♂), Pl. xv, fig. 15; 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 708 and 723, fig. O (♂). ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 5 (♂). FOREL, Mitt. Naturh. Mus. Hamburg, XXIV, p. 14 (♂). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 10 (♂). SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 745 (♂); 1911, Ann. Soc. Ent. Belgique, LIV, p. 282 (♂). FOREL, 1913, Rev. Zool. Afr., II, p. 312 (♂).

Type locality: Amu, Togo.

LIBERIA: (Kieselbach). IVORY COAST: (A. Richard; Lohier). GOLD COAST. NIGERIA: Lower Benue (Lenfant). FRENCH CONGO: Ogowe (Mocquerys). BELGIAN CONGO: Kondué (Luja).

101. Var. **confusus** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 246 (♂).

Type locality: Grand Bassam, IVORY COAST (Lohier).

IVORY COAST: near Imbokro (Posth).

11. **Dorylus helvolus** (LINNÆUS) FABRICIUS, 1793, 'Ent. Syst.,' II, p. 365 (♂). COQUEBERT, 1801, 'Illustr. Iconogr. Ins.,' II, p. 65 (♂), Pl. xvi, fig. 1. LATREILLE, 1802, 'Hist. Nat. Crust. Ins.,' III, p. 352 (♂). FABRICIUS, 1804, 'Syst. Piez.,' p. 427 (♂). LATREILLE, 1805, 'Hist. Nat. Crust. Ins.,' XIII, p. 260 (♂, in part). JURINE, 1807, 'Nouv. Méth. Class. Hym.,' p. 281 (♂). LATREILLE, 1809, 'Gen. Crust. Ins.,' IV, p. 124 (♂). LEPELETIER, 1836, 'Hist. Nat. Ins. Hym.,' I, p. 228 (♂). BLANCHARD, 1840, 'Hist. Nat. Ins.,' III, p. 378 (♂). SCHUCKARD, 1840, Ann. Nat. Hist., V, p. 315 (♂). WESTWOOD, 1842, 'Arcana Ent.,' I, p. 79 (♂). FAIRMAIRE, in J. Thomson, 1858, 'Archives Entomologiques,' II, p. 263. F. SMITH, 1859, 'Cat. Hym. Brit. Mus.,' VII, p. 1, Pl. 1, fig. 5. DUMÉRIL, 1860, Mém. Acad. Sc. Paris, XXXI, p. 901 (♂). ROGER, 1863, 'Verzeich. Formicid.,' p. 41. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 408. TRIMEN, 1880, Trans. Ent. Soc. London, Proc., p. xxiv, fig. and p. xxiii (♀); 1881, Trans. South African Phil. Soc., II, 1, Proc., p. xvii. EMERY, 1887, Ann. Mus. Civ. Genova, XXV, p. 449, footnote (♂, ♂); 1887, Bull. Soc. Ent. Italiana, XIX, p. 349 (♂, ♀, ♂), Pl. xi, figs. 1-3 and 6-7. DISTANT, 1892, 'A Naturalist in the Transvaal,' p. 211. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 11. H. STADELMANN, 1893, Mitth. Deutsch. Schutzgeb., VI, p. 217. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 17 (♂); *ibid.*, Bull. p. lxxiii, fig. (♀); 1895, Ann. Mus. Civ. Genova, XXXV, p. 177 (♂); 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 707, 708, and 714, figs. C-E (♂, ♂). WASMANN, 1897, Deutsch. Ent. Zeitschr., p. 278; 1898, Wien. Ent. Zeitg., XVII, p. 103. H. BRAUNS, *ibid.*, p. 226. WASMANN, 1899, Deutsch. Ent. Zeitschr., pp. 174 and 175; 1900, Zool. Jahrb. Abt. Syst., XIV, pp. 266, 268, and 274. EMERY, 1901, Mem. Accad. Sc. Bologna, (5) IX, pp. 416 and 428 (♂). H. BRAUNS, 1901, Zeitschr. f. Hym. u. Dipt., I, p. 16 (♂). BINGHAM, 1903, Ann. Mag. Nat. Hist., (7) XII, p. 61. WASMANN, 1904, Zool. Jahrb. Abt. Syst., Suppl. VII, p. 676. BINGHAM, 1906, Trans. Ent. Soc. London, Proc., p. xxvi. FOREL, 1907, Mitt. Naturh. Mus. Hamburg, XXIV, p. 14 (♂). MAYR, 1907, in Sjöstedt, 'Expéd. Kilimandjaro, Meru, etc.,' II, 8, p. 7 (♂). DIXEY AND LONGSTAFF, 1907, Trans. Ent. Soc. London, pp. 334 and 376 (♂). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 10 (♂, ♂). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 4 (♂, ♂); 1913, Rev. Zool. Afr., II, p. 311 (♂). BEQUAERT, *ibid.*, p. 423. SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd.,

III, p. 14 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 61 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 332 (♀). H. SCHMITZ, 1914, Zool. Jahrb. Abt. Syst., XXXVII, pp. 522 and 559. LAMBORN, 1914, Trans. Ent. Soc. London, (1913), Proc., p. cxxiii. ARNOLD, 1915, Ann. South African Mus., XIV, p. 116 (2, ♀, ♀, ♂), Pl. IV, figs. 33, 33a, 33b, 34, and 34a. WASMANN, 1916, Ent. Mitt. Deutsch. Ent. Mus. Berlin, V, p. 136; 1917, Zeitschr. Wiss. Zool., CXVII, pp. 304 and 352.

Vespa helvola LINNÆUS, 1764, 'Mus. Ludov. Ulric.', p. 412 (♂).

Mutilla helvola LINNÆUS, 1767, 'Syst. Nat.', Ed. 12, I, 2, p. 967 (♂). PH. L. MÜLLER, in LINNÆUS, 1775, 'Vollst. Natursyst.', V, p. 920 (♂). FABRICIUS, 1787, 'Mantissa Insect.', I, p. 313 (♂). GMELIN, 1790, in LINNÆUS, 'Syst. Nat.', Ed. 13, I, 5, p. 2807 (♂). CHRIST, 1791, 'Naturg. d. Insect.', p. 151 (♂).

Mutilla dorylus LAMARCK, 1817, 'Hist. Nat. Anim. s. Vertébr.', IV, p. 101 (♂); 1835, *ibid.*, 2e éd., IV, p. 316 (♂).

Mutilla (Dorylus) helvolus BLANCHARD, in Cuvier, 1849, 'Règne Animal,' Ed. 3, Insect., II, Pl. CXVIII, fig. 1 (♂).

Typhlopone punctata F. SMITH, 1858, 'Cat. Hym. Brit. Mus.', VI, p. 112 (♀). ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 45; 1863, 'Verzeich. Formicid.', p. 20. MAYR, 1865, 'Reise Novara, Zool.', II, Formicidæ, p. 74. W. F. KIRBY, 1884, Ann. Mag. Nat. Hist., (5) XIII, p. 406 (♀).

Typhlopone europæa ROGER, 1859, Berlin. Ent. Zeitschr., III, p. 248 (♀), Pl. VII, fig. 3. MAYR, 1861, 'Europ. Formicid.', p. 53 (♀). ROGER, 1863, 'Verzeich. Formicid.', p. 20. ERN. ANDRÉ, 1874, Rev. Mag. Zool., (3) II, p. 187 (♀). EMERY, 1876, Stettin. Ent. Zeitg., XXXVII, p. 72 (♀).

Typhlopone europæa MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 457.

Typhlopone punctata MAYR, *ibid.*, p. 457.

Dorylus planifrons MAYR, 1865, 'Reise Novara, Zool.', II, Formicidæ, p. 74 (♂), Pl. III, fig. 21.

Dorylus punctatus ERN. ANDRÉ, 1882, 'Spec. Hym. Europ. Algérie,' II, p. 251 (♀), Pl. xv, fig. 5.

Type locality: Cape of Good Hope.

Togo: Bismarckburg (Büttner). SOUTHERN NIGERIA: Agege (Lamborn). BELGIAN CONGO: Elisabethville (J. Bequaert). SOUTHERN RHODESIA: common (G. Arnold). TRANSVAAL: Barberton (Rendall); Pretoria (Wichgraf; F. Silvestri; Distant); Makapan; Hebron (E. Simon). ORANGE FREE STATE: Kroonstad (H. Brauns); Bloemfontein (E. Simon). CAPE PROVINCE: Cape Town (E. Simon; Schönland; Wilms); Dunbrody; Blue Cliff (O'Neil); East London (Dixey and Longstaff); Port Elizabeth (H. Brauns). NATAL: (Kluckauf); Estcourt (I. Trägårdh); Port Natal (Schultze); Durban (Wilms; Hahn); Ladysmith (Dixey and Longstaff). NYASALAND: Zomba; Fort Johnston (Rendall). GERMAN EAST AFRICA: Mombo, Usambara; Kibonoto, Mt. Kilimanjaro (Sjöstedt). BRITISH EAST AFRICA: Kenia Region, between the Nyeri and Amboni Rivers, 1800 m. (Alluaud and Jeannel). ABYSSINIA: Arussi, Galla, Ganale Gudda (V. Bottego).

11. Var. *impressus* STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 127 (2, ♀).

Type locality: Alen, SPANISH GUINEA (Tessmann).

12. *Dorylus lobatus* SANTSCI, 1919, Rev. Zool. Afr., VI, p. 244 (♂).

Type locality: Lukolela, BELGIAN CONGO (Christy).

BELGIAN CONGO: Kabinda (Schwetë).

13. *Dorylus mandibularis* MAYR, 1896, Ent. Tidskr., XVII, p. 228 (♂). Sjöstedt, 1904, 'I Västafrikas Urskogar,' p. 157, footnote. EMERY, 1910, 'Gen. Insect., Doryline,' p. 10 (♂).

Type locality: CAMEROON (Sjöstedt).

13a. Var. *pulchellus* SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 361 (♂).

Type locality: Dimbroko, IVORY COAST (Le Moulé).

14. *Dorylus maestus* (EMERY) FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 71 (♂). EMERY, 1910, 'Gen. Insect., Doryline,' p. 10 (♂). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 348 (♂). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 374 (♂). See p. 43.

Dorylus affinis var. *maestus* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 708 and 720 (♂). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 53 (♂).

Type locality: CONGO.

BELGIAN CONGO: Sankuru (Luja); Lukula (Daniel); Stanleyville (Lang and Chapin); Kimuenza (Schultze).

14a. Var. *claripennis* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 243 (♂).

Type locality: Lower Kasai, BELGIAN CONGO (Vanderyst).

14b. Var. *morio* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 243 (♂).

Dorylus maestus SANTSCHI, 1910, Rev. Suisse Zool., XVIII, pp. 737 and 741, figs. 1b and 3 (♂).

Type locality: Madingu, FRENCH CONGO (Zimmermann).

14a. Subsp. *schereri* FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 254 (♂).

Dorylus maestus var. *schereri* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 244 (♂).

Type locality: LIBERIA (Scherer).

IVORY COAST: Dimbroko.

15. *Dorylus politus* EMERY, 1901, Mem. Accad. Sc. Bologna, (5) IX, pp. 416, 426, and 428 (♀), Pl. I, figs. 14-17; 1910, 'Gen. Insect., Doryline,' p. 10 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 348 (♀); 1913, Rev. Suisse Zool., XXI, p. 668 (♀).

Type locality: CAMEROON (Conradt).

BELGIAN CONGO: Mayombe (de Brier). CAMEROON: Johann-Albrechtshöhe (Conradt).

16. *Dorylus spininodis* EMERY, 1901, Mem. Accad. Sc. Bologna, (5) IX, pp. 426 and 428 (♀), Pl. I, figs. 18-21. SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 352 (♀). EMERY, 1910, 'Gen. Insect., Doryline,' p. 10 (♀). SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 747 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 332 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicide,' p. 60 (♀).

Type locality: CAMEROON (Conradt).

CAMEROON: Victoria (F. Silvestri). NIGERIA: Benin; Olokemeji (F. Silvestri). FRENCH CONGO: Brazzaville (A. Weiss). UGANDA: Toro Province near Fort Portal (C. Alluaud).

16a. Subsp. *longiceps* VIERMEYER, 1914, Arch. f. Naturg., LXXIX, Abt. A, Heft 12, (1913), p. 26, fig. 1 (♀).

Type locality: Marow, Langenburg, GERMAN EAST AFRICA.

17. *Dorylus stadelmanni* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 708 and 722, figs. N and M (♂). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 53 (♂). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 10 (♂).

Type locality: CONGO.

NORTHEASTERN AFRICA: Terwidja (von Erlanger).

18. *Dorylus staudingeri* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 708 and 721 (♂); 1910, 'Gen. Insect., Dorylinae,' p. 10 (♂). See p. 43.

Type locality: CONGO.

BELGIAN CONGO: Medje (Lang and Chapin).

19. *Dorylus striatidens* SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 745, fig. 5 (♀).

Type locality: CASAMANCE (Claveau).

SENEGAMBIA: Dakar (Melou).

Subgenus 2. *Anomma* SHUCKARD

Anomma SHUCKARD, 1840, Ann. Nat. Hist., V, p. 326.

- Dorylus* subg. *Anomma* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 701 and 710; 1910, 'Gen. Insect., Dorylinae,' p. 10.

Sphegomyrmex IMHOFF, 1852, Verh. Naturf. Ges. Basel, X, p. 175.

Alaopone (part) EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 274.

- Subgenotype: *Anomma burmeisteri* Shuckard, 1840 = *Dorylus nigricans* Illiger (1802) subsp. *burmeisteri* (Shuckard).

20. *Dorylus (Anomma) emeryi* MAYR, 1896, Ent. Tidskr., XVII, p. 225 (♀). EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 462; 1901, Mem. Accad. Sc. Bologna, (5) IX, p. 428 (♀); 1910, 'Gen. Insect., Dorylinae,' p. 11 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 127 (♀). SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 154, figs. A7 and B9 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 333 (♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 402 (♀).

Dorylus emeryi SÖSTEDT, 1904, 'I Västafrikas Urskogar,' p. 157, footnote.

Type locality: CAMEROON (Sjöstedt; Conradt).

- GOLD COAST: Aburi (F. Silvestri). CAMEROON: Mundame (Conradt). BELGIAN CONGO: (Kohl).

- 20a. Var. *pulsi* FOREL, 1904, Ann. Soc. Ent. Belgique, XLVIII, p. 170 (♀). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 11 (♀). SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 154 (♀).

Type locality: West Africa.

- 20a. Subsp. *opacus* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 58 (♀). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 11 (♀). FOREL, 1911, Rev. Suisse Zool., XIX, p. 452 (♂); 1913, Ann. Soc. Ent. Belgique, LVII, p. 348 (♀). See p. 43.

- Dorylus (Anomma) opacus* SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, pp. 154 and 157, figs. A8, B3, and B10 (♂, ♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 402 (♂).

- Dorylus (Anomma) emeryi* var. *opaca* STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 127 (♀, ♀).

Type locality: Lower Congo, in stomach of *Manis temmincki* (Solon).

- BELGIAN CONGO: (Kohl); Mayombe (de Briey); Medje; Ngayu (Lang and Chapin). CAMEROON: Mundame (Conradt). FERNANDO PO: (Conradt). FRENCH CONGO: Gaboon (F. Faure).

21. *Dorylus (Anomma) funereus* EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 11 (♂). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 402 (♂). See p. 44.

Dorylus (Anomma) nigricans var. *funereus* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 708 and 712 (♂); 1899, Ann. Soc. Ent. Belgique, XLIII, p. 461. FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 70 (♂); 1913, Rev. Zool. Afr., II, p. 311 (♂); 1913, Rev. Suisse Zool., XXI, p. 667 (♂); 1913, Ann. Soc. Ent. Belgique, LVII, p. 348 (♂).

Type locality: GOLD COAST.

BELGIAN CONGO: Sankuru; Kondué (Luja); Lukula (Daniel); Medje; Stanleyville; Bolobo to Lukolela (Lang and Chapin). CAMEROON: (Conradt).

22. *Dorylus (Anomma) gerstäckeri* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 707 and 713 (♀), Pl. XVII, fig. 11.

Anomma gerstäckeri EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 462 (♀).

Dorylus (Anomma) gerstäckeri EMERY, 1901, Mem. Accad. Sc. Bologna, (5) IX, pp. 416, 427, and 428 (♀), Pl. I, fig. 22; 1910, 'Gen. Insect., Dorylinae,' p. 11 (♀); SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 332 (♀).

Dorylus (Anomma) gerstäckeri SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 154 (♀).

Type locality: AKRA, GOLD COAST.

GOLD COAST: Aburi (F. Silvestri). CAMEROON: (Conradt).

- 22a. Subsp. *quadratus* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 430 (♀).

Type locality: MANOW, GERMAN EAST AFRICA.

23. *Dorylus (Anomma) kohli* WASMANN, 1904, Zool. Jahrb. Abt. Syst., Suppl. VII, pp. 669 and 675 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 352 (♀). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 11 (♀). FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 254 (♀). SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 154, fig. C1 (♀). WASMANN, 1916, Ent. Mitt. Deutsch. Ent. Mus. Berlin, V, pp. 105 and 147; 1917, Zeitschr. Wiss. Zool., CXVII, pp. 290 and 322. See p. 44.

Anomma kohli H. SCHMITZ, 1914, Zool. Jahrb. Abt. Syst., XXXVII, pp. 515 and 540, footnote; 1915, Deutsch. Ent. Zeitschr., p. 487. WASMANN, 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, p. 204.

Type locality: Stanleyville, BELGIAN CONGO (Kohl).

NIGERIA: Old Calabar (Bates). FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: Akenge; Niangara; Avakubi (Lang and Chapin); St. Gabriel (Kohl).

- 23₁. Var. *chapini* Wm. M. WHEELER. See p. 45 (♀).

Type locality: Stanleyville, BELGIAN CONGO (Lang and Chapin).

- 23₂. Var. *congolensis* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 352, fig. 1 (♀); 1912, Ann. Soc. Ent. Belgique, LVI, pp. 154 and 158, figs. A6, B11, and C2 (♀). See p. 45.

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

FRENCH CONGO: N'Douma (A. Weiss). BELGIAN CONGO: Leopoldville (Lang and Chapin); Thysville (J. Bequaert).

- 23₃. Var. *frenisyi* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 402 (♀).

Type locality: BELGIAN CONGO (Kohl).

- 23₄. Var. *langi* Wm. M. WHEELER. See p. 45 (♀).

Type locality: Malela, BELGIAN CONGO (Lang and Chapin).

23. Var. *minor* SANTSCHI, 1911, Rev. Zool. Afr., I, p. 206 (♀); 1912, Ann. Soc. Ent. Belgique, LVI, p. 155, figs. A9 and B8 (♀).

Type locality: CUCALA, BENGUELA (J. Cruchet).

24. *Dorylus (Anomma) mayri* SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, pp. 155 and 159, figs. A10, B7 and B13 (♀).

Type locality: CAMEROON.

25. *Dorylus (Anomma) nigricans* ILLIGER. EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 706, 708, and 710 (♀, ♂), Pl. XIV, figs. 1a-g. MAYR, 1896, Ent. Tidskr., XVII, p. 225 (♀, ♂). EMERY, 1901, Mem. Accad. Sc. Bologna, (5) IX, pp. 416 and 428 (♀). FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 47 (♂); 1907, *ibid.*, XXIV, p. 15 (♂); 1909, Ann. Soc. Ent. Belgique, LIII, p. 51 (♂). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 352 (♂). FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 248 (♀). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 12 (♀, ♂). STRITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 128 (♀, ♂); 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 375 (♀). FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 362 (♀); 1912, Mitt. Naturh. Mus. Hamburg, XXIX, p. 177, Pl., figs. 1, 3a, 4a, and 6-8 (♀, ♀). SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 156 (♀). FOREL, 1913, *ibid.*, LVII, p. 348 (♀); 1913, Rev. Zool. Afr., II, p. 311 (♂). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 332 (♂); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 62 (♂). FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 216 (♂).

Dorylus nigricans ILLIGER, 1802, Mag. f. Insectenk., I, p. 188 (♂). FABRICIUS, 1804, 'Syst. Piez.,' p. 427 (♂). JURINE, 1807, 'Nouv. Méth. Class. Hym.,' p. 281 (♂). LATREILLE, 1809, 'Gen. Crust. Ins.,' IV, p. 124 (♂). SHUCKARD, 1840, Ann. Nat. Hist., V, p. 271 (♂). WESTWOOD, 1842, 'Arcana Ent.,' I, p. 79 (♂). GUÉRIN, in Lefebvre, 1848, 'Voy. Abyssinie,' IV, Zool., pt. 6, p. 353. F. SMITH, 1859, 'Cat. Hym. Brit. Mus.,' VII, p. 2 (♂), Pl. I, fig. 7. ROGER, 1863, 'Verzeich. Formicid.,' p. 41. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 408. GERSTAECKER, 1871, Arch. f. Naturg., XXXVII, I, p. 355; 1873, in 'v. d. Decken's Reisen in Ost-Afrika, Gliederthiere,' p. 357 (♂). RITSEMA, 1874, Tijdschr. v. Ent., XVII, p. 182 (♂). EMERY, 1892, Ann. Soc. Ent. France, LXI, Bull., p. liv; 1892, *ibid.*, XL, (1891), p. 555. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 12. H. STADELMANN, 1893, Mitth. Deutsch. Schutzgeb., VI, p. 217. ERN. ANDRÉ, 1895, Rev. d'Ent. Cren., XIV, p. 5 (♂). H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 42. MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 1 (♂). F. C. WELLMAN, 1908, Ent. News, XIX, p. 226. ZAVATTARI, 1909, 'Il Ruwenzori, Parte Scientific.,' I, p. 212 (♀). BINGHAM, 1909, Trans. Zool. Soc. London, XIX, p. 180. BEQUAERT, 1913, Rev. Zool. Afr., II, p. 423. LAMBORN, 1914, Trans. Ent. Soc. London, (1913), Proc., p. exxiii; 1914, *ibid.*, (1914), Proc., p. vi. G. D. H. CARPENTER, 1915, *ibid.*, (1914), Proc., p. cvii.

? *Dorylus* sp. WATERHOUSE, 1907, Trans. Ent. Soc. London, Proc., p. vi, figs. (♀, ♀).

? *Dorylus nigrilaris* E. STRAND, 1911, Jahrb. Nassau. Ver. Naturk. Wiesbaden, LXIV, p. 118 (♀).

Type locality: SIERRA LEONE.

LIBERIA: (Kieselbach). FRENCH GUINEA: Konakry (F. Silvestri). IVORY COAST: Assinie (C. Alluaud). NIGERIA: Moor Plantation near Ibadan; Agege (Lamborn). CAMEROON: (Sjöstedt); Mowange (C. Feldmann); Jabassi (Riggen-

bach); Yaunde (Scheunemann); Bipindi (Zenker); Bibundi (Tessmann); Mukonje Farm (R. Rohde). SPANISH GUINEA: Alen (Tessmann). FRENCH CONGO: Brazzaville (A. Weiss). PORTUGUESE CONGO: Landana. BELGIAN CONGO: Elisabethville, Katanga (J. Bequaert); Congo da Lemba (R. Mayné); Kitobola (Rovere); Mayombe (de Briey; Cabra); Chikai (Cabra); Zambi (don Lopez); northwestern shore of Lake Tanganyika (Grauer). NYASALAND: Madona, Nyamias (Montgomery). GERMAN EAST AFRICA: Amani (Zimmer); Tanga (H. Prell). ZANZIBAR: (Stuhlmann). BRITISH EAST AFRICA: Bura Mts. (v. d. Decken). UGANDA: M'Bale (C. Alluaud); Bihunga (Duke of Abruzzi); Mubuku Valley, eastern slopes of Mt. Ruwenzori, 5000-7000 ft. (Woosnam); Bugalla Island; Damba Island (G. D. H. Carpenter). WESTERN ABYSSINIA: (Ilg). ANGOLO-EGYPTIAN SUDAN: Renk (I. Trägårdh).

25a. Subsp. *arcens* (WESTWOOD) EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 706 and 710 (♂); 1910, 'Gen. Insect., Dorylinae,' p. 12 (♂). SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, pp. 157 and 160, figs. A1 and B4 (♂); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 332 (2, ♀). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 373 (♀). See p. 46.

Anomma arcens WESTWOOD, 1847, Trans. Ent. Soc. London, V, p. 17 (♂), Pl. I, figs. 3 and 3a-f. SAVAGE, 1849, Proc. Acad. Nat. Sci. Philadelphia, IV, p. 196 (♂). F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 114, Pl. VIII, fig. 3. ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 46 (♂); 1863, 'Verzeich. Formicid.,' p. 20. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 394. PERKINS, 1869, American Naturalist, III, pp. 360-364 (♀). EMERY, 1892, Ann. Soc. Ent. France, LXI, Bull., p. liv. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 8. M. L. SYKES, 1900, Trans. Manchester Microsc. Soc., XX, (1899), p. 89.

Anomma burmeisteri subsp. *arcens* EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 554 (♀).

Anomma pubescens ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 47 (♂); 1863, 'Verzeich. Formicid.,' p. 21. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 395. EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, p. 711 (♀).

Dorylus (*Anomma*) *nigricans* subsp. *burmeisteri* var. *arcens* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 352 (♀).

Type locality: Cape Palmas, LIBERIA (Savage).

SIERRA LEONE: Kortright Hill (W. G. Clements). IVORY COAST: (Lohier); Assinie (C. Alluaud). GOLD COAST: Aburi (F. Silvestri). FRENCH CONGO: Fort Crampel (Schubotz); Brazzaville (A. Weiss); Gaboon. BELGIAN CONGO: Medje (Lang and Chapin).

25b. Subsp. *burmeisteri* (SHUCKARD) EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 706 and 710-713, figs. A, B (♂, ♂). WASMANN, 1904, *ibid.*, Suppl. VII, pp. 671 and 675 (♂). SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 157, figs. A4, B12, and B16 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 332 (♀). See p. 46.

Anomma burmeisteri SHUCKARD, 1840, Ann. Nat. Hist., V, p. 326 (♂). WESTWOOD, 1840, Ann. Mag. Nat. Hist., VI, p. 88 (♂), Pl. II, figs. 4 and 4a. F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 114, Pl. VIII, fig. 2. ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 46 (♂); 1863, 'Verzeich. Formicid.,' p. 20. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 395. EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 554 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 9. MAYR, 1893, Jahrb. Ham-

burg. Wiss. Anst., X, 2, p. 200. WASMANN, 1894, 'Verzeichn. Myrmecoph. Termitoph. Arthrop.', p. 77. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 5 (♀). WASMANN, 1916, Ent. Mitt. Deutsch. Ent. Mus. Berlin, V, p. 143; 1917, Zeitschr. Wiss. Zool., CXVII, p. 318.

Dorylus (Anomma) burmeisteri STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 373 (♀).

Type locality: SIERRA LEONE (D. F. Morgan).

SENEGAMBIA. NIGERIA: Ibadan (F. Silvestri). FERNANDO PO: on the Peak, at 2400 m. (Schultze). CAMEROON: Grand Batanga (G. Schwab). FRENCH CONGO: Ogowé (Mocquereys). BELGIAN CONGO: Duma; Libenge (Schubotz); Katala (J. Bequaert); Stanleyville; Lukolela to Basoko (Lang and Chapin); GERMAN EAST AFRICA: Mhonda (Stuhlmann). ZANZIBAR: (Stuhlmann).

25b₁. Var. *hybridus* SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, pp. 157 and 161, fig. B14 (♀). F. SILVESTRI, 1913, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 145. SANTSCHI, 1914, *ibid.*, p. 332 (♀). F. SILVESTRI, 1914, 'Report Exped. to Africa in Search Nat. Enem. Fruit Flies,' p. 128.

Type locality: CASAMANCE (Claveau).

FRENCH GUINEA: Konakry; Kakoulima; Kindia (F. Silvestri).

25b₂. Var. *molestus* (GERSTÄCKER) MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 7 (♀). Sjöstedt, 1908, *ibid.*, II, 8, p. 108. SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, pp. 156 and 162, figs. A5 and B15 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 62 (♀).

Anomma molesta GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 262 (♀). ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 47. GERSTÄCKER, in Peters, 1862, 'Reise n. Mossambique,' Zool., V, p. 302 (♀), Pl. xxxii, fig. 2. ROGER, 1863, 'Verzeich. Formicid.,' p. 21. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 395. GERSTÄCKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 355; 1873, in 'v. d. Decken's Reisen in Ost Afrika, Gliederthiere,' p. 347 (♀). EMERY, 1887, Bull. Soc. Ent. Italiana, XIX, p. 348, footnote. WASMANN, 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, pp. 30 and 290.

Anomma burmeisteri EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 271, figs. (♂) (nec Shuckard).

Anomma burmeisteri var. *molesta* EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 554 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 9.

Dorylus (Anomma) nigricans var. *molestus* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, p. 710 (♀); 1901, Mem. Accad. Sc. Bologna, (5) IX, p. 416 (♀). WASMANN, 1904, Zool. Jahrb. Abt. Syst., Suppl. VII, pp. 671-673 and 675 (♀). ZAVATTARI, 1907, Boll. Mus. Zool. Anat. Comp. Torino, XXII, No. 550, p. 2 (♀). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 12. STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 375 (♀, ♂). SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 162 (♀). SWYNNERTON, 1916, Trans. Ent. Soc. London, (1915), pp. 317-348 (♀).

Dorylus (Anomma) nigricans subsp. *molesta* EMERY, 1912, Annuar. Mus. Zool. Univ. Napoli, N. S., III, No. 26, p. 1 (2, ♀).

Alaopone antinorii EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 275, figs. (♂).

Dorylus antinorii EMERY, 1887, Bull. Soc. Ent. Italiana, XIX, p. 349. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 10.

Dorylus (Alaopone) antinorii EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 707 and 736 (♂).

Type locality: Tete, PORTUGUESE EAST AFRICA (Peters).

BENGUELA: (J. Cruchet). PORTUGUESE EAST AFRICA: Mts. of Morumballe on the Zambesi R. (Luja). RHODESIA: Kazungula (Jallá); Mt. Chirinda (Swynnerton); Luapula River (Elena d'Aosta). GERMAN EAST AFRICA: Ruanda (Elena d'Aosta); Mt. Karisimbi; southern Mpororo; Isalinio (Schubotz); Usambara (Vossler); New Moschi, Mt. Kilimanjaro, 800 m.; Kilema, 1400-1500 m. (Alluaud and Jeannel); Kibonoto, Mt. Kilimanjaro; Mt. Meru, 3000 m. (Sjöstedt). UGANDA: Unyoro Province, Lake Albert Region, Waki River; Mt. Ruwenzori, eastern slope, 1600 m. (C. Alluaud). BRITISH EAST AFRICA: Mt. Kenia, 2400-2800 m.; Bura Mts., Wa-Taita Province (v. d. Decken; C. Alluaud); Mombasa, Freretown; Naivasha, Rift Valley (Alluaud and Jeannel). ARYSSINIA: Let Marefia, Schoa (Antinori).

255₃. Var. *ornatus* SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, pp. 157 and 161 (♂).

Type locality: CAMEROON.

255₄. Var. *pallidus* SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, pp. 157 and 160, fig. B1 (♂).

Type locality: CAMEROON.

255₅. Var. *rubellus* (SAVAGE) SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 352 (♂); 1912, Ann. Soc. Ent. Belgique, LVI, p. 157, fig. B2 (♂). See p. 47.

Anomma rubella SAVAGE, 1849, Proc. Acad. Nat. Sci. Philadelphia, IV, p. 196 (♂). F. SMITH, 1858, 'Cat. Hym. Brit. Mus.', VI, p. 199 (♂). ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 47 (♂); 1863, 'Verzeich. Formicid.', p. 21. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 395. F. SMITH, 1863, Trans. Ent. Soc. London, (3) I, p. 470 (♂).

Anomma pubescens EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 273.

Anomma burmeisteri var. *rubella* EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 554 (♂). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 9. WASMANN, 1904, Zool. Jahrb. Abt. Syst., Suppl. VII, pp. 672 and 675 (♂); 1916, Ent. Mitt. Deutsch. Ent. Mus. Berlin, V, pp. 95, 96, 97, 102, 104, 137, 140, and 146; 1917, Zeitschr. Wiss. Zool., CXVII, pp. 271, 275, 298, 314, 319, and 321.

Dorylus (Anomma) nigricans var. *rubellus* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, p. 710 (♂); 1899, Ann. Soc. Ent. Belgique, XLIII, p. 462 (♂). Sjöstedt, 1908, 'Exped. Kilimandjaro, Meru., etc.', II, 8, p. 111. EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 12. FOREL, 1911, Rev. Zool. Afr., I, p. 275 (♂). SANTSCHI, *ibid.*, p. 206. FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 348 (♂).

Dorylus nigricans var. *rubellus* Sjöstedt, 1904, 'I Västafrikas Urskogar,' p. 157, footnote.

Dorylus (Anomma) rubellus FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 77 (♂).

Anomma nigricans subsp. *rubella* WASMANN, 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, p. 289.

Type locality: Mpongwe District, FRENCH CONGO (Savage).

CAMEROON: Buea (German Deep Sea Exp.). FRENCH CONGO: Gaboon. BELGIAN CONGO: Lukula (Daniel); Mayombe (de Briey); St. Gabriel near Stanleyville (H. Kohl); Boma (Lang and Chapin). BENGUELA: (J. Cruchet). GERMAN EAST AFRICA: Mafia Island (Voeltzkow).

25c. Subsp. *sjöstedti* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 461 (♀); 1910, 'Gen. Insect., Dorylinae,' p. 12. SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, pp. 156 and 160, figs. A2, B5, and B17 (♀). See p. 48.

Dorylus nigricans var. *sjöstedti* SJÖSTEDT, 1904, 'I Västafrikas Urskogar,' p. 157, footnote; 1908, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 111.

Anomma nigricans subsp. *arcens* MAYR, 1896, Ent. Tidskr., XVII, p. 225 (♀) (nec Westwood).

Anomma nigricans subsp. *sjöstedti* WASMANN, 1904, Zool. Jahrb. Abt. Syst., Suppl. VII, p. 672 (♀). REICHENSPERGER, 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, p. 124.

Dorylus (Anomma) nigricans subsp. *sjöstedti* STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 128 (2, ♀). WASMANN, 1912, Zool. Anzeiger, XXXIX, p. 473.

Anomma sjöstedti WASMANN, 1913, Ann. Rept. Smiths. Inst. for 1912, p. 471. H. SCHMITZ, 1914, Zool. Jahrb. Abt. Syst., XXXVII, p. 540. WASMANN, 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, pp. 31 and 32; 1916, *ibid.*, V, pp. 96, 102, 107, 142, and 145; 1917, Zeitschr. Wiss. Zool., CXVII, pp. 271, 298, 302, 314, 321, and 330.

Anomma sjöstedti H. SCHMITZ, 1917, Zoolog. Meded. Mus. Leiden, III, p. 125.

Type locality: CAMEROON (Sjöstedt).

FERNANDO PO: (Conradt). CAMEROON: (Mansfeld; Zenker); Mundame (Conradt); Molundu (Reichensperger); Victoria (German Deep Sea Exp.); Lolodorf; Grand Batanga (G. Schwab). FRENCH CONGO: Krebedje, southern Darbanda (Decorse). BELGIAN CONGO: Faradje; Medje; Niangara (Lang and Chapin).

25c1. Var. *rufescens* (WASMANN) EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 12.

Anomma nigricans subsp. *sjöstedti* var. *rufescens* WASMANN, 1904, Zool. Jahrb. Abt. Syst., Suppl. VII, p. 673 (♀).

Dorylus (Anomma) nigricans subsp. *burmeisteri* var. *rufescens* SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, pp. 157 and 161, figs. A3, B6, and B18 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 332 (♀).

Dorylus (Anomma) nigricans subsp. *rufescens* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 247, fig. 3 (♀).

Dorylus (Anomma) nigricans subsp. *rufescens* SANTSCHI, 1917, *ibid.*, LXXXV, (1916), p. 278 (♀).

Anomma sjöstedti var. *rufescens* WASMANN, 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, p. 32; 1916, *ibid.*, V, p. 107; 1917, Zeitschr. Wiss. Zool., CXVII, p. 330.

Type locality: Victoria, CAMEROON (German Deep Sea Exp.; F. Silvestri).

CAMEROON: Yukaduma (Funck); Grand Batanga; Lolodorf (G. Schwab). FRENCH CONGO: Samkita (F. Faure).

25c2. Var. *sjöstedti-wilverthi* (WASMANN).

Anomma sjöstedti var. *sjöstedti-wilverthi* WASMANN, 1916, Ent. Mitt. Deutsch. Ent. Mus. Berlin, V, pp. 136 and 146; 1917, Zeitschr. Wiss. Zool., CXVII, pp. 305 and 321 (♀).

Type locality: Grand Batanga, CAMEROON (G. Schwab).

26. *Dorylus (Anomma) stanleyi* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 52 (♂). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 12 (♂). FOREL, 1912, Ent. Mitt. Deutsch. Ent. Mus. Berlin, I, p. 81 (♂). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 63 (♂). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 402 (♂).

?*Dorylus (Anomma) emeryi* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 52 (♂).

Type locality: Katanga, BELGIAN CONGO (Lemaire).

BELGIAN CONGO: (Kohl); Mawambi. BRITISH EAST AFRICA: Molo, Mau Escarpment, 2420 m.; Wambogo, Wa-Kikuyu, 1750 m.; Mt. Kenia, eastern slope, 1900-2000 m. (Alluaud and Jeannel). UGANDA: Unyoro Province, Lake Albert Region (C. Alluaud).

27. *Dorylus (Anomma) wilverthi* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 459, fig. (♀); 1901, Mem. Accad. Sc. Bologna, (5) IX, pp. 417 and 428 (♀), Pl. I, figs. 23-25. WASMANN, 1904, Zool. Jahrb. Abt. Syst., Suppl. VII, p. 674. EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 12 (♀, ♂). FOREL, 1912, Mitt. Naturh. Mus. Hamburg, XXIX, p. 177 (♀, ♀), Pl., figs. 2, 3b, 4b, 5, 9a, and 9b. See p. 48.

Anomma wilverthi WASMANN, 1900, Zool. Jahrb. Abt. Syst., XIV, pp. 257 and 263 (♀); 1904, *ibid.*, Suppl. VII, pp. 660 and 671; 1911, 1er Congr. Intern. Entom. Bruxelles (1910), II, Mém., pp. 227 and 229, Pl. xxv, fig. 23; 1913, Ann. Rept. Smiths. Inst. for 1912, pp. 469 and 471, Pl. vii, fig. 23; 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, pp. 30, 31, 33, 34, 204, and 289; 1916, *ibid.*, V, pp. 94, 95, 97, 98, 101, 104, 107, 108, 137, 140, 142, 143, and 146; 1917, Zeitschr. Wiss. Zool., CXVII, pp. 262, 271, 275, 276, 282, 296, 313, 318, 319, 321, 330, and 353.

Dorylus (Anomma) wilverthi WASMANN, 1900, Zool. Jahrb. Abt. Syst., XIV, p. 274. FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, pp. 51 and 70 (♀, ♂). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 352 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 128 (♀, ♀). SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 155. FOREL, 1913, Rev. Zool. Afr., II, p. 311 (♂); 1913, Ann. Soc. Ent. Belgique, LVII, p. 348 (♀); 1916, Rev. Suisse Zool., XXIV, p. 402 (♂). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 373 (♀).

Anomma nigricans ERN. ANDRÉ, 1900, Bull. Mus. Hist. Nat. Paris, VI, p. 365, 4 figs. (♀) (*nec* Illiger).

Anomma wilverthi WASMANN, 1907, Atti Accad. Pontif. Nuov. Lincei, LX, Sess. 7, pp. 1-6.

Type locality: Leopoldville, BELGIAN CONGO (Wilwerth).

IVORY COAST: OUOSSOU (Talbot). FRENCH CONGO: Brazzaville; Krebedje (A. Weiss). BELGIAN CONGO: Umangi (Wilwerth); Iringui (Lindemann); Mobeka (Lothaire); Kimpoko (Büttner); Ganda Sundi (R. Mayné); Kondué; Sankuru (Luja); St. Gabriel (Kohl); Akenge; Medje; Faradje; Avakubi (Lang and Chapin); Kwesi to Kilo (Bayer); Beni (Borgerhoff); Kimuenza (Schultze); Duma (Schubotz).

Subgenus 3. *Typhlopone* WESTWOOD

Typhlopone WESTWOOD, 1840, 'Introd. Class. Insects,' II, p. 219.

Dorylus subg. *Typhlopone* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 701 and 723; 1910, 'Gen. Insect., Dorylinae,' p. 12.

Cosmaecetes SPINOLA, 1853, Mem. Accad. Sc. Torino, (2) XIII, p. 70.

Subgenotype: *Typhlopone fulva* Westwood, 1840.

28. *Dorylus (Typhlopone) fulvus* (WESTWOOD) EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 707, 709, and 723, figs. P, Q (♀, ♂); 1899, Ann. Mus. Civ. Genova, XXXIX, p. 499 (♀); 1901, Mem. Accad. Sc. Bologna, (5) IX, pp. 416 and 428 (♀); 1910, 'Gen. Insect., Dorylinae,' p. 13. STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 128 (♀).

Dorylus helveticus LATREILLE, 1805, 'Hist. Nat. Crust. Ins.,' XIII, p. 260 (♂, in part), Pl. c, fig. 10 (*nec* LINNÆUS).

Typhlopone fulva WESTWOOD, 1840, 'Introd. Class. Insects,' II, p. 219 (♀); 1840, Ann. Mag. Nat. Hist., VI, p. 87 (♀), Pl. II, figs. 1 and 1a-g. F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 110. ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 45; 1863, 'Verzeich. Formicid.,' p. 20.

Labidus (Typhlopone) kirbii SHUCKARD, 1840, Ann. Nat. Hist., V, p. 265 (♀).

Labidus (Typhlopone) thewaitsi SHUCKARD, *ibid.*, p. 326 (♀).

Labidus (Typhlopone) spinolæ SHUCKARD, *ibid.*, p. 327 (♀).

Dorylus juvenculus SHUCKARD, *ibid.*, p. 318 (♂). WESTWOOD, 1842, 'Arcana Ent.,' I, p. 80 (♂). LUCAS, 1846, 'Explor. Sc. Algérie,' Zool., III, p. 299 (♂). ERN. ANDRÉ, 1882, 'Spec. Hym. Europ. Algérie,' II, p. 254 (♂), Pl. xv, figs. 10-12. EMERY, 1887, Bull. Soc. Ent. Italiana, XIX, p. 350; 1891, 'Explor. Tunisie, Fourmis,' p. 1 (♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 11. WASMANN, 1899, Deutsch. Ent. Zeitschr., p. 175; 1900, Zool. Jahrb. Abt. Syst., XIV, p. 274. BINGHAM, 1903, Ann. Mag. Nat. Hist., (7) XII, p. 61 (♂). WASMANN, 1904, Zool. Jahrb. Abt. Syst., Suppl. VII, p. 676.

?*Dorylus labiatus* SHUCKARD, 1840, Ann. Nat. Hist., V, p. 319 (♂).

Typhlopone shuckardi WESTWOOD, 1840, Ann. Mag. Nat. Hist., VI, p. 88 (♀), Pl. II, fig. 2. F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 111. ROGER, 1863, 'Verzeich. Formicid.,' p. 20.

Typhlopone dahlbomi WESTWOOD, 1840, Ann. Mag. Nat. Hist., VI, p. 88 (♀), Pl. II, figs. 3a-b. F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 111. ROGER, 1863, 'Verzeich. Formicid.,' p. 20.

Typhlopone oraniensis LUCAS, 1846, 'Explor. Sc. Algérie,' Zool., III, p. 302 (♀), Pl. XVI, figs. 11 and 11a-g; 1853, Ann. Soc. Ent. France, (3) I, Bull., pp. xxxviii-xxxix (♀). NYLANDER, 1856, Ann. Sc. Nat. Zool., (4) V, p. 76 (♀), Pl. III, figs. 24-26.

Cosmaecetes homalinus SPINOLA, 1853, Mem. Accad. Sc. Torino, (2) XIII, p. 71 (♀). EMERY, 1893, Boll. Mus. Zool. Anat. Comp. Torino, VIII, No. 163, p. 1.

Typhlopone oraniensis F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 112. MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 737 (♀). ROGER, 1863, 'Verzeich. Formicid.,' p. 20. EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 381, fig. (♀). PASCOE, 1878, Trans. Ent. Soc. London, Proc., p. xxxviii. ERN. ANDRÉ, 1882, 'Spec. Hym. Europ. Algérie,' II, Pl. I, fig. 13 (♀). FOREL, 1885, Journ. Asiat. Soc. Bengal, LIV, p. 177 (♀).

Typhlopone homalina F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 111. ROGER, 1863, 'Verzeich. Formicid.,' p. 20.

Typhlopone spinolæ F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 111. ROGER, 1863, 'Verzeich. Formicid.,' p. 20.

Typhlopone thewaitsi F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 111. ROGER, 1863, 'Verzeich. Formicid.,' p. 20.

Typhlopone spinolæ MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 457.

Typhlopone thewaitsi MAYR, *ibid.*, p. 457.

Typhlopone dahlbomi MAYR, *ibid.*, p. 457.

Typhlopone shuckardi MAYR, *ibid.*, p. 457.

Typhlopone fulva MAYR, *ibid.*, p. 457.

Typhlopone oraniensis MAYR, *ibid.*, p. 457.

Typhlopone homalina MAYR, *ibid.*, p. 457.

Dorylus oraniensis ERN. ANDRÉ, 1882, 'Spec. Hym. Europ. Algérie,' II, p. 252 (♀), Pl. xv, figs. 1-4.

Typhlopone clausii JOSEPH, 1882, Berlin. Ent. Zeitschr., XXVI, p. 47 (♀).

Dorylus clausii ERN. ANDRÉ, 1882, 'Spec. Hym. Europ. Algérie,' II, p. 252 (♀), Pl. xv, fig. 6.

Dorylus fulvus EMERY, 1895, Ann. Soc. Ent. France, LXIV, Bull., p. lxxiii, fig. (♀, ♂); 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 595 (♀, ♂). H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 42. R. GESTRO, in Vanutelli and Citeri, 1899, 'Seconda Spedizione Bottego. L'Omo,' p. 627, fig. 20 (♀, ♂). F. KOHL, 1907, Denkschr. Ak. Wiss. Wien, LXXI, p. 281 (♂). MAYR, 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. I, p. 387. ZAVATARRI, 1909, 'Il Ruwenzori, Parte Scientific,' I, p. 212 (♂). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 4 (♂). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 333 (♂).

Dorylus fulvus fulvus EMERY, 1919, Ann. Soc. Ent. Belgique, LIX, p. 102.

Type locality: unknown, "in saccharo detecta."

SENEGAMBIA: Thiès (F. Silvestri). CAMEROON: Mundame (Conradt). ANGLO-EGYPTIAN SUDAN: Khartum (F. Werner); Sennar. ABYSSINIA: Ogaden; Bela (Ruspoli); Lakes Bass Marle and Bass Narok (V. Bottego). SOMALILAND: Lugh (Ruspoli). ZANZIBAR: (Stuhlmann). SOUTHERN ARABIA: Makalla (O. Simony). SINAI PENINSULA. SYRIA. NORTH AFRICA. Doubtful record: Pretoria, TRANSVAAL (Wichgraf).

28a. Subsp. *badius* (GERSTÄCKER) EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 13. STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exped. (1907-08),' III, p. 375 (♂). FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 362 (♂); 1913, Ann. Soc. Ent. Belgique, LVII, pp. 111 (♀, ♂) and 350 (♂); 1913, Rev. Zool. Afr., II, p. 312 (♂). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 63 (♀, ♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 125 (♀, ♂), Pl. iv, figs. 31, 31a, 32, 32a, and 32b. EMERY, 1919, Ann. Soc. Ent. Belgique, LIX, p. 102. WHEELER, 1921, Proc. American Ac. Arts Sci. Boston, LVI, p. 303, fig. 5a (♂).

Dorylus badius GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 261 (♂); in Peters, 1862, 'Reise n. Mossambique,' Zool., V, p. 499 (♂), Pl. xxxi, fig. 14. ROGER, 1863, 'Verzeich. Formicid.,' p. 41. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 408. GERSTÄCKER, 1872, Stettin. Ent. Zeitg., XXXIII, p. 257 (♀, ♂).

Dorylus juvenculus var. *badius* EMERY, 1894, Mem. Accad. Sc. Bologna, (5) IV, p. 112 (♂). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 4 (♂).

Dorylus (*Typhlopone*) *fulvus* var. *badius* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 709 and 724, figs. P and R (♀). MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 8 (♂). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 52 (♂).

Type locality: MOZAMBIQUE (Peters).

GOLD COAST: Akra (Unger). BELGIAN CONGO: Kondué, Kasai (Luja); Lukula (Daniel); Kabambare (Flamand); Volcano region N. E. of Lake Kivu (Schubotz). RHODESIA: Bulawayo (G. Arnold). GERMAN SOUTHWEST AFRICA: Windhoek (Lübbert); Okahandja (Casper; Peters); Ontys (Langheld). BECHUANALAND: between Severelela and Khakhea (L. Schultze). CAPE PROVINCE: Cape Town (Wilms). MOZAMBIQUE: Inhambane (Fornasini). GERMAN EAST AFRICA: Tanga; New

Moschi, Mt. Kilimanjaro (Alluaud and Jeannel); Kibonoto, Kilimanjaro and Masai Steppe (Sjöstedt). BRITISH EAST AFRICA: Mt. Kenia, Meranga; Fort Hall, Nairobi; Mombasa Island (Alluaud and Jeannel). UGANDA: Unyoro Province near Hoima, east of Lake Albert (C. Alluaud); Lake Victoria (Zimmer); Entebbe (Duke of Abruzzi).

28a₁. Var. *eurossus* EMERY, 1919, Ann. Soc. Ent. Belgique, LIX, p. 102.

Dorylus fulvus subsp. *eurossus* EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 5, figs. 1 and 2 (♂, ♂).

Type locality: Keren, ERITREA (F. Silvestri).

ERITREA: Ghinda; Nefasit (F. Silvestri).

28a₂. Var. *obscurior* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 333 (♀). EMERY, 1919, Ann. Soc. Ent. Belgique, LIX, p. 102. See p. 49.

Type locality: Konakry, FRENCH GUINEA (F. Silvestri).

BELGIAN CONGO: Vankerkhovenville; Paradje; Garamba; Batama; Stanleyville (Lang and Chapin); Avakubi (Boyton).

28b. Subsp. *dentifrons* WASMANN, 1904, Zool. Jahrb. Abt. Syst., Suppl. VII, p. 674 (♀). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 13 (♀); 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 5, fig. 1 (♂); 1919, Ann. Soc. Ent. Belgique, LIX, p. 102.

Dorylus fulvus var. *stramineus* STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 128 (♀, ♀).

Type locality: Sankuru, BELGIAN CONGO (E. Luja).

CAMEROON: (Zenker).

28c. Subsp. *glabratus* (SHUCKARD) EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 13 (♂).

Dorylus glabratus SHUCKARD, 1840, Ann. Nat. Hist., V, p. 317 (♂). WESTWOOD, 1842, 'Arcana Ent.,' I, p. 79 (♂). F. SMITH, 1859, 'Cat. Hym. Brit. Mus.,' VII, p. 2 (♂), Pl. I, fig. 8. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 408. RADOSZ-KOWSKY, 1876, Hor. Soc. Ent. Rossicae, XII, p. 140. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 11. MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 200 (♂).

Dorylus (*Typhlopone*) *fulvus* ? var. *glabratus* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 709 and 724.

Type locality: GAMBIA.

EGYPT. ZANZIBAR: (Stuhlmann).

28d. Subsp. *rhodesiae* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 111 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 129 (♀). EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 5, fig. 1 (♀); 1919, Ann. Soc. Ent. Belgique, LIX, p. 102.

Dorylus (*Typhlopone*) *fulvus* subsp. *rhodesianus* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 333 (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

TRANSVAAL: Pretoria (F. Silvestri).

Subgenus 4. *Rhogmus* SHUCKARD

Rhogmus SHUCKARD, 1840, Ann. Nat. Hist., V, p. 323.

Dorylus subg. *Rhogmus* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 702 and 736; 1910, 'Gen. Insect., Dorylinae,' p. 13.

Subgenotype: *Rhogmus fimbriatus* Shuckard, 1840.

29. **Dorylus (Rhogmus) fimbriatus** (SHUCKARD) EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 709 and 736, figs. AA and BB (♀, ♂); 1899, Ann. Mus. Civ. Genova, XXXIX, p. 499 (♂); 1901, Mem. Accad. Sc. Bologna, (5) IX, pp. 416, 419, and 428 (♀), Pl. I, figs. 31-38. FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 47 (♂). H. BRAUNS, 1903, Zeitschr. f. Hym. u. Dipt., III, pp. 294-298, fig. (♀). MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro-Meru, etc.', II, 8, p. 8 (♂). FOREL, 1907, Mitt. Naturh. Mus. Hamburg, XXIV, p. 15 (♂); 1909, Ann. Soc. Ent. Belgique, LIII, pp. 52 and 71 (♂). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 14 (♀, ♀, ♂). FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 362 (♂). SANTSCHI, 1911, Rev. Zool. Afr., I, p. 207 (♀). STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 375 (♂). FOREL, 1912, Ent. Mitt. Deutsch. Ent. Mus. Berlin, I, p. 81 (♂). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 333 (♀, ♂); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 65 (♂). ARNÖLD, 1915, Ann. South African Mus., XIV, p. 129 (♀, ♀, ♀, ♂), Pl. IV, figs. 36, 36a. SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 23 (♂).

Rhogmus fimbriatus SHUCKARD, 1840, Ann. Nat. Hist., V, p. 325 (♂). WESTWOOD, 1842, 'Arcana Ent.,' I, p. 80 (♂). F. SMITH, 1859, 'Cat. Hym. Brit. Mus.,' VII, p. 4 (♂), Pl. I, fig. 2. ROGER, 1863, 'Verzeich. Formicid.,' p. 42. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 453. GERSTÄCKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 355; 1873, in 'v. d. Decken's Reisen in Ostafrika, Gliederthiere,' p. 347 (♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 8. FOREL, 1904, Ann. Mus. Zool. Ac. Sc. St. Pétersbourg, VIII, (1903), p. 369 (♂).

Dorylus fimbriatus H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 42. MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 2 (♀). BINGHAM, 1903, Ann. Mag. Nat. Hist., (7) XII, p. 61 (♂). POULTON, in A. J. Hayes, 1905, 'The Source of the Blue Nile,' p. 314 (♂).

Type locality: GAMBIA.

FRENCH GUINEA: Konakry; Mamou (F. Silvestri). GOLD COAST. CAMEROON. FRENCH CONGO: Gaboon. BELGIAN CONGO: Sankuru (Luja); Katanga (Lemaire). BENGUELA: Cucala (J. Cruchet). SOUTHERN RHODESIA: Salisbury (G. Marshall). NYASALAND: Fort Johnston (Rendall). GERMAN EAST AFRICA: Mpororo; Marangu (Zimmer); Tanga, Caves of Kulumuzi (Alluaud); Kibonoto, Mt. Kilimanjaro, 1000-1900 m. (Sjöstedt). ZANZIBAR: (v. d. Decken). BRITISH EAST AFRICA: Molo, Mau Escarpment, 2400 m.; Wambogo, Wa-Kikuyu, 1750 m.; Nairobi; Mt. Kenia, N'daika, 2200 m. (Alluaud and Jeannel). UGANDA: Ubyoro Province, east of Lake Albert (C. Alluaud). ANGLO-EGYPTIAN SUDAN: Kaka, White Nile (I. Trägårdh); Lake Tsano (A. J. Hayes). SOMALILAND. ABYSSINIA: Buditu to Dimé (V. Bottego); Adis-Abeba (Kachovskij).

29a. Var. **crampeli** SANTSCHI, 1919, Rev. Zool. Afr., VI, pp. 231 and 232 (♂).

Type locality: Fort Crampel, FRENCH CONGO.

29b. Var. **laevipodex** SANTSCHI, 1919, Rev. Zool. Afr., VI, pp. 231 and 232 (♂).

Type locality: BRITISH EAST AFRICA.

29c. Var. **poweri** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 217 (♀).

Type locality: Kimberley, CAPE COLONY (Power).

30. **Dorylus (Rhogmus) fuscipennis** EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 709 and 737, figs. CC and DD (♂); 1901, Mem. Accad. Sc. Bologna, (5) IX, pp. 416 and 428 (♀); 1910, 'Gen. Insect., Dorylinae,' p. 14 (♂). FOREL, 1910, Notes Leyden Mus., XXXI, p. 224, footnote (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 349 (♀). SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 231 (♂).

Rhogmus fuscipennis EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 570 (♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 8.

Type locality: WEST AFRICA.

Haitifi. GOLD COAST: Aburi (Fisch). FRENCH CONGO: Ogowé (Mocquers).

30. Var. *lugubris* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 231 (♂); 1919, *ibid.*, VII, p. 91.

Dorylus (Rhogmus) fimbriatus var. *lugubris* SANTSCHI, 1919, *ibid.*, VI, p. 232 (♂).

Type locality: Aburi, GOLD COAST (Fisch).

30a. Var. *marginiventris* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 231 (♂); 1919, *ibid.*, VII, p. 91.

Dorylus (Rhogmus) fimbriatus var. *marginiventris* SANTSCHI, 1919, *ibid.*, VI, p. 233 (♂).

Type locality: Dimbroko, IVORY COAST (Le Mout).

31. *Dorylus (Rhogmus) leo* SANTSCHI, 1919, Rev. Zool. Afr., VI, pp. 231 and 232 (♂).

Type locality: Dimbroko, IVORY COAST.

32. *Dorylus (Rhogmus) ocellatus* (STITZ).

Dichthadia sp. STITZ, 1909, Zool. Anzeiger, XXXV, p. 231, figs. 1-3 (♀).

WASMANN, 1911, Rev. Zool. Afr., I, p. 113.

Dichthadia ocellata STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 128 (♀).

Type locality: Bipindi, CAMEROON (Zenker).

33. *Dorylus (Rhogmus) savagei* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 709 and 738, figs. EE and FF (♂). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 52 (♂). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 14 (♂). FOREL, 1911, Rev. Zool. Afr., I, p. 275 (♂); 1913, Ann. Soc. Ent. Belgique, LVII, p. 348 (♂). SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 231 (♂).

Rhogmus fuscipennis ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 5 (♂) (*nec* Emery).

Type locality: Gaboon, FRENCH CONGO.

BELGIAN CONGO: Lukula (Daniel); Mayombe (Deleval); Tshoa (Cabra).

33a. Var. *mucronatus* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 462 (♂); 1910, 'Gen. Insect., Dorylinae,' p. 14 (♂). SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 231 (♂).

Dorylus (Rhogmus) savagei subsp. *mucronatus* FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 47 (♂).

Type locality: Lagos, NIGERIA.

NIGERIA: Wari, Forcados River (Henry Fischer).

34. *Dorylus (Rhogmus) termitarius* WASMANN, 1911, Rev. Zool. Afr., I, p. 111 (♂, ♀), Pl. I, fig. 4.

Type locality: Romée near Stanleyville, BELGIAN CONGO (H. Kohl).

Subgenus 5. *Alaopone* EMERY

Alaopone EMERY (part), 1881, Ann. Mus. Civ. Genova, XVI, p. 274.

Dorylus subg. *Alaopone* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 702 and 730; 1910, 'Gen. Insect., Dorylinae,' p. 14.

Dorylus subg. *Shuckardia* (part) EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 703 and 740.

Subgenotype: *Alaopone oberthüri* Emery, 1881 = *Dorylus orientalis* Westwood, 1835.

35. *Dorylus (Alaopone) atriceps* SHUCKARD. EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 15 (♂). SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 362 (♂). See p. 51.

Dorylus atriceps SHUCKARD, 1840, Ann. Nat. Hist., V, p. 323 (♂). WESTWOOD, 1842, 'Arcana Ent.,' I, p. 80 (♂). F. SMITH, 1859, 'Cat. Hym. Brit. Mus.,' VII, p. 4 (♂), Pl. I, fig. 14. ROGER, 1863, 'Verzeich. Formicid.,' p. 41. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 408. RADOSZKOWSKY, 1881, Journ. Sci. Ac. Lisboa, VIII, No. 31, p. 198. ERN. ANDRÉ, 1886, 'Spec. Hym. Europ. Algérie,' II, p. 840 (♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 10. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 5 (♂).

Dorylus shuckardi RITSEMA, 1874, Tijdschr. v. Ent., XVII, p. 182 (♂).

Dorylus ritsemæ DALLA TORRE, 1892, Wien. Ent. Zeitg., XI, p. 89 (♂).

Dorylus atriceps var. *shuckardi* DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 10.

Dorylus (Shuckardia) atriceps EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 710 and 740, figs. HH and GG (♂). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 53 (♂).

Type locality: Gambia River, GAMBIA.

SENEGAMBIA. GUINEA. IVORY COAST: Dimbroko (Le Moul). FRENCH CONGO: Ogowe (Mocquerys). BELGIAN CONGO: Vivi; Chikai (Cabra); Faradje; Stanleyville (Lang and Chapin). ANGOLA: (Welwitsch). ABYSSINIA. TUNIS.

35a. Var. *katanensis* STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 375 (♂).

Type locality: Plain south of Lake Albert Edward, BELGIAN CONGO (Schubotz).

35a. Subsp. *æthiopicus* EMERY. FOREL, 1907, Ann. Soc. Ent. Belgique, LI, p. 201 (♀). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 15 (♀, ♂).

Dorylus atriceps EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 381 (♂) (nec Shuckard).

Dorylus (Shuckardia) atriceps subsp. *æthiopicus* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 710 and 741, figs. JJ and KK (♂). SANTSCHI, 1907, Rev. Suisse Zool., XV, p. 327 (♀).

Type locality: Sudan.

ERITREA: Anseba (Beccari). TUNIS: Kairouan (F. Santschi).

36. *Dorylus (Alaopone) attenuatus* SHUCKARD. EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 709 and 733, figs. X, Y (♂); 1910, 'Gen. Insect., Dorylinae,' p. 15 (♂). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 4 (♂); 1913, Rev. Zool. Afr., II, p. 312 (♂). ARNOLD, 1915, (misspelled *attenatus*) Ann. South African Mus., XIV, p. 133 (♂).

Dorylus attenuatus SHUCKARD, 1840, Ann. Nat. Hist., V, p. 322 (♂). WESTWOOD, 1842, 'Arcana Ent.,' I, p. 80 (♂). F. SMITH, 1859, 'Cat. Hym. Brit. Mus.,' VII, p. 3 (♂), Pl. I, fig. 13. ROGER, 1863, 'Verzeich. Formicid.,' p. 41. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 408. RADOSZKOWSKY, 1876, Hor. Soc. Ent. Rossica, XII, p. 140. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 10. BINGHAM, 1903, Ann. Mag. Nat. Hist., (7) XII, p. 61 (♂).

Type locality: Gambia River, GAMBIA.

BELGIAN CONGO: Valley of the Lubumbashi (Buttgenbach). GERMAN SOUTH-WEST AFRICA: (Lübbert). BECHUANALAND: Lobatsi, north of Mafeking (L. Schultze). CAPE PROVINCE: Cape of Good Hope. TRANSVAAL: Pretoria (Distant). ABYSSINIA: (Raffray).

36i. Var. *acuminatus* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 462 (♂). MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 8 (♂). EMERY, 1910, 'Gen. Insect., Dorylinæ,' p. 15 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 133 (♂).

Type locality: ORANGE FREE STATE.

GERMAN EAST AFRICA: Kibondo, Mt. Kilimanjaro, 1000-1200 m. (Sjöstedt).

36e. Var. *umbratipennis* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 52 (♂). EMERY, 1910, 'Gen. Insect., Dorylinæ,' p. 15 (♂).

Type locality: Nini (? probably Vivi, BELGIAN CONGO).

36a. Subsp. *latinodis* FOREL, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 479 (♂).

Type locality: Stanleyville, BELGIAN CONGO (H. Kohl).

37. *Dorylus (Alaopone) buyssoni* SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 748, figs. 7a, b, d, and e (♂); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 65 (♂).

Type locality: Nairobi, Wa-Kikuyu, BRITISH EAST AFRICA (C. Alluaud).

BRITISH EAST AFRICA: Cheteni (Alluaud and Jeannel).

37i. Var. *conjugens* SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 750, figs. 7c and f (♂); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 65 (♂).

Type locality: Mombasa, Athi Plains, BRITISH EAST AFRICA (Fernique).

BRITISH EAST AFRICA: Mwatate, Wa-Taita Province (C. Alluaud).

38. *Dorylus (Alaopone) conradti* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 707 and 734, fig. Z (♀, ♀), Pl. XVI, figs. 1-4, and Pl. XVII, figs. 7-10; 1901, Mem. Accad. Sc. Bologna, (5) IX, p. 428 (♀); 1910, 'Gen. Insect., Dorylinæ,' p. 15 (♀, ♀). WASMANN, 1918, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VII, p. 136. See p. 51.

Type locality: Bismarckburg, Togo (Conradt).

CAMEROON: Soppo (v. Rothkirch). BELGIAN CONGO: Niangara (Lang and Chapin).

39. *Dorylus (Alaopone) diadema* GERSTÄCKER. EMERY, 1910, 'Gen. Insect., Dorylinæ,' p. 15 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 134 (♂).

Dorylus diadema GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 261 (♂); in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 500 (♂), Pl. XXXI, fig. 15. ROGER, 1863, 'Verzeich. Formicid.,' p. 41. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 408. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 10. FOREL, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 309 (♂).

Dorylus (Shuckardia) diadema EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 710 and 742, figs. MM and LL (♂).

Type locality: Tete, MOZAMBIQUE (Peters).

39i. Var. *fusciceps* EMERY, 1910, 'Gen. Insect., Dorylinæ,' p. 15 (♂).

Dorylus (Shuckardia) diadema var. *fusciceps* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 463 (♂).

Type locality: Lake Nyasa.

CONGO.

39a. Subsp. **arnoldi** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 216 (♂).

Type locality: Bulawayo, RHODESIA (G. Arnold).

40. **Dorylus (Alaopone) distinctus** SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 747, fig. 6 (♂).

Type locality: FRENCH GUINEA (Marchand).

41. **Dorylus (Alaopone) montanus** SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 750, fig. 8 (♂); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 66 (♀).

Type locality: Kibosho, Mt. Kilimanjaro, 1400 m., GERMAN EAST AFRICA (C. Alluaud).

GERMAN EAST AFRICA: New Moschi, 800 m.; Kilema, 1440 m. (Alluaud and Jeannel).

41a. Var. **bondroiti** SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 162 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 135 (♂).

Dorylus (Alaopone) montanus subsp. *bondroiti* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 248 (♂).

Type locality: TRANSVAAL.

CAPE PROVINCE: Steynsburg (Ellenberger).

41a. Subsp. **australis** SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 231 (♀).

Dorylus (Alaopone) australis SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 362 (♀).

Type locality: Majuba, Herschel District, CAPE PROVINCE (G. Arnold).

41b. Subsp. **brevis** SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 245, fig. 1 (♂).

Type locality: Wombali, BELGIAN CONGO (Vanderyst).

Ecitonini Forel

Enictogiton EMERY

Enictogiton EMERY, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 49; 1910, 'Gen. Insect., Dorylinæ,' p. 27.

Genotype: *Enictogiton fossiceps* Emery, 1901.

1. **Enictogiton bequaerti** FOREL, 1913, Rev. Zool. Afr., II, p. 314 (♂).

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

2. **Enictogiton elongatus** (SANTSCHI).

Enictogiton elongatus SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 246, fig. 2c (♂).

Type locality: Malela, BELGIAN CONGO (Burgeon).

2a. Var. **attenuatus** SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 247, fig. 2b (♂).

Type locality: 300 km. south of Kindu, BELGIAN CONGO (Burgeon).

3. **Enictogiton emeryi** FOREL, 1913, Rev. Zool. Afr., II, p. 315 (♂).

Type locality: Valley of the Lubumbashi, BELGIAN CONGO (Buttgenbach).

4. **Enictogiton fossiceps** EMERY, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 49, figs. 3 and 4 (♂); 1910, 'Gen. Insect., Dorylinæ,' p. 28 (♂).

Type locality: Congo.

5. **Enictogiton sulcatus** (SANTSCHI).

Enictogiton sulcatus SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 247, fig. 2a (♂).

Type locality: Katakai, BELGIAN CONGO (Gérard).

***Enictus* SHUCKARD¹**

Enictus SHUCKARD, 1840, Ann. Nat. Hist., V, p. 266 (♂). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 28.

Typhlatta F. SMITH, 1858, Journ. Linn. Soc. London, Zool., II, p. 79 (♀).

Eciton (part) F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 153.

Alaopone (part) ERN. ANDRÉ, 1886, 'Spec. Hym. Europ. Algérie,' II, p. 854.

Dorylus subg. *Shuckardia* (part) EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, pp. 703 and 740.

Genotype: *Enictus ambiguus* Shuckard, 1840.

1. *Enictus asperivalvus* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 233 (♂); 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 373, fig. 10 (♂).

Type locality: Dimbroko, IVORY COAST (Le Moult).

2. *Enictus batesi* FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 255 (♀).

Type locality: Old Calabar, NIGERIA (Bates).

3. *Enictus bottegoi* EMERY, 1899, Ann. Mus. Civ. Genova, XXXIX, p. 499, fig. (♂); 1910, 'Gen. Insect., Dorylinae,' p. 31 (♂).

Type locality: Lake Bass Narok, ABYSSINIA (V. Bottego).

3a. Var. *alluaudi* SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 754, fig. 10 (♂); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 67 (♂).

Type locality: Kisumu, Victoria Nyanza, BRITISH EAST AFRICA (C. Alluaud).

BRITISH EAST AFRICA: Nairobi, Wa-Kikuyu; Athi Basin, Ndarugo River (C. Alluaud).

3a. Subsp. *anceps* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 249 (♂).

Enictus anceps SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 374, fig. 11 (♂).

Type locality: Nefasit, ERITREA (K. Escherich).

ERITREA: Ghinda (K. Escherich).

3b. Subsp. *noctivagus* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 303, fig. 1 (♂).

Type locality: ABYSSINIA.

4. *Enictus buttgenbachi* FOREL, 1913, Rev. Zool. Afr., II, p. 313 (♂). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 68 (♀, ♂).

Type locality: Valley of the Lubumbashi, BELGIAN CONGO (Buttgenbach).

GERMAN EAST AFRICA: Kilema, Mt. Kilimanjaro, 1440 m.; New Moschi, 800 m. (Alluaud and Jeannel).

5. *Enictus congolensis* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 277 (♀, ♀), Pl. I.

Enictus rixator var. *congolensis* SANTSCHI, 1911, Rev. Zool. Afr., I, p. 207 (♀).

Type locality: FRENCH CONGO (A. Weiss).

FRENCH CONGO: Lambarene (F. Faure).

6. *Enictus crucifer* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 68, fig. 5 (♂).

Type locality: Tiwi, BRITISH EAST AFRICA (Alluaud and Jeannel).

7. *Enictus decolor* (MAYR) DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 7. EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, p. 746 (♀); 1910, 'Gen. Insect., Dorylinae,' p. 31 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 143 (♀).

¹*Enictus unicolor* "Smith," recorded from Angola (Welwitsch) by Radoszkowsky, Journ. Sci. Ac. Lisbon, VIII, No. 31, 1881, p. 198, has apparently never been described.

Typhlatta decolor MAYR, 1878, Verh. Zool. Bot. Ges. Wien, XXVIII, p. 668 (♀).

Type locality: East Africa.

8. *Enictus eugenii* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 17 (♀), Pl. II, figs. 1-4; 1895, Zool. Jahrb. Abt. Syst., VIII, p. 747 (♀). WASMANN, 1900, ibid., XIV, pp. 268 and 275. H. BRAUNS, 1901, Zeitschr. f. Hym. u. Dipt., I, p. 17. WASMANN, 1904, Zool. Jahrb. Abt. Syst., Suppl. VII, p. 676. EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 31 (♀). FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 255 (♀). WASMANN, 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, pp. 28 and 203.

Enictus eugeniae ARNOLD, 1915, Ann. South African Mus., XIV, p. 139 (♀), Pl. IV, fig. 38.

Type locality: Makapan, TRANSVAAL (E. Simon).

ORANGE FREE STATE: Bothaville (H. Brauns). SOUTHERN RHODESIA: Bulawayo (G. Arnold).

8a. Var. *brazzai* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 355 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

8a. Subsp. *caroli* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 248 (♀). EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 3 (♀).

Type locality: Nefasit, ERITREA (K. Escherich).

ERITREA: Ghinda (F. Silvestri).

9. *Enictus foreli* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 234 (♂); 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 369 (♂).

Type locality: Dimbroko, IVORY COAST (Le Moutt).

10. *Enictus furculatus* SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 91; 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 373, fig. 10 (♂).

Enictus furculatus SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 233 (♂).

Type locality: Saint Louis, SENEGAMBIA (Le Moutt).

11. *Enictus fuscovarius* GERSTAECKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 262 (♂); in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 501 (♂), Pl. XXXII, fig. 1. ROGER, 1863, 'Verzeich. Formicid.,' p. 41. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 394. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 7. EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, p. 747 (♂); 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 597 (♂). FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 249 (♂). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 31 (♂). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 350 (♂). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 67 (♂).

Type locality: Tete, MOZAMBIQUE (Peters).

BELGIAN CONGO: Sainpwe (J. Bequaert). BRITISH EAST AFRICA: Voi, Taita, 600 m.; Taveta, 750 m.; Landjoro, Pori, 900 m. (Alluaud and Jeannel). ABYSSINIA: Ganale; Lake Abaja; Bela; Magala Re Umberto; Webi (Ruspoli). ERITREA: Ghinda (K. Escherich).

11a. Var. *magretti* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, p. 747 (♂). MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 2 (♂); 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. I, p. 387. EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 31. SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 756 (♂); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 67 (♂).

Enictus inconspicuus MAGRETTI, 1884, Ann. Mus. Civ. Genova, XXI, p. 539 (♂) (nec Westwood); 1884, Bull. Soc. Ent. Italiana, XV, (1883) p. 244 (♂).

Enictus magretti EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 569 (♂), Pl. xv, figs. 13, 14; 1892, Ann. Mus. Civ. Genova, XXXII, p. 110, fig. (♂). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 8.

Type locality: ANGLO-EGYPTIAN SUDAN (Magretti).

SENEGAMBIA: Gorée Island (H. Brauns). BRITISH EAST AFRICA: Taveta; Bura, Wa-Taita Province (Alluaud and Jeannel). SOMALILAND: Eter-es-Saghir (Bricchetti-Robecchi). ABYSSINIA: Diré Dava; Bazen (Magretti). ERITREA: Sogodas (Magretti). ANGLO-EGYPTIAN SUDAN: White Nile (I. Trägårdh); Khartum (F. Werner); Kor Guillo; Bahr-el-Salaam; El Hefera (Magretti).

12. *Enictus hamifer* EMERY, 1896, Ann. Mus. Civ. Genova, XXXVII, p. 153, fig. (♂); 1910, 'Gen. Insect., Dorylinae,' p. 31 (♂). FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 250 (♂); 1912, Ent. Mitt. Deutsch. Ent. Mus. Berlin, I, p. 81 (♂). SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 756 (♂).

Type locality: Maddo Wells, BRITISH EAST AFRICA (V. Bottego).

ABYSSINIA: Confluence of the Webi and Ganale Rivers (V. Bottego); Diré Dava (Magretti). ERITREA: Ghinda (K. Escherich). BRITISH EAST AFRICA.

13. *Enictus humeralis* SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 752, figs. 9a-e (♂).

Type locality: Bamako, FRENCH SUDAN (Chevalier).

13i. Var. *chevalieri* SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 754 (♂).

Type locality: CASAMANCE (Chevalier).

13e. Var. *viridans* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 248 (♂).

Type locality: Kouandé, UPPER DAHOMEY (Desanti).

14. *Enictus inconspicuus* WESTWOOD, 1847, Trans. Ent. Soc. London, IV, p. 238 (♂), Pl. xiv, fig. 4. F. SMITH, 1859, 'Cat. Hym. Brit. Mus.,' VII, p. 10 (♂), Pl. II, figs. 15a-c. ROGER, 1863, 'Verzeich. Formicid.,' p. 41. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 394. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 8. EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, p. 747 (♂); 1910, 'Gen. Insect., Dorylinae,' p. 31 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 141 (♂).

Type locality: South Africa (Drège).

15. *Enictus luteus* EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 568 (♂), Pl. xv, figs. 11 and 12. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 8. EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, p. 747 (♂); 1910, 'Gen. Insect., Dorylinae,' p. 32 (♂). SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 373 (♂).

Type locality: SIERRA LEONE.

IVORY COAST: Dimbroko (Le Mout).

16. *Enictus mariae* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 18 (♀), Pl. II, figs. 5-7; 1895, Zool. Jahrb. Abt. Syst., VIII, p. 747 (♀); 1910, 'Gen. Insect., Dorylinae,' p. 31 (♀). FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 212 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 140 (♀).

Type locality: Makapan, TRANSVAAL (E. Simon).

RHODESIA: Bulawayo (G. Arnold).

16i. Var. *natalensis* FOREL, in Emery, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 49 (♀). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 31 (♀). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 14 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 141 (♀).

Type locality: NATAL (Haviland).

ZULULAND: Dukudu (I. Trägårdh).

17. *Enictus möbii* EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, p. 747, fig. NN (♂); 1910, 'Gen. Insect., Dorylinae,' p. 32 (♂).
Enictus möbii SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 269, fig. 6 (♂).
 Type locality: Bismarckburg, Togo (Büttner).
 IVORY COAST: Dimbroko (Le Moul't).
 17a. Subsp. *mutatus* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 304, fig. 2 (♂).
Enictus möbii var. SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 754 (♂).
Enictus mutatus SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 370, fig. 7 (♂).
 Type locality: IVORY COAST (Delafosse).
 IVORY COAST: Dimbroko (Le Moul't). DAHOMEY: Kouandé (Desanti).
 17b. Subsp. *pudicus* (SANTSCHI).
Enictus mutatus subsp. *pudicus* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 234 (♂); 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 371, fig. 8 (♂).
 Type locality: Dimbroko, IVORY COAST (Le Moul't).
 17c. Subsp. *sankisianus* FOREL, 1913, Rev. Zool. Afr., II, p. 312 (♂).
 Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).
 18. *Enictus raptor* FOREL, 1913, Rev. Zool. Afr., II, p. 314 (♂).
 Type locality: Lubumbashi Valley, BELGIAN CONGO (Buttgenbach).
 19. *Enictus rixator* FOREL, in Emery, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 48 (♀). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 31 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 137 (♀).
 Type locality: NATAL (R. C. Wroughton).
 20. *Enictus rotundatus* MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 1 (♀). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 31 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 112 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 138 (♀), Pl. IV, fig. 37.
 Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).
 RHODESIA: Bulawayo (G. Arnold).
 21. *Enictus soudanicus* SANTSCHI, 1910, Rev. Suisse Zool., XVIII, p. 758, fig. 12 (♂); 1913, Ann. Soc. Ent. Belgique, LVII, p. 304 (♀).
 Type locality: Toukola, SUDAN (Conan).
 SENEGAMBIA: St. Louis (Claveau).
 21a. Var. *brunneus* FOREL, 1913, Rev. Zool. Afr., II, p. 313 (♂).
 Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).
 22. *Enictus steindachneri* MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 2 (♀). EMERY, 1910, 'Gen. Insect., Dorylinae,' p. 31 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 69 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 139 (♀).
 Type locality: Reddersburg, ORANGE FREE STATE (H. Brauns).
 BRITISH EAST AFRICA: Mbuyuni, Pori, 1110 m. (Alluaud and Jeannel).
 23. *Enictus togoënsis* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 248 (♂); 1920, ibid., LXXXVIII, (1919), p. 375, fig. 12 (♂).
 Type locality: Togo.
 IVORY COAST: Dimbroko (Le Moul't).

24. *Enictus tuberculatus* ARNOLD, 1915, Ann. South African Mus., XIV, p. 142 (♂).

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

25. *Enictus weissi* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 354, fig. 2 (♀). FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 212 (♀). SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 9 (♀).

Type locality: Gomba, FRENCH CONGO (A. Weiss).

CAMEROON: Grand Batanga (G. Schwab). FRENCH CONGO: Samkita (F. Faure).

CERAPACHYINÆ Forel

Cerapachyini Forel

Sphinctomyrmex MAYR

Sphinctomyrmex MAYR, 1866, Verh. Zool. Bot. Ges. Wien, XVI, p. 895. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 7.

Cerapachys (part) FOREL, in Grandidier, 1892, 'Hist. Phys. Nat. Madagascar,' XX, 2, p. 243.

Genotype: *Sphinctomyrmex stali* Mayr, 1866.

The two following African forms are probably not true members of this genus; being described from the male alone they cannot be properly classified, since no male of the genus is known with certainty.

1. *Sphinctomyrmex rufiventris* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 244, fig. 1 (♂).

Type locality: Djougou, Kouandé, UPPER DAHOMY (Desanti).

UPPER DAHOMY: Kika (Ribot). FRENCH SUDAN: Sikosso (Chevalier).

1a. Subsp. *chariensis* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 245 (♂).

Type locality: Fort Archambault, FRENCH EQUATORIAL AFRICA (Decorse).

Cerapachys F. SMITH

Cerapachys F. SMITH, 1858, Journ. Linn. Soc. London, Zool., II, p. 74. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 8.

Ceratopachys W. A. SCHULZ, 1906, 'Spolia Hymenopterologica,' p. 155.

Genotype: *Cerapachys antennatus* F. Smith, 1858.

The African species all belong to the subgenus *Cerapachys*, *sensu stricto*.

1. *Cerapachys afer* FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 9 (♀, ♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 9.

Type locality: Mto-ya-Kifaru, Arusha-chini, GERMAN EAST AFRICA (Katona).

2. *Cerapachys arnoldi* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 211 (♀).

Type locality: CAPE COLONY.

3. *Cerapachys cooperi* ARNOLD, 1915, Ann. South African Mus., XIV, p. 14 (♀, ♀).

Type locality: Umgeni, Durban, NATAL (C. B. Cooper).

4. *Cerapachys cribrinodis* EMERY, 1902, Rend. Accad. Sc. Bologna, N. S., VI, p. 24; 1911, 'Gen. Insect., Ponerinae,' p. 9 (♀). FOREL, 1913, Rev. Suisse Zool., XXI, p. 667 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 12 (♀). See p. 53.

Parasyscia cribrinodis EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 463, fig. (♂). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 128 (♂).

Type locality: CAMEROON (Conradt).

CAMEROON: Johann-Albrechtshöhe (Conradt). BELGIAN CONGO: Medje (Lang and Chapin).

4. Var. *natalensis* FOREL, 1901, Rev. Suisse Zool., IX, p. 335 (♂); 1910, Ann. Soc. Ent. Belgique, LIV, p. 423 (♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 9. FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 211 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 13 (♂).

Type locality: NATAL (Haviland).

NATAL: Estcourt (R. C. Wroughton).

5. *Cerapachys peringueyi* EMERY, 1902, Rend. Accad. Sc. Bologna, N. S., VI, p. 24; 1911, 'Gen. Insect., Ponerinae,' p. 9, Pl. I, fig. 2. ARNOLD, 1915, Ann. South African Mus., XIV, p. 13 (♂), Pl. I, figs. 1, 1a.

Parasyscia peringueyi EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 360 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 17. EMERY, 1895, Zool. Jahrb. Abt. Syst., VIII, p. 757, fig. PP (♂, ♀).

Type locality: Cape of Good Hope (L. Péringuey).

TRANSVAAL.

5a. Var. *latiuscula* EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 9. ARNOLD, 1915, Ann. South African Mus., XIV, p. 14 (♂).

Parasyscia peringueyi var. *latiuscula* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 19 (♂).

Type locality: Makapan, TRANSVAAL (E. Simon).

6. *Cerapachys roberti* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 212 (♂).

Type locality: Estcourt, NATAL (R. C. Wroughton).

7. *Cerapachys wroughtoni* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 422 (♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 9 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 16 (♂).

Type locality: Mountains of NATAL (R. C. Wroughton).

7a. Var. *rhodesiana* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 212 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 16 (♂).

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

Phyracaces EMERY

Phyracaces EMERY, 1902, Rend. Accad. Sc. Bologna, N. S., VI, p. 23; 1911, 'Gen. Insect., Ponerinae,' p. 10.

Cerapachys (part) FOREL, in Grandidier, 1892, 'Hist. Phys. Nat. Madagascar,' XX, 2, p. 243.

Genotype: *Cerapachys mayri* Forel, 1892.

1. *Phyracaces braunsi* EMERY, 1902, Rend. Accad. Sc. Bologna, N. S., VI, p. 27 (♂); 1911, 'Gen. Insect., Ponerinae,' p. 11 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 17 (♂).

Type locality: Willowmore, CAPE PROVINCE (H. Brauns).

2. *Phyracaces cooperi* ARNOLD, 1915, Ann. South African Mus., XIV, p. 18 (♂).

Type locality: Umgeni, Durban, NATAL (C. B. Cooper).

2a. Subsp. **congolensis** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 401 (♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

3. **Phyracaces foreli** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 309 (♀); 1915, (part) Ann. Soc. Ent. France, LXXXIV, p. 245 (♀), fig. 2b (nec ♀, fig. 2a).

Type locality: Aburi, GOLD COAST (F. Silvestri).

4. **Phyracaces langi** WM. M. WHEELER. See p. 54 (♀, ♀).

Type locality: Lubila, BELGIAN CONGO (Lang and Chapin).

5. **Phyracaces nkomoensis** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 400 (♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

6. **Phyracaces santschii** WM. M. WHEELER. See p. 56 (♀).

Phyracaces foreli (part) SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 245 (♀), fig. 2a (nec ♀, fig. 2b).

Type locality: Samkita, FRENCH CONGO (F. Faure).

Lioponera MAYR

Lioponera MAYR, 1878, Verh. Zool. Bot. Ges. Wien, XXVIII, p. 666. EMERY, 1911, 'Gen. Insect., Ponerinæ,' p. 11.

Cerapachys subg. *Lioponera* FOREL, in Grandidier, 1892, 'Hist. Phys. Nat. Madagascar,' XX, 2, p. 244.

Genotype: *Lioponera longilarsus* Mayr, 1878.

The Ethiopian and North African species being described only from the male, it is very doubtful whether the genus really occurs in those regions.

1. **Lioponera decorsei** SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 150 (♂).

Type locality: Krebedje, southern Darbanda, FRENCH CONGO (Decorse).

2. **Lioponera nigra** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 46 (♂).

Type locality: Molo, Mau Escarpment, 2420 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

PONERINÆ Lepeletier

Cylindromyrmecini Emery

Simopone FOREL

Simopone FOREL, in Grandidier, 1891, 'Hist. Phys. Nat. Madagascar,' XX, 2, p. 139. EMERY, 1911, 'Gen. Insect., Ponerinæ,' p. 15.

Cerapachys subg. *Simopone* FOREL, in Grandidier, 1892, 'Hist. Phys. Nat. Madagascar,' XX, 2, p. 249.

Genotype: *Simopone grandidieri* Forel, 1891.

1. **Simopone conradti** EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 475 (♀); 1911, 'Gen. Insect., Ponerinæ,' p. 16 (♀), Pl. I, fig. 7.

Type locality: CAMEROON (Conradt).

2. **Simopone marleyi** ARNOLD, 1915, Ann. South African Mus., XIV, p. 20 (♀).

Type locality: Stella Bush, Durban, NATAL (Marley).

Amblyoponini Forel**Mystrium ROGER**

Mystrium ROGER, 1862, Berlin. Ent. Zeitschr., VI, p. 245. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 22.

Genotype: *Mystrium mysticum* ROGER, 1862.

1. **Mystrium silvestrii** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 310, fig. 1 (♂, ♀).

Type locality: Victoria, CAMEROON (F. Silvestri).

Xymmer (SANTSCHI)

Stigmatomma subg. *Xymmer* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 311.

Genotype: *Stigmatomma (Xymmer) muticum* Santschi, 1914.

1. **Xymmer muticum** (SANTSCHI).

Stigmatomma (Xymmer) muticum SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 311, fig. 2 (♂).

Type locality: Ibadan, NIGERIA (F. Silvestri).

Platythyreini Emery**Platythyrea ROGER**

Platythyrea ROGER, 1863, Berlin. Ent. Zeitschr., VII, p. 172. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 28.

Ponera (part) GERSTAECKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 262.

Pachycondyla (part) F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 108.

Genotype: *Pachycondyla punctata* F. Smith, 1858.

1. **Platythyrea arnoldi** FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 206 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 28 (♂).

Type locality: Shiloh, SOUTHERN RHODESIA (G. Arnold).

2. **Platythyrea conradti** EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 464 (♂, ♂); 1900, Bull. Soc. Ent. Italiana, XXXII, p. 107, fig. 2a (♂). FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 45 (♂); in Schulze, 1910, 'Forschungsreise in Südafrika,' IV, p. 3 (♂). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 129 (♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 29 (♂, ♂), Pl. II, figs. 1, 1b. FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 335 (♂, ♂); 1916, Rev. Suisse Zool., XXIV, p. 398 (♂, ♂). See p. 59.

Type locality: CAMEROON (Conradt).

WEST AFRICA: (Fülleborn). FRENCH CONGO: Gaboon (H. Petersen). FERNANDO PO: (Conradt). BELGIAN CONGO: St. Gabriel (Kohl); Risimu (Lang and Chapin).

3. **Platythyrea cooperi** ARNOLD, 1915, Ann. South African Mus., XIV, p. 29 (♂, ♂).

Type locality: Stella Bush, Durban, NATAL (C. B. Cooper).

4. **Platythyrea cribrinodis** (GERSTAECKER) ROGER, 1863, Berlin. Ent. Zeitschr., VII, p. 173; 1863, 'Verzeich. Formicid.,' p. 17. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 442. GERSTAECKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 355; 1873, in 'v. d. Decken's Reisen in Ost Afrika, Gliederthiere,' p. 346 (♂).

EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 270; 1892, *ibid.*, XXXII, p. 111 (♀). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 27. MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 200; 1895, Ann. Naturh. Hofmus. Wien, X, p. 125. EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 177; 1896, *ibid.*, XXXVII, p. 154 (♀, ♂); 1897, *ibid.*, XXXVIII, p. 597 (♀, ♂). H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 41. FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 76 (♀); 1907, Ann. Mus. Nat. Hungarici, V, p. 2 (♀). MAYR, 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. 1, p. 387; in Sjöstedt, 1907, 'Exped. Kili-mandjaro, Meru, etc.,' II, 8, p. 10 (♀, ♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 29 (♀). FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 361 (♀). EMERY, 1912, Annuar. Mus. Zool. Univ. Napoli, N. S., III, No. 26, p. 1 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 110 (♀); 1913, Deutsch. Ent. Zeitschr., Beih., p. 208 (♂). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 47 (♀, ♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 23 (♀, ♂), Pl. I, fig. 4.

Ponera cribrinodis GERSTAECKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 262 (♀); in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 504 (♀), Pl. XXXII, fig. 3. ROGER, 1860, Berlin. Ent. Zeitschr., IV, p. 295.

Type locality: MOZAMBIQUE (Peters).

RHODESIA: Bulawayo (G. Arnold); Lake Bangweolo (Elena d'Aosta). GERMAN EAST AFRICA: Lake Tanganyika (Reichardt); Tanga (H. Brauns); Mt. Lettema (Katona); Buiko (Zimmer); Ngare-na-Nyuki, Mt. Meru (Sjöstedt); Kihenga; Korogwe; Mbusini (Stuhlmann); Mt. Ugoni. BRITISH EAST AFRICA: Manda Island; Patta Island (Voeltzkow); Mombasa (v. d. Decken; C. Alluaud); Bura (H. Prell); Nairobi; Masai Plain; Tiwi; Lusinga Island, Victoria Nyanza; Fort Hall, Mt. Kenia, Meranga District (C. Alluaud); Maddo Wells (V. Bottego). SOMALI-LAND: Lugh; Matagoi to Lugh (V. Bottego); Mogadiscio; Obbia; Milmil (Bricchetti-Robecchi). ABYSSINIA: wells of Laffarugh to wells of Aberio, Ogaden (V. Bottego); Kaka, Schoa (Antinori); Salole; Lake Abaja; Bela; Daua (Ruspoli). UGANDA: Gondokoro (F. Werner).

4. Var. *brevidentata*, new name.

Platythyrea cribrinodis var. *punctata* ARNOLD, 1915, Ann. South African Mus., XIV, p. 24 (♀); *nec* *Platythyrea punctata* (F. Smith).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

4a. Var. *brevinodis* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 47 (♀).

Type locality: Samburu, Wa-Nyika District, BRITISH EAST AFRICA (C. Alluaud). BRITISH EAST AFRICA: Mwatate, Wa-Taita District; Kibwezi, Wa-Kamba District (C. Alluaud).

5. *Platythyrea crucheti* SANTSCHI, 1911, Rev. Zool. Afr., I, p. 205 (♀).

Type locality: CUCALA, BENGUELA (J. Cruchet).

6. *Platythyrea frontalis* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 466 (♀, ♀); 1911, 'Gen. Insect., Ponerinae,' p. 29 (♀, ♀).

Type locality: CAMEROON (Conradt).

7. *Platythyrea gracillima* WM. M. WHEELER. See p. 59 (♀).

Type locality: AVAKUBI, BELGIAN CONGO (Lang and Chapin).

8. *Platythyrea lamellosa* ROGER, 1863, Berlin. Ent. Zeitschr., VII, p. 173; 1863, 'Verzeich. Formicid.,' p. 17. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 442. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 27. MAYR, 1895, Ann. Naturh.

Hofmus. Wien, X, p. 125. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 29 (♀). FOREL, 1913, Rev. Zool. Afr., II, p. 307 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 25 (♀).

Ponera lamellosa ROGER, 1860, Berlin. Ent. Zeitschr., IV, p. 295 (♀).

Simaethiops var. *grisea* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 5 (♂).

Type locality: Cape of Good Hope.

CAPE PROVINCE: (G. Arnold). TRANSVAAL. ORANGE FREE STATE: (O. Schneider). BECHUANALAND: between Khakhea and Kang (L. Schultze).

8a. Subsp. *longinoda* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 75 (♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 3 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 29 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 26 (♀).

Type locality: Valdezia, TRANSVAAL (P. Berthoud).

BECHUANALAND: Kooa to Sekgoma (L. Schultze).

8a. Var. *rhodesiana* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 347 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 27 (♀, ♀, ♂), Pl. 1, fig. 3.

Platythyrea lamellosa var. *rhodesiana* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 110 (♀, ♂).

Type locality: SOUTHERN RHODESIA (G. Arnold).

BELGIAN CONGO: Kasongo (Pons).

8b. Subsp. *suturalis* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 64 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 29 (♀).

Type locality: BENGUELA (C. Wellman).

9. *Platythyrea mocquersyi* EMERY, 1900, Bull. Soc. Ent. Italiana, XXXI, p. 270 (♀, ♀). See p. 1007.

9. Var. *matopopensis* ARNOLD, 1915, Ann. South African Mus., XIV, p. 32 (♀).

Type locality: Matopopo Hills, SOUTHERN RHODESIA (G. Arnold).

10. *Platythyrea modesta* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 467 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 129 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 29 (♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 398 (♀). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp., 1910-11,' I, p. 371 (♀).

Type locality: CAMEROON.

CAMEROON: Mundame (Conradt). SPANISH GUINEA: Uelleburg (Tessmann). BELGIAN CONGO: Duma (Schubotz).

11. *Platythyrea occidentalis* ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 315 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 27. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 3 (♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 29 (♀, ♂).

Type locality: SIERRA LEONE (Mocquers).

FRENCH CONGO: Ogowe (Mocquers).

12. *Platythyrea schultzei* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 3 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 29 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 25 (♀).

Type locality: Salem, GERMAN SOUTHWEST AFRICA (L. Schultze).

SOUTHERN RHODESIA: Lonely Mine (Swale). GERMAN SOUTHWEST AFRICA: Windhoek (L. Schultze).

12. Var. *bequaerti* FOREL, 1913, Rev. Zool. Afr., II, p. 307 (♀, ♂).

Type locality: Nyangwe, BELGIAN CONGO (J. Bequaert).

BELGIAN CONGO: Sankisia (J. Bequaert).

13. *Platythyrea tenuis* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 467 (♀, ♀); 1911, 'Gen. Insect., Ponerinae,' p. 29 (♀, ♀).

Type locality: CAMEROON (Conradt).

14. *Platythyrea viehmeyeri* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 429 (♀).

Type locality: Khutu Steppe, GERMAN EAST AFRICA (K. Schwartze).

Proceratiini Emery

Sysphincta ROGER, *emend.*

Sysphincta ROGER, 1863, Berlin. Ent. Zeitschr., VII, p. 175.

Sysphincta MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidae, p. 12. EMERY, 1909, Deutsch. Ent. Zeitschr., p. 359; 1911, 'Gen. Insect., Ponerinae,' p. 50.

Proceratium (part) MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 437.

Ponera (part) ROGER, 1860, Berlin. Ent. Zeitschr., IV, p. 291.

Genotype: *Sysphincta micrommata* Roger, 1863.

1. *Sysphincta arnoldi* (FOREL) ARNOLD, 1915, Ann. South African Mus., XIV, p. 35 (♀).

Proceratium (*Sysphincta*) *arnoldi* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 210 (♀).

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

Discothyrea ROGER

Discothyrea ROGER, 1863, Berlin. Ent. Zeitschr., VII, p. 176. EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 51; 1911, 'Gen. Insect., Ponerinae,' p. 51.

Genotype: *Discothyrea testacea* Roger, 1863.

1. *Discothyrea hewitti* ARNOLD, 1916, Ann. South African Mus., XIV, p. 160 (♀).

Type locality: GRAHAMSTOWN, CAPE PROVINCE (Hewitt).

2. *Discothyrea oculata* EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 52 (♀, ♀, ♂). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 351 (♀, ♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 52 (♀, ♀, ♂). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 312 (♀).

Type locality: CAMEROON (L. Conradt).

FRENCH CONGO: Brazzaville (A. Weiss). FRENCH GUINEA: Mamou (F. Silvestri).

2. Var. *sculptor* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 302 (♀).

Type locality: FRENCH CONGO (A. Weiss).

3. *Discothyrea tragaordhi* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 3 (♀).

Type locality: Pietermaritzburg, NATAL (I. Trägårdh).

Probolomyrmex MAYR

Probolomyrmex MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 2. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 52.

Genotype: *Probolomyrmex filiformis* Mayr, 1901.

1. **Probolomyrmex filiformis** MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 3 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 52 (♀), Pl. II, figs. 10, 10b. SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 312 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 34 (♀), Pl. I, figs. 2, 2a.

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

FRENCH GUINEA: Kakoulima (F. Silvestri).

Escherichia FOREL

- Escherichia* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 245. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 53.

Genotype: *Escherichia brevisrostris* Forel, 1910.

1. **Escherichia brevisrostris** FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 246 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 53 (♀).

Type locality: Ghinda, ERITREA (K. Escherich).

Pseudosphincta ARNOLD

- Pseudosphincta* ARNOLD, 1916, Ann. South African Mus., XIV, p. 161.

Genotype: *Pseudosphincta poweri* Arnold, 1916.

1. **Pseudosphincta poweri** ARNOLD, 1916, Ann. South African Mus., XIV, p. 162, figs. 9, 9a (♀).

Type locality: Kimberley, CAPE PROVINCE (Power).

Ponerini FOREL

Centromyrmex MAYR

- Centromyrmex* MAYR, 1866, Verh. Zool. Bot. Ges. Wien, XVI, p. 894. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 57.

Spalacomyrmex EMERY, 1889, Ann. Mus. Civ. Genova, XXVII, p. 489.

Ponera (part) ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 5.

Genotype: *Centromyrmex bohemani* Mayr, 1866.

1. **Centromyrmex constanciae** ARNOLD, 1915, Ann. South African Mus., XIV, p. 38 (♀; ♀), Pl. II, figs. 14, 14a, 14b, 14c.

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

SOUTHERN RHODESIA: Bembesi (G. Arnold).

1. Var. **arnoldi** (SANTSCHI).

Centromyrmex arnoldi SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 229, fig. (♀, ♂).

- Centromyrmex constanciae* var. *arnoldi* SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 8.

Type locality: Amatongas Forest, PORTUGUESE EAST AFRICA (G. Arnold).

2. **Centromyrmex sellaris** MAYR, 1896, Ent. Tidskr., XVII, p. 230 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 58 (♀).

Type locality: CAMEROON (Sjöstedt).

2. Var. **longiventris** SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 229 (♀).

- Centromyrmex sellaris* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 313 (♀) (nec Mayr).

Type locality: Victoria, CAMEROON (F. Silvestri).

Streblognathus MAYR

Streblognathus MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 716. EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 468; 1911, 'Gen. Insect., Ponerinae,' p. 61.

Ponera (part) F. SMITH, ROGER.

Genotype: *Ponera aethiopica* F. Smith, 1858.

1. *Streblognathus aethiopicus* (F. SMITH) MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 716 (♀), Pl. XIX, figs. 10 and 11; 1863, *ibid.*, XIII, p. 454. ROGER, 1863, 'Verzeich. Formicid.,' p. 19. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 31. R. T. LEWIS, 1896, Journ. Quekett Micr. Club, (2) VI, p. 274, Pl. XI, figs. 1-8 (♀). EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 468 (♂); 1900, Bull. Soc. Ent. Italiana, XXXII, p. 105, fig. 1 (♂). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 51 (♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 3 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 61 (♀). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 3 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 41 (♀, ♂), Pl. I, fig. 5.

Ponera aethiopica F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 91 (♀), Pl. VI, figs. 19, 20. ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 11 (♀).

Type locality: South Africa.

BASUTOLAND. NATAL: VAN REENEN (I. Trägårdh); Durban (Wilms; v. Stuckrad); Richmond (J. R. Ward). CAPE PROVINCE: Willowmore (H. Brauns); Uitenhage. TRANSVAAL: (Heinemann).

Paltothyreus MAYR

Paltothyreus MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 735; 1866, *ibid.*, XVI, p. 894. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 62.

Formica (part) FABRICIUS. *Ponera* (part) GUÉRIN, F. SMITH, ROGER.

Pachycondyla (part) F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 105.

Genotype: *Formica tarsata* Fabricius, 1798.

1. *Paltothyreus tarsatus* (FABRICIUS) MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 736. ROGER, 1863, 'Verzeich. Formicid.,' p. 17. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 440; 1866, *ibid.*, XVI, p. 894 (♂), Pl. XX, fig. 6. FOREL, in Grandidier, 1891, 'Hist. Phys. Nat. Madagascar,' XX, 2, p. 136 (♂, ♀, ♂). EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 557 (♀, ♀, ♂); 1892, Ann. Mus. Civ. Genova, XXXII, p. 111 (♀). FOREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 352 (♀, ♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 32. MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 200. H. STADELMANN, 1893, Mitth. Deutsch. Schutzgeb., VI, p. 217. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 74. EMERY, 1894, Mem. Accad. Sc. Bologna, (5) IV, p. 112 (♀). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 126 (♀, ♀, ♂). EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 177. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 3 (♀, ♀). MAYR, 1896, Ent. Tidskr., XVII, p. 231. EMERY, 1896, Ann. Mus. Civ. Genova, XXXVII, p. 115 (♀, ♀, ♂); 1897, *ibid.*, XXXVIII, p. 597. FOREL, 1897, Abhandl. Senckenberg. Naturf. Ges., XXI, p. 188 (♀). H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 41. A. v. SCHULTHESS, 1899, Bull. Soc. Vaudoise Sc. Nat., (5) XXXV, p. 253 (♀). EMERY, 1899, Ann. Mus. Civ. Genova, XXXIX, p. 500 (♀, ♀, ♂); 1900, Bull. Soc. Ent. Italiana, XXXII, p. 112, fig. 5 (♂, ♀). SJÖSTEDT, 1904, 'I Västafrikas Urskogar,' p. 210. SCHÖNLAND, 1904, Trans. Ent. Soc. London, Proc., p.

xl. ZAVATTARI, 1907, Boll. Mus. Zool. Anat. Comp. Torino, XXII, No. 548, p. 2 (♀); 1907, *ibid.*, XXII, No. 550, p. 2 (♀). DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, pp. 354 and 357. MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.', II, 8, p. 8. FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika', II, p. 75 (♀); 1909, Ann. Soc. Ent. Belgique, LIII, p. 51 (♀, ♀) and p. 64 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 350 (♀, ♀). ZAVATTARI, 1909, 'Il Ruwenzori, Parte Scientif.', I, p. 213 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 129 (♀, ♀, ♂). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika', IV, p. 3 (♀); 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 361 (♀). STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08)', III, p. 377 (♀, ♀). EMERY, 1911, 'Gen. Insect., Ponerinae', p. 62 (♀), Pl. II, fig. 14; 1912, Annuar. Mus. Zool. Univ. Napoli, N. S., III, No. 26, p. 1 (♀). FOREL, 1913, Rev. Zool. Afr., II, p. 308 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 347 (♀, ♀, ♂); 1913, Rev. Suisse Zool., XXI, p. 666 (♀, ♂). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 313 (♀, ♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae', p. 47 (♀, ♀); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 4 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 44 (♀, ♀, ♂), Pl. I, figs. 6 and 10. STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. (1910-11)', I, p. 371 (♀). SANTSCHI, 1919, Rev. Zool. Afr., VII, pp. 79 and 81 (♀, ♀, ♂). CARPENTER, 1920, 'A Naturalist on Lake Victoria', pp. 50 and 281. See p. 61.

Formica tarsata FABRICIUS, 1798, 'Suppl. Ent. Syst.', p. 280 (♀); 1804, 'Syst. Piez.', p. 408 (♀).

Ponera tarsata LEPELETIER, 1836, 'Hist. Nat. Ins. Hym.', I, p. 194 (♀). ROGER, 1860, Berlin. Ent. Zeitschr., IV, p. 310; 1861, *ibid.*, V, p. 166.

Ponera gagates GUÉRHIN, 1845, 'Iconogr. Règne Anim.', VII, Insect., p. 423 (♀). FAIRMAIRE, in J. THOMSON, 1858, 'Archives Entomologiques', II, p. 263. F. SMITH, 1858, 'Cat. Hym. Brit. Mus.', VI, p. 90.

Ponera pestilentia F. SMITH, 1858, 'Cat. Hym. Brit. Mus.', VI, p. 92 (♀).

Ponera spiniventris F. SMITH, 1858, *ibid.*, p. 92 (♂).

Pachycondyla similima F. SMITH, 1858, *ibid.*, p. 105 (♀), Pl. VII, fig. 17. ROGER, 1863, 'Verzeich. Formicid.', p. 18.

Paltothyreus pestilentia A. GIRARD, in Capello and Ivens, 1882, 'From Benguela to the Territory of Yacca', II, p. 294. M. L. SYKES, 1900, Trans. Manchester Microsc. Soc., XX, (1899), p. 90. W. F. KIRBY, 1900, Proc. Zool. Soc. London, p. 47.

Paltothyreus pestilentia RADOSZKOWSKY, 1881, Journ. Sci. Ac. Lisboa, VIII, No. 31, p. 198. A. GIRARD, *ibid.*, p. 227.

Paltothyreus ? similima MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 358.

Type locality: Gorée Island, SENEGAMBIA.

SENEGAMBIA: Dakar (C. Alluaud); Thiès (F. Silvestri). CASAMANCE: (Claveau). PORTUGUESE GUINEA: Bissis Island (Lucas). FRENCH GUINEA: Los Islands (C. Alluaud), Konakry; Kakoulima; Kindia (F. Silvestri). SIERRA LEONE: Samlia Falls, River N'Gamie (Mocquerys). IVORY COAST: Dimbroko (Le Mout); Jacqueville; Grand Bassam (Lohier); Assinie (C. Alluaud). GOLD COAST: Akra. TOGO: Bismarckburg (Conradt; Büttner). CAMEROON: (Sjöstedt; F. Silvestri; Conradt); Mundame (Conradt); Bibundi (Tessmann). SPANISH GUINEA: Alen (Tessmann). FRENCH CONGO: Ogowe (Mocquerys); Fort Crampel (Le Mout); Brazzaville. BELGIAN CONGO: Kwesi to Kilo (Bayer); Kondué (Laja); Kiniati; Eala (R.

Mayné; Mayombe (Deleval); village of Denge, Niangara (Huttereau); Katanga (Lemaire); Ikelemba (Kinsberger); Boga; Beni; Avakubi; Ubangi District (Schubotz); Kindu (Grauer); Lake Leopold II (Leyder); Garamba, Yakuluku; Stanleyville; Medje; Risimu; Leopoldville; Bafwasende; Faradje; Gamangui; Bafwabaka; Niangara; Niapu; Ngayu; Akenge; Avakubi (Lang and Chapin). ANGOLA: (Welwitsch). BENGUELA: (C. Wellman). GERMAN SOUTHWEST AFRICA: (Lübbert); Okahandja (Peters); Omaheke (Trotha). RHODESIA: Luapula River (Elena d'Aosta); Bulawayo; Bembesi; Victoria Falls; Matoppos Hills (G. Arnold); Kazungula (Jallé). CAPE PROVINCE: Willowmore (H. Brauns). NATAL: Stamford Hill (I. Trägårdh); Port Natal (H. Brauns). ZULULAND: (Marley). PORTUGUESE EAST AFRICA: Delagoa (Liengme); Inhambane (Fornasini). GERMAN EAST AFRICA: Tanga; Rosako, Usaramo (Stuhlmann); Buiko; Monga (H. Prell); Amani (Zimmer); Mombo, Usambara (Sjöstedt); Lake Tanganyika (Reichardt). ZANZIBAR: (Voeltzkow). BRITISH EAST AFRICA: Patta Island (Voeltzkow); Shimoni; Mbuyuni, Pori; Kisumu, Victoria Nyanza; Nyangnori, western Nandi (Alluaud and Jeannel); Mombasa (H. Prell). UGANDA: Buzubizi (Alluaud and Jeannel); Butiti; Ibanda (Duke of Abruzzi). SOMALILAND: (C. Keller); Bulari (Peel); Merca (Taramasso); Mogadiscio; Obbia; Ellahelaj, Uorandi; Las Ej (Bricchetti-Robecchi); Hargeisa (V. Bottego). ABYSSINIA: Buditu to Dimé; Arussi Galla (V. Bottego); Webi (Bricchetti-Robecchi); Ogaden; Daua (Ruspoli).

Has been recorded from Madagascar, but undoubtedly through an error.

1. Var. **medianus** SANTSCHI, 1919, Rev. Zool. Afr., VII, pp. 80 and 82 (♀, ♂).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

CAMEROON: Molundu (Reichensperger). FRENCH CONGO: Ubangi (Augustin); Borda (Charleux). BENGUELA: Cucala (J. Cruchet).

12. Var. **subopacus** SANTSCHI, 1919, Rev. Zool. Afr., VII, pp. 80 and 82 (♀, ♀).

Type locality: Samkita, FRENCH CONGO (F. Faure).

13. Subsp. **delagoensis** (EMERY) SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 48 (♀, ♀).

Pallothyreus tarsatus var. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 126.

Pallothyreus tarsatus var. *delagoensis* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 468 (♀, ♀); 1911, 'Gen. Insect., Ponerinae,' p. 62 (♀, ♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 45 (♀). SANTSCHI, 1919, Rev. Zool. Afr., VII, pp. 80 and 82 (♀).

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA.

RHODESIA: (G. Arnold). GERMAN EAST AFRICA: Mt. Kilimanjaro, River Himo (C. Alluaud); Pangani (Reichensperger). BRITISH EAST AFRICA: Mt. Kenia, Meranga, Fort Hall; Bura, Wa-Taita; Kibwezi, Wa-Kamba; Mombasa; Maji-yachumvi, Wa-Nyika; Samburu, Wa-Nyika (Alluaud and Jeannel).

14. Var. **robustus** (SANTSCHI).

Pallothyreus tarsatus var. *robusta* SANTSCHI, 1919, Rev. Zool. Afr., VII, pp. 81 and 82 (♀, ♀).

Type locality: Buarsangueli, SOMALILAND (Revoil).

15. Var. **striatidens** (SANTSCHI).

Pallothyreus tarsatus var. *striatidens* SANTSCHI, 1919, Rev. Zool. Afr., VII, pp. 81 and 82 (♀, ♀).

Type locality: Kibwezi, BRITISH EAST AFRICA (C. Alluaud).

Glyphopone FOREL

Glyphopone FOREL, 1913, Rev. Zool. Afr., II, p. 308.

Genotype: *Glyphopone bequaerti* FOREL, 1913.

1. ***Glyphopone bequaerti*** FOREL, 1913, Rev. Zool. Afr., II, p. 308, fig. 1 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 347 (♀). BEQUAERT, 1913, Rev. Zool. Afr., II, p. 422.

Type locality: Kibombo, BELGIAN CONGO (J. Bequaert).

BELGIAN CONGO: Kasongo (Pons).

Leptopone (ARNOLD)

Glyphopone subg. *Leptopone* ARNOLD, 1916, Ann. South African Mus., XIV, p. 163.

Genotype: *Glyphopone (Leptopone) rufigaster* ARNOLD, 1916.

1. ***Leptopone rufigaster*** (ARNOLD).

Glyphopone (Leptopone) rufigaster ARNOLD, 1916, Ann. South African Mus., XIV, p. 163, figs. 10, 10a (♀).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

Megaponera MAYR

Megaponera MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 734. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 67.

Megaloponera ROGER, 1863, 'Verzeich. Formicid.,' p. 17.

Ponera (part) GUÉRIN, GERSTAECKER, ROGER. *Formica* (part) FABRICIUS, LATREILLE.

Genotype: *Formica fœtens* Fabricius, 1793.

1. ***Megaponera fœtens*** (FABRICIUS) MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 735 (♀); 1863, *ibid.*, XIII, p. 428. MAGRETTI, 1884, Ann. Mus. Civ. Genova, XXI, p. 538 (♀); 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 244 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 30. MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 200; 1895, Ann. Naturh. Hofmus. Wien, X, p. 125. EMERY, 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 597 (♀, ♂). H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 41. GESTRO, 1901, Ann. Mus. Civ. Genova, XL, p. 729. MAYR, 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. 1, p. 387. STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 130 (♀); 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exped. (1907-08),' III, p. 377 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 68 (♀); 1912, Annuar. Mus. Zool. Univ. Napoli, N. S., III, No. 26, p. 1 (♀). ARNOLD, 1913, Proc. Rhodesia Sc. Assoc., XII, p. 12; 1914, *ibid.*, XIII, p. 26. H. BRAUNS, 1914, *ibid.*, XIII, pp. 33, 35, 39, and 42. SANTSCI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 48 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 313 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 46, fig. 6 (♀, ♀, ♂), Pl. I, figs. 7, 8, and 8a. POULTON, 1915, Trans. Ent. Soc. London, pp. v-vii and lvi-lxi. BICKHARDT, 1916, Entom. Blätter, XII, p. 2. STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 371 (♀, ♂). CARPENTER, 1918, Trans. Ent. Soc. London, (1917), Proc., p. lx; 1920, 'A Naturalist on Lake Victoria,' p. 282. See p. 64.

Formica fœtens FABRICIUS, 1793, 'Ent. Syst.,' II, p. 354 (♀); 1804, 'Syst. Piez.,' p. 401 (♀). F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 32.

Formica analis LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 282 (♀).

Ponera abyssinica GUÉRIN, in Lefebvre, 1848, 'Voy. Abyssinie,' IV, Zool., part 6, p. 352 (♀), Pl. VII, fig. 6. F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 90. RADOSZKOWSKY, 1876, Hor. Soc. Ent. Rossice, XII, p. 140. CANNAVIELLO, 1900, Bull. Soc. Ent. Italiana, XXXII, p. 297.

Ponera laviuscula GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 262 (♀); in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 505 (♀), Pl. XXXII, fig. 4.

Ponera fatens GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 262; 1859, Arch. f. Naturg., XXV, 2, p. 431. ROGER, 1860, Berlin. Ent. Zeitschr., IV, p. 310 (♀). GERSTÄCKER, 1862, in Peters, 'Reise n. Mossambique, Zool.,' V, p. 504 (♀).

Megaloponera fatens ROGER, 1863, 'Verzeich. Formicid.,' p. 17. EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 368. RADOSZKOWSKY, 1881, Journ. Sci. Ac. Lisboa, VIII, No. 31, p. 197. FOREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 352 (♀); 1894, ibid., IX, p. 75. EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 177; 1899, ibid., XXXIX, p. 500 (♂). FOREL, 1904, Ann. Mus. Zool. Ac. Sc. St. Pétersbourg, VIII, (1903), p. 369 (♀, ♂); 1907, Ann. Mus. Nat. Hungarici, V, p. 2 (♀). MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 9 (♀, ♂). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 64 (♀); 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 244 (♀). PRELL, 1911, Zool. Anzeiger, XXXVIII, p. 244, fig. 1. FOREL, 1911, Rev. Zool. Afr., I, p. 274 (♀); 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 361 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, pp. 108 (♀) and 347 (♀, ♂); 1913, Rev. Zool. Afr., II, p. 308 (♀); 1913, Deutsch. Ent. Zeitschr., Beih., p. 206. J. BEQUAERT, 1913, Rev. Zool. Afr., II, p. 422 (♀).

Ponera crassicornis GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 262 (♀); 1862, in Peters, 'Reise n. Mossambique, Zool.,' V, p. 506 (♀), Pl. XXXII, fig. 5. ROGER, 1860, Berlin. Ent. Zeitschr., IV, p. 309; 1863, 'Verzeich. Formicid.,' p. 23. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 448.

Megaponera crassicornis MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 200. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 30. EMERY, 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 597 (♀). H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 41.

Megaloponera crassicornis EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 368 (♀). FOREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 352 (♀); 1907, Ann. Mus. Nat. Hungarici, V, p. 2 (♀).

Megaponera dohrni EMERY, 1902, Rend. Accad. Sc. Bologna, N. S., VI, p. 30 (♀).

Type locality: Guinea (Isert).

SENEGAMBIA: Thiès (F. Silvestri). SIERRA LEONE: (F. Silvestri). GOLD COAST: Akropong (Imhoff). TOGO: (H. Brauns); Bismarekburg (Conradt). SOUTHERN NIGERIA: Agege; Ibadan (Farquharson). CAMEROON: Metit (G. Schwab). FRENCH CONGO: Fort de Pessel; Fort Crampel (Schubotz). BELGIAN CONGO: Lukula (Daniel); Kondué (Lujia); Kambove (S. A. Neave); Malela; Sankisia (J. Bequaert); Kwesi to Kilo (Bayer); Kindu (Burgeon); Kwidjwi Island, Lake Kivu; Libenge (Schubotz); Zambi; Niangara; Rungu; Avakubi; Faradje; Panga to Banjala; Boyulu; Niapu; Garamba; Akenge; Gamangui (Lang and Chapin). ANGOLA: (Welwitsch). BENGUELA: (C. Wellman). RHODESIA: Mumbwa (Dollman); Leangwa Valley (S. A. Neave); Bulawayo (G. Arnold); Luapula River (Elena d'Aosta). PORTUGUESE EAST AFRICA: Tete; Mozambique (Peters). GERMAN EAST AFRICA: Mt. Kilimanjaro (Bornemisza); Moschi (Katona); Njussi

(H. Prell); Amani; near Buiko (Zimmer); Mbusini; Bondei (Stuhlmann); Kibonoto, Mt. Kilimanjaro (Sjöstedt); Itigi; Kyaka Fort on the Kagera River (G. D. H. Carpenter); Usambara. ZANZIBAR. BRITISH EAST AFRICA: Pori of Serengeti between Bura and Taveta; Fort Hall; Mbuyuni (Alluaud and Jeannel). UGANDA: Unyoro Province, Lake Albert (C. Alluaud); Gondokoro (F. Werner). ABYSSINIA: Arussi Galla, Ganale Gudda; Saneur to Amarr Burgi; Dimé to Bass Narok (V. Bottego); Sciotel (Beccari); Daua; Ogaden; Degabolla; Lake Abaja; Brava (Ruspoli); Gogfale; Harar (Kachovskij); southern Abyssinia (Ilg); Adua; Makalle. SOMALILAND: (C. Keller); Lower Ganana (V. Bottego). ERITREA: Keren (Beccari; Magretti); Bogos (Antinori); Ghinda (K. Escherich). ANGLO-EGYPTIAN SUDAN: Sennar.

1. Var. **rapax** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 49 (♀).

Type locality: New Moschi, Mt. Kilimanjaro, 800 m., GERMAN EAST AFRICA (Alluaud and Jeannel).

GERMAN EAST AFRICA: Moschi and Kilema, 1400-1500 m. (Alluaud and Jeannel).

Ophthalmopone FOREL

Ophthalmopone FOREL, 1890, Ann. Soc. Ent. Belgique, XXXIV, C. R., p. cxi. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 69.

Pachycondyla (part) EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 360.

Genotype: *Ophthalmopone berthoudi* Forel, 1890.

1. **Ophthalmopone berthoudi** FOREL, 1890, Ann. Soc. Ent. Belgique, XXXIV, C. R., p. cxii (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 31. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 76 (♀, ♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 69 (♀, ♂). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 108 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 50 (♀, ♂), Pl. III, figs. 11, 13.

Type locality: Valdezia, TRANSVAAL (P. Berthoud).

SOUTHERN RHODESIA: Bulawayo (G. Arnold). PORTUGUESE EAST AFRICA: Delagoa (Liengme).

2. **Ophthalmopone depilis** EMERY, 1902, Rend. Accad. Sc. Bologna, N. S., VI, p. 29 (♀); 1911, 'Gen. Insect., Ponerinae,' p. 69 (♀).

Type locality: SAN THOMÉ (Mocquereys).

3. **Ophthalmopone hottentota** EMERY, 1902, Rend. Accad. Sc. Bologna, N. S., VI, p. 30; 1911, 'Gen. Insect., Ponerinae,' p. 70 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 52 (♀, ♂), Pl. II, fig. 12.

Pachycondyla hottentota EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 360 (♀); 1890, Ann. Soc. Ent. France, (6) X, p. 72 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 34.

Ophthalmopone lanceolata MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 125 (♀).

Type locality: Cape of Good Hope (L. Péringuey).

SOUTHERN RHODESIA: Sebakwe (G. Arnold). CAPE PROVINCE: generally (G. Arnold); Willowmore (H. Brauns).

4. **Ophthalmopone ilgi** FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 76 (♀). EMERY, 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 597 (♀); 1911, 'Gen. Insect., Ponerinae,' p. 70 (♀).

Type locality: SOUTHERN ABYSSINIA (Ilg).

SOMALILAND: (Ruspoli).

5. *Ophthalmopone mocquersyi* EMERY, 1902, Rend. Accad. Sc. Bologna, N. S., VI, p. 29 (♀). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 422 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 70 (♀).

Type locality: SAN THOMÉ (Mocquers).

ANGOLA: Mossamedes.

Bothroponera MAYR

Bothroponera MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 717.

Pachycondyla subg. *Bothroponera* EMERY, 1911, 'Gen. Insect., Ponerinae,' p.

75. FOREL, ARNOLD, SANTSCHI.

Pachycondyla (part) F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 105.

Ponera (part) JERDON, F. SMITH, ROGER.

Genotype: *Bothroponera pumicosa* Roger, 1860.

1. *Bothroponera cariosa* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 20 (♀).

Pachycondyla (*Bothroponera*) *cariosa* EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 45; 1911, 'Gen. Insect., Ponerinae,' p. 76 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 59 (♀).

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA.

CAPE PROVINCE. TRANSVAAL.

2. *Bothroponera cavernosa* (ROGER) MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 717; 1863, *ibid.*, XIII, p. 397. ROGER, 1863, 'Verzeich. Formicid.,' p. 16. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 36.

Ponera cavernosa ROGER, 1860, Berlin. Ent. Zeitschr., IV, p. 288 (♀).

Pachycondyla (*Bothroponera*) *cavernosa* EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 45; 1911, 'Gen. Insect., Ponerinae,' p. 76. ARNOLD, 1915, Ann. South African Mus., XIV, p. 60 (♀).

Type locality: Caffraria.

CAPE PROVINCE.

3. *Bothroponera crassa* EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 111 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 36. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 75. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 2 (♀).

Ponera crassa EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 366, fig. (♀).

Ponera (*Bothroponera*) *crassa* EMERY, 1895, *ibid.*, XXXV, p. 177.

Pachycondyla (*Bothroponera*) *crassa* EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 46. FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 244 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 77 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 347 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 50 (♀).

Type locality: Sciotel, ABYSSINIA (Beccari).

FRENCH CONGO: Ogowe (Mocquers). BELGIAN CONGO: Mayombe (de Briey). BRITISH EAST AFRICA: Nairobi (C. Alluaud). ERITREA: Nefasit (Ilg). ABYSSINIA: (Ilg); Arussi Galla, Ganale Gudda; Wells of Laffarugh to Wells of Aberio; Ogaden (V. Bottego); Webi (Bricchetti-Robecchi). SOMALILAND: Obbia (Bricchetti-Robecchi); Erdal.

3. *Var. ilgi* (FOREL).

Pachycondyla (*Bothroponera*) *crassa* var. *ilgi* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 244 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 77 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 50 (♀).

Type locality: Schoa, ABYSSINIA (Ilg).

BRITISH EAST AFRICA: Kenia Region between the Amboni and Naremurur Rivers, 1800-2000 m. (Alluaud and Jeannel).

4. *Bothroponera cribrata* (SANTSCHI).

Pachycondyla (*Bothroponera*) *cribrata* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 349 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 76 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

5. *Bothroponera fugax* (FOREL).

Pachycondyla (*Bothroponera*) *fugax* FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 7 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 76 (♀).

Type locality: Arusha-chini, Lake Djiye, GERMAN EAST AFRICA (Katona).

6. *Bothroponera granosa* (ROGER) MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 717; 1863, *ibid.*, XIII, p. 397. ROGER, 1863, 'Verzeich. Formicid.,' p. 16. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 36.

Ponera granosa ROGER, 1860, Berlin. Ent. Zeitschr., IV, p. 290 (♀).

Pachycondyla (*Bothroponera*) *granosa* EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 45; 1911, 'Gen. Insect., Ponerinae,' p. 76 (♀). FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 214 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 61 (♀), Pl. II, fig. 16.

Type locality: Cape of Good Hope (Drège).

CAPE PROVINCE: George, Knysna (H. Brauns). NATAL: SOUTHERN RHODESIA: Victoria Falls (L. Péringuey).

7. *Bothroponera krügeri* (FOREL).

Pachycondyla (*Bothroponera*) *krügeri* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 28 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 56 (♀).

Pachycondyla (*Bothroponera*) *kruegeri* EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 77 (♀). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 4 (♀). TRÄGÅRDH, *ibid.*, p. 43.

Type locality: Valdezia, TRANSVAAL (P. Berthoud).

ZULULAND: Umfolosi (I. Trägårdh). TRANSVAAL: (G. Arnold).

7. *Var. rhodesiana* (FOREL).

Pachycondyla (*Bothroponera*) *krügeri* var. *rhodesiana* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 109 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 57 (♀, ♂).

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

7a. Subsp. *asina* (SANTSCHI).

Pachycondyla (*Bothroponera*) *asina* SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 153 (♀).

Pachycondyla (*Bothroponera*) *kruegeri* subsp. *asina* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 4 (♀).

Type locality: BRITISH EAST AFRICA (Demarchi).

ZULULAND: Umfolosi (I. Trägårdh).

8. *Bothroponera laevissima* (ARNOLD).

Pachycondyla (Bothroponera) levissima ARNOLD, 1915, Ann. South African Mus., XIV, p. 58 (♀), Pl. II, figs. 15, 15a.

Type locality: Saldanha Bay, CAPE PROVINCE (G. Arnold).

9. ***Bothroponera nasica* (SANTSCHI).**

Pachycondyla (Bothroponera) nasica SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 6 (♀).

Type locality: Samkita, FRENCH CONGO (F. Faure).

10. ***Bothroponera pachyderma* (EMERY).** See p. 73.

Pachycondyla (Bothroponera) pachyderma EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 49 (♀, ♀). MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 9 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 76 (♀, ♀).

Type locality: CAMEROON (Conradt).

GERMAN EAST AFRICA: Kibonoto, Mt. Kilimanjaro, 1000-2000 m.; Natron Lakes (Sjöstedt). BELGIAN CONGO: Manamaba; Bafwasende; Medje; Ngayu; Niapu; Niangara; Akengé (Lang and Chapin).

10. **Var. *attenata* (SANTSCHI).**

Pachycondyla (Bothroponera) pachyderma var. *attenata* SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 7 (♀).

Type locality: N'gazi, BELGIAN CONGO (Elskens).

10. **Var. *funerea* Wm. M. WHEELER.** See p. 73 (♀).

Type locality: Medje, BELGIAN CONGO (Lang and Chapin).

10. **Var. *postsquamosa* (SANTSCHI).**

Pachycondyla (Bothroponera) pachyderma var. *postsquamosa* SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 7 (♀).

Type locality: Lobay, FRENCH CONGO (Riggenbach).

11. ***Bothroponera picardi* (FOREL).**

Pachycondyla (Bothroponera) picardi FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 304 (♀); in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 559 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 77 (♀):

Type locality: between the Cubango and Cuito Rivers, MOSSAMEDES (Baum and Van der Kellen).

12. ***Bothroponera pumicosa* (ROGER) MAYR,** 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 717; 1863, *ibid.*, XIII, p. 397. ROGER, 1863, 'Verzeich. Formicid.,' p. 16. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 36.

Poner a pumicosa ROGER, 1860, Berlin. Ent. Zeitschr., IV, p. 290 (♀).

Pachycondyla pumicosa FOREL, 1901, Rev. Suisse Zool., IX, p. 344.

Pachycondyla (Bothroponera) pumicosa EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 45. STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 130 (♀). FOREL, in Schultze, 1910, 'Forschungsreise in Sudafrica,' IV, p. 4 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 76 (♀). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 4 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 62 (♀, ♂).

Type locality: Caffraria.

CAMEROON: Mundame (Conradt). CAPE PROVINCE: (H. Brauns; Drège); Cape Town (Wilms). NATAL: (Haviland); Van Reenen (I. Trägårdh).

12. **Var. *berthoudi* (FOREL).**

Pachycondyla berthoudi FOREL, 1901, Rev. Suisse Zool., IX, p. 344 (♀).

Pachycondyla (Bothroponera) berthoudi EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 76 (♀). FOREL, 1913, Rev. Zool. Afr., II, p. 306 (♀, ♂).

Pachycondyla (Bothroponera) pumicosa subsp. *berthoudi* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 109 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 63 (♂, ♂).

Type locality: Valdezia, TRANSVAAL (P. Berthoud).

CAPE PROVINCE: Willowmore (H. Brauns).

12. Var. *sculpturata* (SANTSCHI).

Pachycondyla (Bothroponera) sculpturata SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 151 (♀).

Type locality: Zambésie, RHODESIA (Demarchi).

MOZAMBIQUE: (G. Vasse).

13. *Bothroponera sanguinea* (SANTSCHI).

Pachycondyla (Bothroponera) sanguinea SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 7 (♀).

Type locality: CONGO (obtained from Le Mout).

14. *Bothroponera silvestrii* (SANTSCHI).

Pachycondyla (Bothroponera) silvestrii SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 313 (♀).

Type locality: Aburi, GOLD COAST (F. Silvestri).

15. *Bothroponera sjöstedti* (MAYR). See p. 74.

Ponera sjöstedti MAYR, 1896, Ent. Tidskr., XVII, p. 231 (♀, ♀, ♂).

Pachycondyla (Bothroponera) sjöstedti EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 45; 1911, 'Gen. Insect., Ponerinae,' p. 77 (♀, ♀, ♂).

Pachycondyla (Bothroponera) sjöstedti SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 315, fig. 3 (♀).

Type locality: CAMEROON (Sjöstedt).

CAMEROON: Victoria (F. Silvestri); Mbalmajo to Ekeneli (G. Schwab). BELGIAN CONGO: Malela (J. Bequaert).

16. *Bothroponera soror* (EMERY). See p. 74.

Ponera (Bothroponera) soror EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 472 (♀, ♀).

Pachycondyla (Bothroponera) soror EMERY, 1901, *ibid.*, XLV, p. 46. SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 350 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 129 (♀, ♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 78 (♀, ♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 110 (♀); 1913, Rev. Zool. Afr., II, p. 306 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 59 (♀).

Pachycondyla soror BEQUAERT, 1913, Rev. Zool. Afr., II, p. 421.

Type locality: CAMEROON (Conradt).

TOGO: Bismarckburg (Conradt). CAMEROON: Metit (G. Schwab); Mundame (Conradt); Yaunde (Zenker). FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: Elisabethville; Sankisia (J. Bequaert); Akenge; Medje; Ngayu; Niangara; Avakubi; Niapu (Lang and Chapin). RHODESIA: Springvale; Bulawayo (G. Arnold).

16. Var. *ancilla* (EMERY). See p. 74.

Ponera (Bothroponera) soror var. *ancilla* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 472 (♀).

Pachycondyla (Bothroponera) soror var. *ancilla* EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 46; 1911, 'Gen. Insect., Ponerinae,' p. 78 (♀).

Type locality: River Kuilu, FRENCH CONGO.

BELGIAN CONGO: Isangi (Lang and Chapin).

16a. Subsp. *suturalis* (FOREL).

Pachycondyla (*Bothroponera*) *soror* subsp. *suturalis* FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 133 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 78 (♀).

Type locality: KARSSA, ABYSSINIA (de Rothschild).

17. *Bothroponera strigulosa* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 19 (♀).

Pachycondyla (*Bothroponera*) *strigulosa* EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 45; 1911, 'Gen. Insect., Ponerinae,' p. 76 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 61 (♀).

Type locality: Kimberley, CAPE PROVINCE (E. Simon).

18. *Bothroponera talpa* ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 316 (♂); DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 37. See p. 73.

Ponera (*Bothroponera*) *talpa* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 473 (♀).

Pachycondyla (*Bothroponera*) *talpa* EMERY, 1901, *ibid.*, XLV, p. 45. SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 314 (♀).

Type locality: SIERRA LEONE (Mocquers).

FRENCH GUINEA: Kakoulima (F. Silvestri). GOLD COAST: Aburi (F. Silvestri). CAMEROON. BELGIAN CONGO: Niapu; Niangara; Avakubi; Medje (Lang and Chapin).

18a. Subsp. *variolata* (SANTSCHI).

Pachycondyla (*Bothroponera*) *variolata* SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 151 (♀).

Pachycondyla (*Bothroponera*) *talpa* subsp. *variolata* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 314.

Type locality: FRENCH CONGO (Zimmermann).

Phrynoponera WM. M. WHEELER

Phrynoponera WM. M. WHEELER, 1920, Psyche, XXVII, p. 53.

Genotype: *Bothroponera gabonensis* ERN. ANDRÉ, 1892.

1. *Phrynoponera armata* (SANTSCHI).

Pachycondyla (*Bothroponera*) *armata* SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 82 (♀).

Type locality: Kitempuka, BELGIAN CONGO (Gérard).

2. *Phrynoponera bequaerti* WM. M. WHEELER. See p. 79 (♀).

Type locality: Ngayu, BELGIAN CONGO (Lang and Chapin).

3. *Phrynoponera gabonensis* (ERN. ANDRÉ). See p. 76.

Bothroponera gabonensis ERN. ANDRÉ, 1892, Rev. d'Ent. Caen, XI, p. 50 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 36.

Pachycondyla (*Bothroponera*) *gabonensis* EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 45. STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 130, fig. 1 (♀, ♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 76 (♀).

Type locality: Gaboon, FRENCH CONGO.

BELGIAN CONGO: Kimpoko (Büttner); Bafwasende; Medje; Akenge (Lang and Chapin).

31. Var. **esta** Wm. M. WHEELER. See p. 77 (♀, ♀).
Type locality: Medje, BELGIAN CONGO (Lang and Chapin).
BELGIAN CONGO: Gamangui; Ngayu (Lang and Chapin).
32. Var. **fecunda** Wm. M. WHEELER. See p. 78 (♀, ♀).
Type locality: Akenge, BELGIAN CONGO (Lang and Chapin).
BELGIAN CONGO: Medje; Ngayu; Avakubi (Lang and Chapin).
33. Var. **robustior** (SANTSCHI).
Pachycondyla (*Bothroponera*) *gabonensis* var. *robustior* SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 82 (♀).
Type locality: Banalia, BELGIAN CONGO (J. Bequaert).
34. Var. **striatidens** (SANTSCHI). See p. 78.
Pachycondyla (*Bothroponera*) *gabonensis* var. *striatidens* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 315, fig. 4 (♀).
Type locality: Victoria, CAMEROON (F. Silvestri).
BELGIAN CONGO: Medje; Akenge; Ngayu (Lang and Chapin).
35. Var. **umbrosa** Wm. M. WHEELER. See p. 78 (♀).
Type locality: Medje, BELGIAN CONGO (Lang and Chapin).
4. **Phrynoponera heterodus** Wm. M. WHEELER. See p. 78 (♀).
Type locality: Stanleyville, BELGIAN CONGO (Lang and Chapin).
5. **Phrynoponera sveni** (FOREL). See p. 79.
Pachycondyla (*Bothroponera*) *sveni* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 398 (♀).
Type locality: BELGIAN CONGO (Kohl).
BELGIAN CONGO: Medje (Lang and Chapin).

Ectomomyrmex MAYR

- Ectomomyrmex* MAYR, 1867, Tijdschr. v. Ent., X, p. 83.
Pachycondyla subg. *Ectomomyrmex* EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 78.
Pachycondyla (part) F. SMITH. *Bothroponera* (part) EMERY, FOREL. *Ponera* (part) ERN. ANDRÉ, FOREL.
Genotype: *Ectomomyrmex javanus* Mayr, 1867.
1. **Ectomomyrmex brunoi** (FOREL).
Pachycondyla (*Ectomomyrmex*) *brunoi* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 205 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 53 (♀, ♂).
Type locality: Bulawayo, RHODESIA (G. Arnold).

Euponera FOREL

- Ponera* subg. *Euponera* FOREL, in Grandidier, 1891, 'Hist. Phys. Nat. Madagascar,' XX, 2, p. 126 (*sensu stricto*).
Euponera EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 43 (*sensu latiore*).
FOREL, 1901, *ibid.*, XLV, p. 141. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 79.
Ponera (part) F. SMITH, ROGER, MAYR, etc. *Formica* (part) FABRICIUS.
Genotype: *Euponera sikora* Forel, 1891.

Subgenus 1. Mesoponera EMERY

- Euponera* subg. *Mesoponera* EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 43; 1911, 'Gen. Insect., Ponerinae,' p. 80.

Mesoponera BINGHAM, 1903, 'Fauna Brit. India, Hym.,' II, p. 99.

Ponera subg. *Xiphopelta* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 108 (misspelled *Hiphopelta* on the same page).

Euponera subg. *Xiphopelta* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 206.

Pachycondyla (part) MAYR, 1866, Verh. Zool. Bot. Ges. Wien, XVI, p. 890.

Ponera (part) F. SMITH, ROGER, MAYR, etc.

Subgenotype: *Ponera cafraria* F. Smith, 1858.

1. ***Euponera (Mesoponera) cafraria*** (F. SMITH) EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 46; 1911, 'Gen. Insect., Ponerinae,' p. 81 (♀, ♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 108 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 65 (♀, ♀, ♂), Pl. II, figs. 17, 17a and Pl. III, fig. 22.

Ponera cafraria F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 91 (♀). ROGER, 1860, Berlin. Ent. Zeitschr., IV, p. 285; 1863, 'Verzeich. Formicid.,' p. 16. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 447; 1886, *ibid.*, XXXVI, p. 358. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 38. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 124 (♀, ♀). EMERY, 1899, Mem. Accad. Sc. Bologna, (5) VIII, p. 3 (larva), Pl. I, fig. 2.

Ponera guineensis ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 318 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 39. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 20.

Mesoponera cafraria DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, pp. 332 and 342 (♀).

Euponera (Brachyponera) cafraria SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 350.

Euponera (Mesoponera) cafra SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 315 (♀).

Type locality: Port Natal, NATAL.

FRENCH GUINEA: Kakoulima (F. Silvestri). SIERRA LEONE: (Mocquerys). CAMEROON: (H. Brauns). FRENCH CONGO: Brazzaville (A. Weiss). RHODESIA: Springvale (G. Arnold). BECHUANALAND: Serui (Dixey and Longstaff). TRANSVAAL: Pretoria (E. Simon). NATAL: (C. B. Cooper and Marley); Colenso (Dixey and Longstaff). CAPE PROVINCE: Willowmore (H. Brauns).

2. ***Euponera (Mesoponera) elisæ*** (FOREL). See p. 1008.

2a. Var. ***redbankensis*** (FOREL).

Ponera (Xiphopelta) arnoldi var. *redbankensis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 109 (♀).

Euponera (Mesoponera) elisæ var. *redbankensis* ARNOLD, 1915, Ann. South African Mus., XIV, p. 71 (♀).

Type locality: Redbank, RHODESIA (G. Arnold).

2a. Subsp. ***divaricata*** EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 6, fig. 3 (♀).

Type locality: Keren, ERITREA (F. Silvestri).

ERITREA: Ghinda; Nefasit (F. Silvestri).

2b. Subsp. ***rotundata*** (EMERY) FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 206 (♀).

Ponera elisæ var. *rotundata* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 20 (♀).

Euponera (Mesoponera) elisæ var. *rotundata* EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 46; 1911, 'Gen. Insect., Ponerinae,' p. 81 (♀).

Euponera (*Mesoponera*) *eliza* var. *rotundata* ARNOLD, 1915, Ann. South African Mus., XIV, p. 70 (♀, ♀).

Ponera (*Xiphopelta*) *arnoldi* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 108 (♀).

Type locality: Bloemfontein, ORANGE FREE STATE (E. Simon).

RHODESIA: Bulawayo (G. Arnold).

3. *Euponera* (*Mesoponera*) *escherichi* (FOREL) EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 3 (♀).

Pachycondyla (*Bothropsonera*) *escherichi* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 244 (♀).

Euponera (*Mesoponera*?) *escherichi*, EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 82.

Type locality: Nefasit, ERITREA (K. Escherich; F. Silvestri).

4. *Euponera* (*Mesoponera*) *fossigera* MAYR, ARNOLD, 1915, Ann. South African Mus., XIV, p. 71 (♀), Pl. II, fig. 20.

Euponera fossigera MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 3 (♀).

Euponera (*Mesoponera*?) *fossigera* EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 82.

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

5. *Euponera* (*Mesoponera*) *haviglandi* (FOREL).

Megaloponera (*Hagensia*) *haviglandi* FOREL, 1901, Rev. Suisse Zool., IX, p. 333 (♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 3 (♀).

Megaloponera (*Hagensia*) *haviglandi* EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 69 (♀).

Euponera (*Mesoponera*) *sulcigera* MAYR, 1904, Verh. Zool. Bot. Ges. Wien, LIV, p. 593 (♀).

Euponera haviglandi ARNOLD, 1915, Ann. South African Mus., XIV, p. 67 (♀, ♂).

Type locality: NATAL (Haviland, R. C. Wroughton).

NATAL: Durban (Wilms). CAPE PROVINCE: Knysna (H. Brauns)¹; Grahams-town (Hewitt), King William's Town (R. Godfrey).

5i. Var. *fochi* FOREL, 1918, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 155.

Type locality: Durban, NATAL.

6. *Euponera* (*Mesoponera*) *ingesta* Wm. M. WHEELER, See p. 82 (♀).

Type locality: Akenge, BELGIAN CONGO (Lang and Chapin).

BELGIAN CONGO: Niapu; Faradje; Lubila (Lang and Chapin).

7. *Euponera* (*Mesoponera*) *peringueyi* EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 81. ARNOLD, 1915, Ann. South African Mus., XIV, p. 66 (♀), Pl. II, fig. 19.

Ponera peringueyi EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 474 (♀).

Euponera (*Brachypsonera*) *peringueyi* EMERY, 1901, *ibid.*, XLV, p. 47.

Type locality: Nord Hook, CAPE PROVINCE (L. Péringuey).

CAPE PROVINCE: Saldanha Bay (L. Péringuey).

8. *Euponera* (*Mesoponera*) *scolopax* EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 46; 1911, 'Gen. Insect., Ponerinae,' p. 81. SRTZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 371 (♀).

Ponera scolopax EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 473 (♀).

Type locality: CAMEROON (Conradt).

BELGIAN CONGO: Libenge (Schubotz).

¹Specimens received from Dr. H. Brauns.

9. *Euponera* (*Mesoponera*) *senegalensis* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 815, fig. 5 (♀).

Euponera (*Xiphopelta*) *senegalensis* EMERY, 1915, *ibid.*, X, p. 7.

Type locality: Thiès, SENEGAMBIA (F. Silvestri).

10. *Euponera* (*Mesoponera*) *subiridescens* WM. M. WHEELER. See p. 83 (♀).

Type locality: Akenge, BELGIAN CONGO (Lang and Chapin).

BELGIAN CONGO: Medje; Ngayu (Lang and Chapin).

Subgenus 2. *Brachyponera* EMERY

- Euponera* subg. *Brachyponera* EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 43; 1911, 'Gen. Insect., Ponerinae,' p. 83.

Brachyponera BINGHAM, 1903, 'Fauna Brit. India, Hym.,' II, p. 101.

Pachycondyla (part) ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 316.

Ponera (part) ROGER, MAYR, F. SMITH, EMERY, FOREL.

Subgenotype: *Ponera sennaarensis* Mayr, 1862.

11. *Euponera* (*Brachyponera*) *ambigua* (ERN. ANDRÉ) EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, pp. 47 and 49; 1911, 'Gen. Insect., Ponerinae,' p. 84.

Pachycondyla ambigua ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 316 (♀).

DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 32.

Type locality: SIERRA LEONE (Mocquerys).

12. *Euponera* (*Brachyponera*) *sennaarensis* (MAYR) EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 47. FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 75 (♀); 1907, Ann. Mus. Nat. Hungarici, V, p. 5 (♀). MAYR, 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. 1, p. 387. SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 350 (♀, ♀). FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 245 (♀, ♀, ♂). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 130 (♀). FOREL, 1911, Rev. Zool. Afr., II, p. 307 (♀). KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 3 (♀, ♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 84 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, pp. 108 (♀, ♀) and 347 (♀, ♂). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 50 (♀, ♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 318. ARNOLD, 1915, Ann. South African Mus., XIV, p. 72 (♀, ♀, ♂), Pl. II, fig. 21. EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 3 (♀, ♀). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 372 (♀). See p. 84.

Ponera sennaarensis MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 721 (♀); 1863, *ibid.*, XIII, p. 450. ROGER, 1863, 'Verzeich. Formicid.,' p. 16. MAYR, 1878, Verh. Zool. Bot. Ges. Wien, XXVIII, p. 662 (♀). EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 366 (♀). MAGRETTI, 1884, *ibid.*, XXI, p. 338 (♀). EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 557 (♀); 1881, Ann. Mus. Civ. Genova, XVI, p. 528; 1892, *ibid.*, XXXII, p. 111 (♀, ♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 42. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 75. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 125. EMERY, 1895, Ann. Soc. Ent. Belgique, XXXIX, p. 337; 1895, Ann. Mus. Civ. Genova, XXXV, p. 177; 1897, *ibid.*, XXXVIII, p. 597; 1899, Ann. Soc. Ent. Belgique, XLIII, p. 474.

Ponera sorghi ROGER, 1863, Berlin. Ent. Zeitschr., VII, p. 160 (♀); 1863, 'Verzeich. Formicid.,' p. 16. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 450.

Ponera senaarensis MAGRETTI, 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 244 (§). WASMANN, 1894, 'Verzeichn. Myrmecoph. Termitoph. Arthrop.', pp. 202 and 221.

?*Ponera senaarensis* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 474.

Euponera senaarensis MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.', p. 2. BEQUAERT, 1913, Rev. Zool. Afr., II, p. 421. ARNOLD, 1913, Proc. Rhodesia Sc. Assoc., XII, p. 13.

Euponera (Brachyponera) sorghi EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 84 (§).

Type locality: Sennar, ANGLO-EGYPTIAN SUDAN.

SENEGAMBIA: Dakar (C. Alluaud); Thiès (F. Silvestri). FRENCH GUINEA: Konakry; Kindia; Kakoulina (F. Silvestri). SIERRA LEONE: Samlia Falls, River N'Garnie (Mocquerys). GOLD COAST: Kitta; Akra. SOUTHERN NIGERIA: Ibadan; Olokemeji (F. Silvestri). CAMEROON: Metit; Mbalmajo to Ekeneli (G. Schwab). SPANISH GUINEA: Alen (Tessmann). FRENCH CONGO: Cape Lopez; Brazzaville; Mindouli (A. Weiss); Fort Crampel (Schubotz). BELGIAN CONGO: Leopoldville (J. Maes); Avakubi; Niapu; Stanleyville; Medje; Faradje; Thysville; Zambi (Lang and Chapin); Bukama (J. Bequaert). RHODESIA: Bulawayo (G. Arnold). UGANDA: Unyoro Province, region of Lake Albert, Kadjura (C. Alluaud); Gondokoro (F. Werner). BRITISH EAST AFRICA: Voi, Taita district (Alluaud and Jeannel). GERMAN EAST AFRICA: Pemba Island; Fundu Island (Voeltzkow). SOMALILAND: Lower Ganana (V. Bottego); Obbia; Mogadiscio; Gubbet-(Bricchetti-Robecchi). ABYSSINIA: Bela; Ogaden (Ruspoli); Sciotel (Beccari); southern Abyssinia (Ilg); Webi (Bricchetti-Robecchi). ERITREA: Kor Lebka (Magretti); Nefasit; Ghinda (K. Escherich); Keren; Ghinda (F. Silvestri). ANGLO-EGYPTIAN SUDAN: Khartum (I. Trägårdh); Karawaiew; White Nile (Gredler); Kassala; Bahr-el-Salaam (Magretti). ARABIA: Mascat (Biro); Aden; Tes; Cheik Isman (Doria and Beccari).

12. Var. *uginota* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 372 (§).

Type locality: Yaunde, CAMEROON (v. Sommerfeld).

Subgenus 3. *Trachymesopus* EMERY

Euponera subg. *Trachymesopus* EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 84.

Pachycondyla subg. *Pseudoponera* (part) EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 42.

Euponera subg. *Pseudoponera* (part) FOREL, 1901, Ann. Soc. Ent. Belgique, XLV, p. 141.

Pseudoponera (part) BINGHAM, 1903, 'Fauna Brit. India, Hym.,' II, p. 91.

Subgenotype: *Formica stigma* Fabricius, 1804.

13. *Euponera (Trachymesopus) darwini* (FOREL) EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 86. See p. 1009.

Belonopelta darwini FOREL, 1893, Ann. Soc. Ent. Belgique, XXXVII, p. 460 (♀).

Type locality: Port Darwin, AUSTRALIA.

13. Var. *africana* FOREL. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 86. SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 318 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 51 (♀). See p. 85.

Euponera (Pseudoponera) darwini var. *africana* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 51 (♀).

Type locality: Luki, BELGIAN CONGO (A. Jullien).

SOUTHERN NIGERIA: Lagos (F. Silvestri). BRITISH EAST AFRICA: Taveta (Alluaud and Jeannel). BELGIAN CONGO: Stanleyville (Lang and Chapin).

14. *Euponera* (*Trachymesopus*) *lamarki* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 303 (♀).

Type locality: Southern Darbanda, FRENCH CONGO (Decorse).

15. *Euponera* (*Trachymesopus*) *nigeriensis* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 316 (♀).

Type locality: Olokemeji, NIGERIA (F. Silvestri).

GOLD COAST: Aburi (F. Silvestri).

16. *Euponera* (*Trachymesopus*) *suspecta* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 51, fig. 1 (♀).

Type locality: New Moschi, Mt. Kilimanjaro, 800 m., GERMAN EAST AFRICA (Alluaud and Jeannel).

17. *Euponera* (*Trachymesopus*) *wroughtoni* FOREL. EMERY, 1911, 'Gen. Insect., Ponerinæ,' p. 85 (♀, ♂). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 52 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 75 (♀, ♂), Pl. II, fig. 18.

Euponera (*Pseudoponera*) *wroughtoni* FOREL, 1901, Rev. Suisse Zool., IX, p. 341 (♀, ♂).

Type locality: NATAL (R. C. Wroughton).

CAPE PROVINCE: Knysna (H. Brauds). BRITISH EAST AFRICA: Voi, Wataita District (C. Alluaud); Tiwi (Alluaud and Jeannel).

17. Var. *crudelis* FOREL. EMERY, 1911, 'Gen. Insect., Ponerinæ,' p. 85 (♀, ♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 76 (♀, ♂).

Euponera (*Pseudoponera*) *wroughtoni* var. *crudelis* FOREL, 1901, Rev. Suisse Zool., IX, p. 341 (♀, ♂).

Type locality: NATAL (Haviland).

Pseudoponera EMERY

Pachycondyla subg. *Pseudoponera* (part) EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 42.

Pseudoponera EMERY, 1911, 'Gen. Insect., Ponerinæ,' p. 86.

Genotype: *Ponera amblyops* Emery, 1887.

Subgenus **Promyopias** SANTSCHI

Myopias subg. *Promyopias* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 323.

Promyopias EMERY, 1915, *ibid.*, X, p. 26.

Subgenotype: *Myopias* (*Promyopias*) *silvestrii* Santschi, 1914.

1. *Pseudoponera* (*Promyopias*) *asili* (CRAWLEY).

Promyopias asili CRAWLEY, 1916, Entomologist, XLIX, p. 30, fig. (♀).

Type locality: NYASALAND (S. A. Neave).

2. *Pseudoponera* (*Promyopias*) *silvestrii* (SANTSCHI).

Myopias (*Promyopias*) *silvestrii* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 324, fig. 10 (♀).

Promyopias silvestrii EMERY, 1915, *ibid.*, X, p. 26, fig. 13b (♀).

Type locality: Mamou, FRENCH CONGO (F. Silvestri).

Cryptopone EMERY

Cryptopone EMERY, 1892, Ann. Soc. Ent. France, LXI, C. R., p. cclxxv; 1911, 'Gen. Insect., Ponerinae,' p. 88.

?*Amblyopone* (part) MOTSCHULSKY, 1863, Bull. Soc. Nat. Moscou, XXXVI, p. 15.

Genotype: *Amblyopone* (?) *testacea* Motschulsky, 1863.

1. ***Cryptopone angustata*** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 319, fig. 7 (♂).

Type locality: Kakoulima, FRENCH GUINEA (F. Silvestri).

Ponera LATREILLE¹

Ponera LATREILLE, 1805, 'Hist. Nat. Crust. Ins.,' XIII, p. 257. EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 41; 1911, 'Gen. Insect., Ponerinae,' p. 88.

Ponera (part) LATREILLE, 1809, 'Gen. Crust. Ins.,' IV, p. 128.

Formica (part) LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 195.

Bothroponera (part) FOREL, 1900, Journ. Bombay Nat. Hist. Soc., XIII, p. 322. Genotype: *Formica coarctata* Latreille, 1802.

1. ***Ponera abeillei*** ERN. ANDRÉ, 1881, Ann. Soc. Ent. France, (6) I, C. R., p. xlviii (♂).

Type locality: Ajaccio, CORSICA.

1_a. Var. ***cammerunensis*** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 321 (♂).

Type locality: Victoria, CAMEROON (F. Silvestri).

2. ***Ponera coarctata*** (LATREILLE) DALLA TORRE, 1892, Wien. Ent. Zeitg., XI, p. 93.

Formica coarctata LATREILLE, 1802, Bull. Soc. Philomath. Paris, III, p. 65 (♂, ♀), Pl. III, fig. 1.

Type locality: "Jardin du Luxembourg, près de Gentilly," FRANCE (Latreille).

The typical form is found in Southern Europe (northward to Paris), the Mediterranean subregion and the Caucasus.

2_a. Subsp. ***boerorum*** FOREL, 1901, Rev. Suisse Zool., IX, p. 339 (♂); 1907, Ann. Mus. Nat. Hungarici, V, p. 2 (♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 90 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 83 (♂).

Type locality: NATAL, 1600 m. (Haviland).

GERMAN EAST AFRICA: Kibosho (Katona).

2_a. Var. ***aemula*** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 53 (♂).

Ponera dulcis var. *aemula* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 351 (♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 90 (♂).

Type locality: Kibosho, Mt. Kilimanjaro, 1400 m., GERMAN EAST AFRICA (C. Alluaud).

GERMAN EAST AFRICA: Tanga, cave C (C. Alluaud).

2_b. Subsp. ***natalensis*** SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 7 (♂).

Type locality: Richmond, NATAL (I. Trägårdh).

¹*Ponera denticulata* F. Smith, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 90 (♀), Pl. vi, figs. 13 and 14, was described by error from the "Cape of Good Hope." It is a synonym of the Malayan *Odontoponera transperes* (F. Smith).

3. **Ponera cœca** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 322, fig. 9 (♂, ♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 397 (♀).

Type locality: Victoria, CAMEROON (F. Silvestri).

BELGIAN CONGO: St. Gabriel (Kohl).

4. **Ponera diderotii** FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 203 (♂, ♀, ♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 78 (♂, ♀, ♂).

Type locality: Knysna, CAPE PROVINCE (H. Brauns).

5. **Ponera dulcis** FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 2 (♂); 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 245 (♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 90 (♂). FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 203 (♂); 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 213 (♂). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 53, fig. 2 (♂). ALLUAUD and JEANNEL, 1914, Arch. Zool. Gén. Exp., LIII, p. 366. ARNOLD, 1915, Ann. South African Mus., XIV, p. 82 (♂, ♀).

Type locality: Arusha-chini, GERMAN EAST AFRICA (Katona).

NATAL: Durban (Marley). RHODESIA: Bulawayo (G. Arnold). GERMAN EAST AFRICA: New Moschi, Mt. Kilimanjaro, 800 m. (Alluaud and Jeannel). BRITISH EAST AFRICA: Cave A, Shimoni (C. Alluaud). ERITREA: Nefasit (K. Escherich).

- 5i. Var. **uncta** SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 7 (♂).

Type locality: Dukudu, ZULULAND (I. Trägårdh).

6. **Ponera ergatandria** FOREL, 1893, Trans. Ent. Soc. London, p. 365 (♂, ♀, ♂). EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 44. WHEELER, 1908, Bull. American Mus. Nat. Hist., XXIV, p. 125 (♂, ♀, ♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 92. SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 154. FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 213 (♂). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 53 (♂).

Type locality: St. Vincent, WEST INDIES (H. Smith).

CANARY ISLANDS: Tenerife. NATAL: Durban (Marley). BRITISH EAST AFRICA: Shimoni (Alluaud and Jeannel).

Tropicopolitan; introduced in hothouses of temperate regions.

- 6a. Subsp. **cognata** SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 153 (♂). FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 203 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 81 (♂).

Type locality: Cucala, BENGUELA (J. Cruchet).

RHODESIA: Bulawayo (G. Arnold).

- 6b. Subsp. **petri** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 397 (♂).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

7. **Ponera gleadowi** FOREL, in Emery, 1895, Mem. Accad. Sc. Bologna, (5) V, p. 292, footnote (♂).

Type locality: Poona, INDIA (R. C. Wroughton).

- 7a. Subsp. **æthiopica** FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 132 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 91. ARNOLD, 1915, Ann. South African Mus., XIV, p. 83 (♂).

Type locality: Tschafianani, southern ABYSSINIA (de Rothschild).

CAPE PROVINCE: Willowmore (H. Brauns).

8. **Ponera inaudax** SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 83 (♂).

Type locality: Yambuya, BELGIAN CONGO (J. Bequaert).

9. *Ponera incisa* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 320, fig. 8 (♀).

Type locality: Lagos, SOUTHERN NIGERIA (F. Silvestri).

10. *Ponera myrmicarise* WASMANN, 1918, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VII, p. 144 (♀).

Type locality: Soppo, CAMEROON (v. Rothkirch).

11. *Ponera orba* EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 7, fig. 4 (♀).

Type locality: Ghinda, ERITREA (F. Silvestri).

12. *Ponera punctatissima* (part?) ROGER, 1859, Berlin. Ent. Zeitschr., III, p. 246 (♀, ♀). FOREL, 1874, 'Fourmis de la Suisse,' pp. 64, 66 (♀, ♀, ♂). ERN. ANDRÉ, 1882, 'Spec. Hym. Europ. Algérie,' II, pp. 241, 242, 245 (♀, ♀, ♂), Pl. XIV, figs. 15, 16. EMERY, 1895, Mem. Accad. Sc. Bologna, (5) V, p. 294, figs. 9, 10 (♀, ♀); 1901, Ann. Soc. Ent. Belgique, XLV, p. 44; 1909, Deutsch. Ent. Zeitschr., p. 373, fig. 11 (♀, ♀, ♂); 1911, 'Gen. Insect., Ponerinae,' p. 91, Pl. III, fig. 5. SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 320 (♀). See p. 1009.

Ponera androgyna ROGER, 1859, Berlin. Ent. Zeitschr., III, p. 254, footnote (♂).

Ponera contracta (part) MEINERT, 1860, Danske Vidensk. Selsk. Skrift., naturv. math. Afd., (5) V, p. 322 (♀; nec ♀, ♂).

Ponera tarda CHARLALEY, 1877, Ent. Monthly Mag., XIV, p. 162.

Type locality: Rauden, GERMANY (in hothouses).

FRENCH GUINEA: Konakry (F. Silvestri). NIGERIA: Olokemeji; Lagos (F. Silvestri).

13. *Ponera ragusæ* EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 91. See p. 1010.

Ponera ragusæ EMERY, 1894, II Naturalista Siciliano, XIV, p. 28 (♀); 1895, Mem. Accad. Sc. Bologna, (5) V, pp. 292, 297, fig. 18 (♀); 1901, Ann. Soc. Ent. Belgique, XLV, p. 44. FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 2 (♀). SANTSCHI, 1908, Ann. Soc. Ent. France, LXXVII, p. 519 (♀, ♂). EMERY, 1909, Deutsch. Ent. Zeitschr., p. 370, fig. 9a (♀).

Type locality: SICILY (Ragusæ).

EGYPT: Cairo (Boisard). GERMAN EAST AFRICA: Boma Gombe (Katona).

131. Var. *sordida* (SANTSCHI).

Ponera ragusæ var. *sordida* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 54 (♀).

Type locality: Shimoni, BRITISH EAST AFRICA (Alluaud and Jeannel).

13a. Subsp. *bulawayensis* (FOREL) ARNOLD, 1915, Ann. South African Mus., XIV, p. 80 (♀).

Ponera ragusæ subsp. *bulawayensis* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 203 (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

14. *Ponera rothkirchi* WASMANN, 1918, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VII, p. 145 (♀).

Type locality: Soppo, CAMEROON (v. Rothkirch).

15. *Ponera spei* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 423 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 92 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 77 (♀).

Type locality: Mountains of NATAL (R. C. Wroughton).

CAPE PROVINCE: Algoa Bay (H. Brauns).

- 15a. Subsp. **devota** SANTSCHI, 1913, Med. Göteborgs Mus. Zool. Afd., III, p. 5 (♀).
 Type locality: Richmond, NATAL (I. Trägårdh).
 15a1. Var. **sancta** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 213 (♀).
 Type locality: Durban, NATAL (G. Arnold).
 16. **Ponera sulcatinasis** SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 4 (♀, ♀).
 Type locality: Richmond, NATAL (I. Trägårdh).
 16a. Subsp. **durbanensis** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 213 (♀).
 Type locality: Durban, NATAL (G. Arnold).
 17. **Ponera trægaordhi** SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 6 (♀).
 Type locality: Pietermaritzburg, NATAL (I. Trägårdh).

Asphinctopone SANTSCHI

- Asphinctopone* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 318.
 Genotype: *Asphinctopone silvestrii* Santschi, 1914.
 1. **Asphinctopone silvestrii** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 318, fig. 6 (♀).
 Type locality: Olokemeji, NIGERIA (F. Silvestri).

Plectroctena F. SMITH

- Plectroctena* F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 101. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 94.
Formica (part) SPINOLA, 1853, Mem. Accad. Sc. Torino, XIII, p. 79.
 Genotype: *Plectroctena mandibularis* F. Smith, 1858.
 1. **Plectroctena cristata** EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 470 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 129 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 95 (♀). See p. 88.
 Type locality: CAMEROON (Conradt).
 Togo: (Conradt; Schröder). CAMEROON: Mundame (Conradt); Barombi (Preuss). FERNANDO PO: (Conradt). BELGIAN CONGO: Medje; Akenge (Lang and Chapin).
 2. **Plectroctena gabonensis** SANTSCHI, 1919, Bull. Soc. Vaudoise Sci. Nat., (5) LII, p. 336 (♀).
 Type locality: Libreville, FRENCH CONGO (Chalot).
 FRENCH CONGO: Samkita (F. Faure).
 3. **Plectroctena mandibularis** F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 101 (♀, ♂), Pl. VII, figs. 1-5. GERSTÄCKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 355; 1873, in 'v. d. Decken's Reisen in Ost Afrika, Gliederthiere,' p. 346. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 19. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 124. EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 469 (♀ erg.). MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 18. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 95 (♀, ♂), Pl. III, fig. 7; 1912, Annuar. Mus. Zool. Univ. Napoli, N. S., III, No. 26, p. 2 (♀). FOREL, 1913, Ann. Soc. Ent.

Belgique, LVII, p. 108 (♀); 1913, Rev. Zool. Afr., II, p. 308 (♀); 1913, Rev. Suisse Zool., XXI, p. 666 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 54 (♀, ♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 86 (♀, ♀ erg., ♂).

Ponera caffra (KLUG) SPINOLA, 1853, Mem. Accad. Sc. Torino, XIII, p. 69 (without description). EMERY, 1893, Boll. Mus. Zool. Anat. Comp. Torino, VIII, No. 163, p. 2.

Plectroctena caffra ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 41 (♀, ♀, ♂); 1863, 'Verzeich. Formicid.,' p. 19. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 442. EMERY, 1891, Ann. Soc. Ent. France, LX, p. 556 (♀), Pl. xv, figs. 3, 4. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 31. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 74. DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, p. 336. ZAVATTARI, 1909, 'Il Ruwenzori, Parte Scientific.,' I, p. 213 (♂). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 51 (♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 3 (♀, ♂); 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 361 (♀).

Plectroctena caffra subsp. *major* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 74 (♀ erg.).

Plectroctena caffra H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 41.

Plectroctena mandibularis var. *major* EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 95. SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 8 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 54 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 86 (♀, ♀ erg., ♂).

Type locality: Port Natal, NATAL (Gueinzius).

CAMEROON: (Conradt). BELGIAN CONGO: Katanga (Lemaire); valley of the Lubumbashi (Buttgenbach). RHODESIA: Bulawayo (G. Arnold); Lupula River (Elena d'Aosta). GERMAN SOUTHWEST AFRICA: (Lübbert). KALAHARI Desert (L. Schultze). ORANGE FREE STATE. CAPE PROVINCE: Cape Town (Wilms); Mafeking (L. Schultze). NATAL: Durban (Wilms). TRANSVAAL: Makapan (E. Simon); Valdezia (Creux and P. Berthoud); Johannesburg (Dixey and Longstaff). ZULULAND: Umfolosi (I. Trägårdh). DELAGOA: (P. Berthoud). GERMAN EAST AFRICA: Kibonoto, Kilimanjaro (Sjöstedt); Lake Tanganyika (Reichardt); Moschi, Mt. Kilimanjaro, 1120 m. (Alluaud and Jeannel). BRITISH EAST AFRICA: Wanga (v. d. Decken); Nairobi (H. Prell); Fort Hall; Nairobi; Bura, Wa-Taita; River Tchania (Alluaud and Jeannel); Samburu, Wa-Nyika District (C. Alluaud). UGANDA: Ibanda (Duke of Abruzzi).

3. Var. *strigosa* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 469 (♀); 1911, 'Gen. Insect., Ponerinae,' p. 95 (♀). FOREL, 1912, Ent. Mitt. Deutsch. Ent. Mus. Berlin, I, p. 81 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 88 (♀).

Type locality: NATAL.

Changmane, Africa.

4. *Plectroctena minor* EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 556 (♀), Pl. xv, figs. 1, 2. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 31. EMERY, 1902, Rend. Accad. Sc. Bologna, N. S., VI, p. 32 (♀, ♀); 1911, 'Gen. Insect., Ponerinae,' p. 95 (♀, ♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 325. FOREL, 1916, Rev. Suisse Zool., XXIV, p. 398 (♀, ♀). See p. 88 (♀ ergat., ♂).

Type locality: Assinie, IVORY COAST (C. Alluaud).

SIERRA LEONE. GOLD COAST: Aburi (F. Silvestri). CAMEROON: Victoria (F. Silvestri). FERNANDO PO: (Conradt). BELGIAN CONGO: St. Gabriel (Kohl); Akenge; Stanleyville; Niapu (Lang and Chapin).

4. Var. *dentata* SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 150; 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 54 (§).

Plectroctena minor SANTSCHI, 1911, Rev. Zool. Afr., I, p. 205 (§).

Type locality: BENGUELA (J. Cruchet).

UGANDA: Ibanda, Mt. Ruwenzori, 1400 m. (C. Alluaud).

4a. Subsp. *conjugata* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 8 (§).

Type locality: Stamford Hill, NATAL (I. Trägårdh).

NATAL: Charlestown (I. Trägårdh). ZULULAND: (I. Trägårdh).

Myopias ROGER

Myopias ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 39.

Genotype: *Myopias amblyops* Roger, 1861.

1. *Myopias subterranea* (ARNOLD).

Plectroctena subterranea ARNOLD, 1915, Ann. South African Mus., XIV, p. 84 (§, ♀), Pl. III, figs. 23, 23a.

Type locality: Bulawayo, RHODESIA (G. Arnold).

RHODESIA: Shiloh (G. Arnold).

1a. Subsp. *gabonensis* (SANTSCHI).

Plectroctena subterranea subsp. *gabonensis* SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 90 (§).

Type locality: Gaboon, FRENCH CONGO (F. Faure).

Psalidomyrmex ERN. ANDRÉ

Psalidomyrmex ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 313. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 95.

Genotype: *Psalidomyrmex foveolatus* Ern. André, 1890.

1. *Psalidomyrmex foveolatus* ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 314 (§). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 31. STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 129 (§, ♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 96 (§). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 325 (§).

Type locality: SIERRA LEONE (Mocquerys).

FRENCH GUINEA: Kakoulina (F. Silvestri). CAMEROON: (Conradt); Bibundi (Tessmann).

2. *Psalidomyrmex longiscapus* SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 8 (♀).

Type locality: Samkita, FRENCH CONGO (F. Faure).

3. *Psalidomyrmex obesus* WM. M. WHEELER. See p. 92 (§).

Type locality: Medje, BELGIAN CONGO (Lang and Chapin).

4. *Psalidomyrmex procerus* EMERY, 1901, Ann. Soc. Ent. Belgique, XLV, p. 50 (§). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 129 (§). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 96, Pl. III, fig. 8. See p. 90.

Psalidomyrmex foveolatus EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 471 (♀, ♂) (nec André).

Type locality: CAMEROON (Conradt).

Togo: Bismarckburg (Conradt). CAMEROON: Mundame (Conradt). BELGIAN CONGO: Medje; Akenge; Niapu (Lang and Chapin).

5. *Psalidomyrmex reichenspergeri* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 302 (♀); 1914, Deutsch. Ent. Zeitschr., p. 288 (♂). See p. 90.

Type locality: Molundu, CAMEROON (Reichensperger).

CAMEROON: Soppo (v. Rothkirch). BELGIAN CONGO: Akenge (Lang and Chapin).

Cacopone SANTSCHI

Cacopone SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 325.

Genotype: *Cacopone hastifer* Santschi, 1914.

1. *Cacopone hastifer* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 325, fig. 11 (♀).

Type locality: Aburi, GOLD COAST (F. Silvestri).

Leptogenyini Forel

Leptogenys ROGER

Leptogenys ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 41 (*sensu stricto*).

Leptogenys MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 198 (*sensu latiore*). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 97.

Lobopelta MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 733.

Ponera (part) JERDON, F. SMITH, ROGER, BUCKLEY.

Genotype: *Leptogenys falcigera* Roger, 1861.

Subgenus 1. *Leptogenys* ROGER, *sensu stricto*¹

Subgenotype: same as genotype.

1. *Leptogenys bellii* EMERY, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 59 (♀); 1911, 'Gen. Insect., Ponerinae,' p. 99 (♀).

Type locality: Ghinda, ERITREA (Belli).

2. *Leptogenys conradti* FOREL, 1913, Rev. Suisse Zool., XXI, p. 666 (♀).

Type locality: CAMEROON (Conradt).

3. *Leptogenys cribrata* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 20 (♀); 1911, 'Gen. Insect., Ponerinae,' p. 99 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 91 (♀).

Type locality: Vrijburg, CAPE PROVINCE (E. Simon).

4. *Leptogenys crustosa* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 326, fig. 12 (♀).

Type locality: Konakry, FRENCH GUINEA (F. Silvestri).

5. *Leptogenys ferrarii* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 209 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 94 (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

5a. Subsp. *dentatula* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 246 (♀).

Type locality: Katakai, BELGIAN CONGO (Gérard).

¹According to Forel, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 81, *Leptogenys pruinosa* Forel, originally described from Ceylon, was found at Hamburg in wood imported from East Africa.

6. *Leptogenys jeanneli* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 55, fig. 3 (♀). ALLUAUD and JEANNEL, 1914, Arch. Zool. Gén. Exp., LIII, p. 375.

Type locality: Cave B, Kulumuza, Tanga, GERMAN EAST AFRICA (Alluaud and Jeannel).

7. *Leptogenys longiceps* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 328, fig. 13 (♀).

Type locality: Dakar, SENEGAMBIA (F. Silvestri).

8. *Leptogenys maxillosa* (F. SMITH) ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 43 (♀); 1863, 'Verzeich. Formicid.,' p. 19. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 426; 1886, *ibid.*, XXXVI, p. 358. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 46. EMERY, 1911, 'Gen. Insect., Ponerinæ,' p. 99. FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 110 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 89 (♂, ♂), Pl. III, fig. 24. See p. 1011.

Ponera maxillosa F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 93 (♀, ♂).

Type locality: MAURITIUS (Beke).

RHODESIA: Bulawayo (G. Arnold). MALAGASY REGION. WEST INDIES: St. Thomas.

9. *Leptogenys pavesii* EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 111 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 46. EMERY, 1911, 'Gen. Insect., Ponerinæ,' p. 99 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 55 (♀). LAMBORN, 1920, Trans. Ent. Soc. London, (1919), Proc., pp. liv and iv.

Type locality: Obbia, SOMALILAND (Bricchetti-Robecchi).

BRITISH EAST AFRICA: Shimoni (Alluaud and Jeannel); Voi (Alluaud). SOMALILAND: Erdal (Pavesi). GERMAN EAST AFRICA: Lindi (Lamborn).

10. *Leptogenys schwabi* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 208 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 93 (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

11. *Leptogenys stuhlmanni* MAYR, H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 39. FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 51 (♂). EMERY, 1911, 'Gen. Insect., Ponerinæ,' p. 100 (♀). FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 214 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 55 (♀, ♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 91 (♀). LAMBORN, 1920, Trans. Ent. Soc. London, (1919), Proc., p. lv.

Leptogenys (Lobopelta) stuhlmanni MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 198 (♀).

Type locality: Quilimane, MOZAMBIQUE (Stuhlmann).

NATAL: Durban (G. Arnold). BRITISH EAST AFRICA: Mombasa (C. Alluaud); Shimoni; Tiwi; Likoni; Cheteni (Alluaud and Jeannel). NORTH EAST AFRICA: (von Erlanger). GERMAN EAST AFRICA: Lindi (Lamborn).

11a. Subsp. *camerunensis* (STITZ) FOREL, 1916, Rev. Suisse Zool., XXIV, p. 400.

Leptogenys camerunensis STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 130 (♀). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 421. EMERY, 1911, 'Gen. Insect., Ponerinæ,' p. 99 (♀).

Type locality: Barombi, CAMEROON (Preuss).

11a₁. Var. *angusticeps* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 399 (♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

11a₂. Var. *opalescens* WM. M. WHEELER. See p. 94 (♀).

Type locality: Akenge, BELGIAN CONGO (Lang and Chapin).

11b. Subsp. *erythraea* EMERY, 1902, Rend. Accad. Sc. Bologna, N. S., VI, p. 33 (♀). FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 131 (♀, ♂); 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 247 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 100 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 92 (♂).

Leptogenys stuhlmanni EMERY, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 60 (♀) (nec Mayr).

Type locality: Ghinda, ERITREA (Belli).

ERITREA: Nefasit (K. Escherich). ABYSSINIA: Bourka (de Rothschild).

Subgenus 2. *Lobopelta* (MAYR)

Lobopelta MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 733.

Leptogenys subg. *Lobopelta* EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 101.

Subgenotype: *Ponera diminuta* F. Smith, 1857.

12. *Leptogenys (Lobopelta) arnoldi* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 110 (♀, ♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 99 (♀, ♂), Pl. III, figs. 27, 27a, and 28.

Type locality: Plumtree, RHODESIA (G. Arnold).

13. *Leptogenys (Lobopelta) attenuata* (F. Smith) MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 124 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 102 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 97 (♀, ♂).

Ponera attenuata F. Smith, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 91 (♀). ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 14 (♀). MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 447. EMERY, 1893, Boll. Mus. Zool. Anat. Comp. Torino, VIII, No. 163, p. 2.

Lobopelta? attenuata ROGER, 1863, 'Verzeich. Formicid.,' p. 19.

Lobopelta attenuata DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 43.

Type locality: Cape of Good Hope.

CAPE PROVINCE: Algoa Bay (H. Brauns). NATAL: Durban (C. B. Cooper).

PORTUGUESE EAST AFRICA: Delagoa Bay.

13₁. Var. *jagerskiöldi* (SANTSCHI).

Leptogenys (Lobopelta) jagerskiöldi SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 9 (♀).

Leptogenys (Lobopelta) attenuata var. *jagerskiöldi* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 279.

Type locality: Richmond, NATAL (I. Trägårdh).

14. *Leptogenys (Lobopelta) buyssoni* FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 131 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 102 (♀).

Type locality: Hiéka Bourka, southern ABYSSINIA (de Rothschild).

15. *Leptogenys (Lobopelta) castanea* (MAYR) EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 102 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 101 (♀).

Lobopelta castanea MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 734 (♀); 1863, ibid., XIII, p. 427. ROGER, 1863, 'Verzeich. Formicid.,' p. 19. MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidæ, p. 72 (♀), Pl. III, fig. 20; 1878, Verh. Zool. Bot. Ges. Wien, XXVIII, p. 664 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 44.

Leptogenys castanea EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 20 (♀).

Type locality: Cape of Good Hope (Novara Expedition).

CAPE PROVINCE: Algoa Bay (H. Brauns). TRANSVAAL: Makapan (E. Simon).

16. *Leptogenys (Lobopelta) ergatogyna* WM. M. WHEELER. See p. 95 (♀).

Type locality: Medje, BELGIAN CONGO (Lang and Chapin).

17. *Leptogenys (Lobopelta) guineensis* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 329, fig. 14 (♀).

Type locality: FRENCH GUINEA (F. Silvestri).

18. *Leptogenys (Lobopelta) havilandi* FOREL, 1901, Rev. Suisse Zool., IX, p. 332 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 102 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 96 (♀).

Type locality: NATAL (Haviland).

18a. Subsp. *peringueyi* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 210 (♀).

Leptogenys (Lobopelta) peringueyi ARNOLD, 1915, Ann. South African Mus., XIV, p. 96 (♀).

Type locality: Table Mt., CAPE PROVINCE (L. Péringuey).

19. *Leptogenys (Lobopelta) intermedia* EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 102 (♀). FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 214 (♀).

Leptogenys intermedia EMERY, 1902, Rend. Accad. Sc. Bologna, N. S., VI, p. 32 (♀).

Type locality: Willowmore, CAPE PROVINCE (H. Brauns).

NATAL: Durban, Umgeni (G. Arnold).

20. *Leptogenys (Lobopelta) nitida* (F. SMITH) WASMANN, 1899, Deutsch. Ent. Zeitschr., p. 404. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 102 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 110; 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 214 (♀). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 11 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 101 (♀; part).

Ponera nitida F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 92 (♀). ROGER, 1863, 'Verzeich. Formicid.,' p. 23. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 449.

Lobopelta nitida MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 358 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 45.

Leptogenys tenuis STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 376, fig. 1 (♀).

Type locality: Port Natal, NATAL.

NATAL: Estcourt (R. C. Wroughton); Stamford Hill (I. Trägårdh); Shiyvte, 4000 ft. (Haviland). CAPE PROVINCE: Ladismith (H. Brauns). BELGIAN CONGO: Lake Kivu (Schubotz).

20₁. Var. *adpressa* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 214 (♀).

Type locality: Grahamstown, CAPE PROVINCE (Hewitt).

20₂. Var. *ana* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 215 (♀).

Leptogenys (Lobopelta) nitida ARNOLD, 1915, Ann. South African Mus., XIV, p. 101 (♀; part).

Type locality: Willowmore, CAPE PROVINCE (H. Brauns).

- 20s. Var. **gracilis** SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 11 (♀).
Type locality: Lake Kivu, BELGIAN CONGO (Schubotz).
- 20a. Var. **grandior** FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 335 (♀).
Type locality: Krantzklouf, NATAL (Marley).
- 20a. Subsp. **brevinodis** FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 335 (♀).
Type locality: George, CAPE PROVINCE (H. Brauns).
- 20b. Subsp. **insinuata** SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 11 (♀).
Type locality: Richmond, NATAL (I. Trägårdh).
21. **Leptogenys (Lobopelta) parva** FOREL, 1901, Rev. Suisse Zool., IX, p. 332 (♀, ♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 102 (♀, ♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 102 (♀).
Type locality: NATAL (Haviland).
- 21i. Var. **bellua** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 214 (♀).
Type locality: Durban, NATAL (G. Arnold).
- 21s. Var. **dispar** SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 10 (♀).
Type locality: ZULULAND (I. Trägårdh).
22. **Leptogenys (Lobopelta) piroskæ** FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 247 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 102 (♀); 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 3 (♀).
Type locality: Ghinda, ERITREA (K. Escherich; F. Silvestri).
23. **Leptogenys (Lobopelta) sulcinoda** (ERN. ANDRÉ) EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 102 (♀).
Lobopelta sulcinoda ERN. ANDRÉ, 1892, Rev. d'Ent. Caen, XI, p. 48 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 46.
Type locality: Gaboon, FRENCH CONGO.

Odontomachini Mayr

Anochetus MAYR

Anochetus MAYR, 1861, 'Europ. Formicid.,' p. 53. EMERY, 1909, Deutsch. Ent. Zeitschr., p. 374; 1911, 'Gen. Insect., Ponerinae,' p. 107.

Stenomyrme MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 711.

Myrmecia (part) FABRICIUS. *Odontomachus* (part) ILLIGER, F. SMITH, SPINOLA, ROGER.

Genotype: *Odontomachus ghilianii* Spinola, 1851.

1. **Anochetus africanus** (MAYR) FOREL, 1887, Mitth. Schweiz. Ent. Ges., VII, p. 382. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 47. FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 1. SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 351 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 131 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 108 (♀, ♀, ♂). FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 203 (♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 331 (♀). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 372 (♀). See pp. 97 and 1012.

Stenomyrmex africanus MAYR, 1865, 'Reise Novara, Zool., II, Formicidae,' p. 11, footnote (♀). ERN. ANDRÉ, 1892, Rev. d'Ent. Caen, XI, p. 47 (♀, ♀).

Anochetus africanus var. *camerunensis* MAYR, 1896, Ent. Tidskr., XVII, p. 236 (♀, ♂). EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 476 (♀, ♀, ♂).

Type locality: GOLD COAST.

GOLD COAST: Aburi (F. Silvestri). CAMEROON: (Sjåstedt); Mundame (Conradt); Victoria (F. Silvestri); Bibundi (Tessmann); Molundu (Schultze). FRENCH CONGO: Brazzaville (Weiss); Gaboon. BELGIAN CONGO: Medje (Lang and Chapin). GERMAN EAST AFRICA: Kibosho (Katona). SOMALILAND: (C. Keller).

1. Var. *obscuratus* (SANTSCHI) EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 108 (♀, ♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 56 (♀, ♀).

Anochetus madagascariensis var. *obscuratus* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 351 (♀, ♀).

Type locality: Kibosho, Mt. Kilimanjaro, GERMAN EAST AFRICA (C. Alluaud).

GERMAN EAST AFRICA: New Moschi (Alluaud and Jeannel). BRITISH EAST AFRICA: Mwatate, Wa-Taita Province (C. Alluaud).

2. *Anochetus bequaerti* FOREL, 1913, Rev. Zool. Afr., II, p. 309 (♀); 1913, Deutsch. Ent. Zeitschr., Beih., p. 203 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 103 (♀, ♀, ♂). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 400 (♀). See p. 99.

Type locality: Bukama, BELGIAN CONGO (J. Bequaert).

BELGIAN CONGO: (Kohl); Garamba (Lang and Chapin). RHODESIA: Bulawayo (G. Arnold).

2a. Subsp. *abstractus* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 12 (♀).

Type locality: Amanzimtoti, NATAL (I. Trägårdh).

3. *Anochetus concinnus* SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 9 (♀).

Type locality: Lugombe, BELGIAN CONGO (Gérard).

4. *Anochetus estus* Wm. M. WHEELER. See p. 98 (♀).

Type locality: Akenge, BELGIAN CONGO (Lang and Chapin).

5. *Anochetus grandidieri* FOREL. See p. 1013.

5a. Var. *katonaë* FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 1 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 109 (♀).

Type locality: Moschi, GERMAN EAST AFRICA (Katona).

6. *Anochetus levaillanti* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 21 (♀). FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 248 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 108 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 108 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 107 (♀, ♀), Pl. III, figs. 25, 25a, 25b.

Type locality: Hamman's Kraal, TRANSVAAL (E. Simon).

SOUTHERN RHODESIA: Springvale (G. Arnold). ERITREA: Nefasit (K. Escherich).

7. *Anochetus opaciventris* Wm. M. WHEELER. See p. 98 (♀).

Type locality: Akenge, BELGIAN CONGO (Lang and Chapin).

8. *Anochetus parvus* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 330 (♀).

Type locality: Olokemeji, NIGERIA (F. Silvestri).

81. Var. *longiceps* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 331 (♀).

Type locality: Victoria, CAMEROON (F. Silvestri).

9. *Anochetus pellucidus* EMERY, 1902, Rend. Accad. Sc. Bologna, N. S., VI, p. 33 (♂); 1911, 'Gen. Insect., Ponerinae,' p. 109 (♂).

Type locality: CAMEROON (Conradt).

91. Var. *aurifrons* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 351 (♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 109 (♂).

Type locality: FRENCH CONGO.

92. Var. *maynei* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 347 (♀).

Type locality: Congo de Lemba, BELGIAN CONGO (R. Mayné).

10. *Anochetus punctaticeps* MAYR, 1901, Ann. Naturb. Hofmus. Wien, XVI, p. 4 (♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 109 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 105 (♂), Pl. III, fig. 26. See p. 99.

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

BELGIAN CONGO: Babeyru (Lang and Chapin).

11. *Anochetus punctatus* SANTSCHI, 1914, Med. Côtéborgs Mus. Zool. Afd., III, p. 13 (♂).

Type locality: Lake Sibayi, ZULULAND (I. Trägårdh).

111. Var. *occidentalis* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 330 (♂).

Type locality: Victoria, CAMEROON (F. Silvestri).

12. *Anochetus rothschildi* FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 129 (♂, ♂ ?). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 109 (♂).

Type locality: Daouélé, SOMALILAND (de Rothschild).

ABYSSINIA: Diré Daus; Upper Aouache, Endessa (de Rothschild).

13. *Anochetus talpa* FOREL, 1901, Rev. Suisse Zool., IX, p. 351 (♂). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 110 (♂). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 331 (♂). ARNOLD, 1915, Ann. South African Mus., XIV, p. 106 (♂).

Type locality: NATAL (Haviland).

NIGERIA: Ibadan (F. Silvestri).

14. *Anochetus tragaordhi* MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 2 (♂). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 351 (♂). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 400 (♂).

Anochetus tragaordhi EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 110 (♂).

Anochetus tragaordhi KARAWAEW, 1911, Rev. Russe Ent., XI, p. 3 (♂).

Type locality: Khartum, ANGLO-EGYPTIAN SUDAN (I. Trägårdh; Karawaiew).

FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: (Kohl).

Odontomachus LATREILLE

Odontomachus LATREILLE, 1805, 'Hist. Nat. Crust. Ins.,' XIII, p. 257. MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 710. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 111.

Pedetes BERNSTEIN, 1861, Verh. Zool. Bot. Ges. Wien, XI, Sitzb., p. 7.

Formica (part) LINNAEUS, LATREILLE, etc. *Myrmecia* (part) FABRICIUS. *Ponera* (part) LATREILLE, LEPELETIER.

Atta PATTON, 1894, *American Naturalist*, XXVIII, p. 618.

Genotype: *Formica hematoda* Linnaeus, 1758.

1. *Odontomachus assiniensis* EMERY, 1892, *Ann. Soc. Ent. France*, LX, (1891), pp. 558, 560 (♀). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 49. EMERY, 1899, *Ann. Soc. Ent. Belgique*, XLIII, p. 476 (♀). SANTSCHI, 1910, *Ann. Soc. Ent. France*, LXXVIII, (1909), p. 350 (♀). STITZ, 1910, *Mitt. Zool. Mus. Berlin*, V, p. 130 (♀). FOREL, 1910, *Ann. Soc. Ent. Belgique*, LIV, p. 27. EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 113 (♀). FOREL, 1911, *Bull. Soc. Vaudoise Sc. Nat.*, (5) XLVII, p. 362 (♀, ♀); 1913, *Rev. Zool. Afr.*, II, p. 309 (♀); 1913, *Rev. Suisse Zool.*, XXI, p. 666 (♀, ♀, ♂). SANTSCHI, 1914, *Boll. Lab. Zool. Gen. Agrar. Portici*, VIII, p. 332 (♀); 1914, *Deutsch. Ent. Zeitschr.*, p. 288 (♂); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 57 (♀, ♂). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 372 (♀). See p. 100.

Odontomachus intermedius STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exped. (1907-08),' III, p. 378 (♀).

Odontomachus assiniensis subsp. *intermedius* SANTSCHI, 1914, *Med. Göteborgs Mus. Zool. Afd.*, III, p. 14.

Type locality: Assinie, IVORY COAST (C. Alluaud).

GOLD COAST: Aburi (F. Silvestri). TOGO: Bismarekburg (Conradt; Büttner); Misahohe (Baumann). CAMEROON: (Bartsch); Molundu (Schultze); Mundame (Conradt); Soppo (von Rothkirch). FERNANDO PO: (Conradt). FRENCH CONGO: between Fort de Possel and Fort Crampel (Schubotz); Brazzaville (A. Weiss). BELGIAN CONGO: Beni (Bergerhoff); Kimpoko (Büttner); Bakaie, between Nyangwe and Stanleyville (Fauconnet); Akenge; Medje; Ngayu; Niangara; Niapu (Lang and Chapin). GERMAN EAST AFRICA: Buiko (H. Prell); Amani (Zimmer); Buddu Forest (Schubotz). UGANDA: Toro; Unyoro, Mubende Region; Ibando, Mt. Ruwenzori, 1400 m. (C. Alluaud).

1. Var. *aterrimus* Wm. M. WHEELER. See p. 102 (♀).

Type locality: Niapu, BELGIAN CONGO (Lang and Chapin).

1. Var. *furvior* Wm. M. WHEELER. See p. 101 (♀, ♀).

Type locality: Faradje, BELGIAN CONGO (Lang and Chapin).

BELGIAN CONGO: Yakuluku; Stanleyville; Bafwasende; Medje; Ngayu; Akenge; Boyulu; Niangara (Lang and Chapin); Thysville (J. Bequaert).

1. Var. *fuscus* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 372 (♀, ♀).

Type locality: Duma, BELGIAN CONGO (Schubotz).

1a. Subsp. *caffrorum* SANTSCHI, 1914, *Med. Göteborgs Mus. Zool. Afd.*, III, p. 14 (♀).

Type locality: Dukudu, ZULULAND (I. Trägårdh).

1b. Subsp. *fauconneti* FOREL, 1910, *Ann. Soc. Ent. Belgique*, LIV, p. 27 (♀). EMERY, 1911, 'Gen. Insect., Ponerinae,' p. 113 (♀).

Type locality: Bakaie, between Nyangwe and Stanleyville, BELGIAN CONGO (Fauconnet).

2. *Odontomachus hematoda* (LINNAEUS) DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 50. ERN. ANDRÉ, 1895, *Rev. d'Ent. Caen*, XIV, p. 2 (♀). MAYR, 1896, *Ent. Tidskr.*, XVII, p. 238 (♂). EMERY, 1899, *Ann. Mus. Civ. Genova*, XXXIX, p. 500; 1899, *Mem. Accad. Sc. Bologna*, (5) VIII, p. 3 (larva), Pl. I, fig. 4; 1911, 'Gen. Insect.,

Ponerinae,' p. 114, Pl. III, fig. 18. ARNOLD, 1915, Ann. South African Mus., XIV, p. 108 (♂, ♀, ♂). STITZ, 1916, 'Ergebn. Zweite. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 373 (♂, ♀). See pp. 102 and 1013.

Formica haematoda LINNAEUS, 1758, 'Syst. Nat.,' Ed. 10, I, p. 582 (♀). P. F. GMELIN, 1766, 'Onomatol. Hist. Nat.,' III, p. 925. LINNAEUS, 1767, 'Syst. Nat.,' Ed. 12, I, 2, p. 965 (♂). FABRICIUS, 1775, 'Syst. Ent.,' p. 395 (♀). PH. L. MÜLLER, in LINNAEUS, 1775, 'Vollst. Natursyst.,' V, 2, p. 916 (♀), Pl. XXVII, fig. 17. FABRICIUS, 1781, 'Spec. Insect.,' I, p. 494 (♂); 1787, 'Mantissa Insect.,' I, p. 311 (♀). GMELIN, in LINNAEUS, 1790, 'Syst. Nat.,' Ed. 13, I, 5, p. 2803 (♂). CHRIST, 1791, 'Naturg. d. Insect.,' p. 516 (♀). OLIVIER, 1791, 'Encycl. Méthod. Insect.,' VI, p. 502 (♀). FABRICIUS, 1793, 'Ent. Syst.,' II, p. 364 (♀). LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 192 (♀).

Formica mazillosa DEGEER, 1773, 'Mém. Hist. Ins.,' III, p. 601 (♀), Pl. XXXI, figs. 3-5. GÖZE, in DeGeer, 1780, 'Abh. Gesch. Insekt.,' III, p. 390 (♀), Pl. XXXI, figs. 3-5. RETZIUS, 1783, 'Gen. et Spec. Insect.,' p. 75 (♀).

Formica unispinosa FABRICIUS, 1793, 'Ent. Syst.,' II, p. 359 (♀). LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 193 (♀), Pl. VIII, fig. 53.

Myrmecia haematoda FABRICIUS, 1804, 'Syst. Piez.,' p. 425 (♀).

Myrmecia unispinosa FABRICIUS, 1804, *ibid.*, p. 423 (♀). OLIVIER, 1811, 'Encycl. Méthod. Insect.,' VIII, p. 114 (♀).

Odontomachus haematodes LATREILLE, 1805, 'Hist. Nat. Crust. Ins.,' XIII, p. 257 (♀). ILLIGER, 1807, Mag. f. Insectenk., VI, p. 194. LEPELETIER, 1836, 'Hist. Nat. Ins. Hym.,' I, p. 187 (♀). F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 76 (♀, ♀, ♂), Pl. v, figs. 4-7. GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 262; 1859, Arch. f. Naturg., XXV, 2, p. 431. ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 24 (♀, ♀, ♂). GERSTÄCKER, in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 503 (♀, ♂). MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 711; 1863, *ibid.*, XIII, p. 436. ROGER, 1863, 'Verzeich. Formicid.,' p. 22. MAYR, 1867, Tijdschr. v. Ent., V, p. 79 (♀, ♀); 1867, Journ. Mus. Godeffroy, XII, p. 85. W. F. KIRBY, 1884, Ann. Mag. Nat. Hist., (5) XIII, p. 405. MAGRETTI, 1884, Ann. Mus. Civ. Genova, XXI, p. 538 (♀); 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 244 (♀). WASMANN, 1892, Wien. Ent. Zeitg., XI, p. 317. EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), pp. 557 and 561 (♀, ♀). FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 74. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 126. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 22. BINGHAM, 1903, 'Fauna Brit. India, Hym.,' II, p. 47 (♀, ♀). MAYR, 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. I, p. 387. ZAVATTARI, 1909, 'Il Ruwenzori, Parte Scientif.,' I, p. 213 (♀, ♂). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXXVIII, (1909), p. 350 (♀, ♀). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 51 (♀, ♂). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 130 (♀, ♀). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 4 (♀, ♂); 1912, Mitt. Naturh. Mus. Hamburg, XXIX, p. 179 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, pp. 108 and 347 (♀, ♀); 1913, Rev. Zool. Afr., II, p. 309 (♀). LAMBORN, 1914, Trans. Ent. Soc. London, (1913), p. 442 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicid.,' p. 58 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 331 (♀, ♀); 1920, 'Études Maladies Parasites Cacaoyer S. Thomé,' X, p. 2 (♀).

Odontomachus unispinosus ILLIGER, 1807, Mag. f. Insectenk., VI, p. 194. LEPELETIER, 1836, 'Hist. Nat. Ins. Hym.,' I, p. 187 (♀, ♀). F. SMITH, 1853, Trans. Ent. Soc. London, (2) II, Proc., p. 134.

Ponera (Odontomachus) hæmatoda LATREILLE, 1809, 'Gen. Crust. Ins.,' IV, p. 128.

Ponera (Odontomachus) unispinosa LATREILLE, 1809, *ibid.*, IV, p. 128.

Ponera hæmatoda LEPELETIER, 1825, 'Encycl. Méthod. Insect.,' X, p. 184 (♀).

Ponera unispinosa LEPELETIER, 1825, *ibid.*, X, p. 184 (♀).

Formica (Odontomachus) hæmatodes BLANCHARD, in Cuvier, 1849, 'Règne Animal,' Ed. 3, Insect., Pl. cxvii, fig. 6 (♀).

Odontomachus similimus F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 80 (♀), Pl. v, figs. 8-9; 1860, Journ. Proc. Linn. Soc. London, Zool., IV, Suppl., p. 72.

Odontomachus hæmatodes var. *microcephalus* EMERY, 1890, Bull. Soc. Ent. Italiana, XXII, p. 104, Pl. v, fig. 1 (mermithergate).

Type locality: "in America meridionali."

Tropicopolitan. SENEGAMBIA: Thiès (F. Silvestri). FRENCH GUINEA: Kindia (F. Silvestri). LIBERIA: Junk River (H. Brauns). IVORY COAST: Assinie (C. Alluaud). GOLD COAST: Kitta (H. Brauns); Aburi (F. Silvestri). NIGERIA: Ibadan; Lagos; Olokemeji (F. Silvestri); Oni Camp, 70 miles east of Lagos (Lamborn). CAMEROON: (H. Brauns); Barombi (Freyer); Bibundi (Tessmann); Yaunde (Zenker); Moliwe region (Conradt); Victoria (F. Silvestri). FERNANDO PO: (Conradt). SPANISH GUINEA: Alen (Tessmann). SAN THOMÉ: (de Seabra). FRENCH CONGO: Ogowe (Mocquers); Brazzaville (A. Weiss). BELGIAN CONGO: Kimpoko (Büttner); Boma (Leboutte); Leopoldville (Lamarche); Congo da Lemba (R. Mayné); Libenge (Schubotz); Mayombe (de Briey); Duma (Montchal); Stanleyville; Malela; Faradjé; Zambi; Avakubi; Leopoldville; Vankerkhovenville; Garamba; Akenge (Lang and Chapin); Matadi; Katala (J. Bequaert). SOUTHERN RHODESIA: Bulawayo (G. Arnold). BECHUANALAND: Severelela; Khakhea to Kang (L. Schultze). TRANSVAAL: Hamman's Kraal (E. Simon); Lydenburg. DELAGOA: (Liengme). MOZAMBIQUE: (Peters). UGANDA: Ibanda; Kitagueti; Butiti; Nakitawa (Duke of Abruzzi); Entebbe (H. Schultze). BRITISH EAST AFRICA: Shimoni; Kisumu; Mombasa (Alluaud and Jeannel). ABYSSINIA: (Ilg); Dimé to Bass Narok (V. Bottego). ANGLO-EGYPTIAN SUDAN: Khor Attar (F. Werner); Bahr-el-Salaam (Magretti). ERITREA: Sogodas; Kor Lebka (Magretti).

21. Var. *stanleyi* Wm. M. WHEELER. See p. 102 (♀).

Type locality: Stanleyville, BELGIAN CONGO (Lang and Chapin).

22. Var. *trogodytes* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 58 (♀).

Type locality: Shimoni, BRITISH EAST AFRICA (Alluaud and Jeannel).

PSEUDOMYRMINE Emery

Pseudomyrmini Emery

Tetraponera F. SMITH

Tetraponera F. SMITH, 1852, Ann. Mag. Nat. Hist., (2) IX, p. 44; 1877, Trans. Ent. Soc. London, p. 68. DONISTHORPE, 1916, Ent. Record, XXVIII, p. 244.

Sima ROGER, 1863, Berlin. Ent. Zeitschr., VII, p. 178. EMERY, 1915, Zool. Anzeiger, XLV, pp. 265-266; 1917, Bull. Soc. Ent. France, p. 94.

Tetraponera EMERY (as a subgenus). *Pseudoponera* (part) SMITH. *Eciton* JERDON (nec Latreille).

Genotype: *Eciton nigrum* Jerdon, 1851 = *Tetraponera atrata* Smith, 1852.

1. *Tetraponera ambigua* (EMERY).

Sima ambigua EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 23 (♀, ♀). FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 89 (without description); 1901, *ibid.*, X, p. 311; in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 564; in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 5 (♀, ♀, ♂).

Sima (Tetraponera) ambigua ARNOLD, 1916, Ann. South African Mus., XIV, p. 185 (♀, ♀, ♂).

Type locality: Hamman's Kraal, TRANSVAAL (E. Simon).

TRANSVAAL: Makapan (E. Simon). BECHUANALAND: Kang to Kgokong (L. Schultze). MOSSAMEDES: between the Cubango and Cuito Rivers (Baum and Van der Kellen). SOUTHERN AETHIOPIA: (Ilg).

1a. Subsp. *erythræa* (EMERY).

Sima ambigua subsp. *erythræa* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 23 (♀).

Type locality: Aden, ARABIA (E. Simon).

1b. Subsp. *rhodesiana* (FOREL).

Sima ambigua subsp. *rhodesiana* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 112 (♀). ARNOLD, 1914, Proc. Rhodesia Sc. Ass., XIII, p. 31, Pl., fig. 1 (♀).

Sima (Tetraponera) ambigua subsp. *rhodesiana* ARNOLD, 1916, Ann. South African Mus., XIV, p. 186 (♀, ♀).

Type locality: Plumtree, SOUTHERN RHODESIA (G. Arnold).

SOUTHERN RHODESIA: Bembesi (G. Arnold).

2. *Tetraponera andrei* (MAYR).

Sima andrei MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 144 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 179 (♀).

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA (H. Brauns).

3. *Tetraponera anthracina* (SANTSCHI). See p. 106.

Sima anthracina SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 355 (♀); 1910, *ibid.*, LXXIX, p. 352, fig. 3 (♀). FOREL, 1911, Rev. Zool. Afr., I, p. 275 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 350 (♀); 1916, Rev. Suisse Zool., XXIV, p. 403 (♀).

Type locality: Combra Tora, FRENCH CONGO (A. Weiss).

BELGIAN CONGO: St. Gabriel (Kohl); Kisantu; Congo da Lemba (R. Mayné); Thysville; Lubutu (Bequaert); Stanleyville (Lang and Chapin).

4. *Tetraponera bifoveolata* (MAYR).

Sima bifoveolata MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 146 (♀); in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 4 (♀).

Sima (Tetraponera) bifoveolata ARNOLD, 1916, Ann. South African Mus., XIV, p. 184 (♀).

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA (H. Brauns).

ANGLO-EGYPTIAN SUDAN: Kaka, White Nile (I. Trägårdh). ZANZIBAR: (H. Brauns).

A subspecies is known from Palestine.

4a. Subsp. *maculifrons* (SANTSCHI).

Sima foveolata subsp. *maculifrons* SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 162 (♀).

Type locality: Obock, FRENCH SOMALILAND (Maindrón).

5. **Tetraponera capensis** (F. SMITH) ROGER, 1863, 'Verzeich. Formicid.,' p. 24.

Pseudomyrma capensis F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 160 (♀). MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 451. F. SMITH, 1877, Trans. Ent. Soc. London, p. 60.

Sima capensis DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 53.

Type locality: Cape of Good Hope.

6. **Tetraponera claveauui** (SANTSCHI).

Sima claveauui SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 304 (♀).

Type locality: Senegal (Claveau).

7. **Tetraponera clypeata** (EMERY).

Sima clypeata EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 361 (♀), Pl. xvii, figs. 4 and 5. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 53. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 24 (♀); 1899, Mem. Accad. Sc. Bologna, (5) VIII, p. 4 (larva).

Sima (Tetraponera) clypeata ARNOLD, 1916, Ann. South African Mus., XIV, p. 182 (♀, ♀), Pl. v, figs. 50, 50a.

Type locality: Cape of Good Hope (L. Péringuey).

CAPE PROVINCE: Cape Town; Matjesfontein (E. Simon).

7a. Subsp. **braunsi** (FOREL).

Sima braunsi FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 112 (♀, ♀).

Sima (Tetraponera) clypeata subsp. *braunsi* ARNOLD, 1916, Ann. South African Mus., XIV, p. 184 (♀, ♀), Pl. v, fig. 49.

Type locality: Willowmore, CAPE PROVINCE (H. Brauns).

7a. Var. **durbanensis** (FOREL).

Sima braunsi var. *durbanensis* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 218 (♀).

Sima (Tetraponera) clypeata subsp. *braunsi* var. *durbanensis* ARNOLD, 1916, Ann. South African Mus., XIV, p. 184 (♀).

Type locality: Durban, NATAL (G. Arnold).

7a. Var. **equidentata** (ARNOLD).

Sima (Tetraponera) clypeata subsp. *braunsi* var. *equidentata* ARNOLD, 1916, Ann. South African Mus., XIV, p. 184 (♀), Pl. v, fig. 48.

Type locality: Cape Town, CAPE COLONY (G. Arnold).

8. **Tetraponera emeryi** (FOREL).

Sima capensis EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 23 (♀) (*nec* F. Smith).

Sima emeryi FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 367 (♀).

Sima (Tetraponera) emeryi ARNOLD, 1916, Ann. South African Mus., XIV, p. 187 (♀).

Type locality: Pretoria, TRANSVAAL (E. Simon).

9. **Tetraponera encephala** (SANTSCHI).

Sima encephala SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 84 (♀).

Type locality: Saint Louis, SENEGAMBIA (Claveau).

10. **Tetraponera gerdæ** (STITZ).

Sima gerdæ STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 381, fig. 4 (♀).

Type locality: Amsari, GERMAN EAST AFRICA (Vosseler).

11. **Tetraponera le moulti** (SANTSCHI).

Sima le moulti SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 375 (♀).

Type locality: Fort Crampel, FRENCH CONGO (Le Moul't).

12. **Tetraponera liengmei** (FOREL).

Sima liengmei FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 88 (♀, ♂).

Sima (*Tetraponera*) *liengmei* ARNOLD, 1916, Ann. South African Mus., XIV, p. 181 (♀, ♂).

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA (Liengme).

13. **Tetraponera mayri** (FOREL).

Sima mayri FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 53 (♀).

Type locality: CAMEROON (H. Brauns).

14. **Tetraponera mocquersyi** (ERN. ANDRÉ).

Sima mocquersyi ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 319 (♀). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 54. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 23 (♀); 1895, Ann. Mus. Civ. Genova, XXXV, p. 178. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 146. STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 131 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 352, fig. 2 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 334 (♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 403 (♀). SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 10.

Type locality: SIERRA LEONE (Mocquersys).

GOLD COAST: Chama (H. Brauns). TOGO: Bismarckburg (Conradt). FRENCH GUINEA: Kakoulima (F. Silvestri). SPANISH GUINEA: Alen (Tessmann). FRENCH CONGO: Benda (Charleux). BELGIAN CONGO: (Kohl). ABYSSINIA: Arussi Galla, Canale Gudda (V. Bottego).

14a. Var. **elongata** (STITZ).

Sima mocquersyi var. *elongata* STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exped. (1907-08)', III, p. 378, fig. 2 (♀).

Type locality: Buddu Forest, GERMAN EAST AFRICA (N. W. of Bukoba) (Schubotz).

14b. Var. **lepida** WM. M. WHEELER. See p. 106 (♀, ♀).

Type locality: Faradje, BELGIAN CONGO (Lang and Chapin).

BELGIAN CONGO: Yakuluku; Garamba (Lang and Chapin).

14c. Var. **lutea** (STITZ).

Sima mocquersyi var. *lutea* STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exped. (1907-08)', III, p. 381, fig. 3 (♀, ♀).

Type locality: Kwidjwi Island, Lake Kivu, BELGIAN CONGO (Schubotz).

14a. Subsp. **emacerata** (SANTSCHI). See p. 107.

Sima mocquersyi subsp. *emacerata* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 352, fig. 1 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae', p. 70 (♀, ♀).

Sima emacerata SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 9 (♀).

Type locality: Nakuru, Rift Valley, BRITISH EAST AFRICA (C. Alluaud).

CENTRAL UGANDA: (C. Alluaud). BELGIAN CONGO: Stanleyville; Faradje (Lang and Chapin); Lubutu; Kasonsero on the Semliki River (J. Bequaert); Ngazi (Elskens).

15. **Tetraponera natalensis** (F. SMITH) ROGER, 1863, 'Verzeich. Formicid.', p. 24. F. SMITH, 1877, Trans. Ent. Soc. London, p. 69.

Pseudomyrma natalensis F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 160 (♀). MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 452. F. SMITH, 1876, Trans. Ent. Soc. London, p. 604.

Sima capensis MAYR, 1866, Verh. Zool. Bot. Ges. Wien, XVI, p. 906 (♀), Pl. XX, fig. 14. EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 113 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 53. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 43 (*nec Sima capensis* Smith).

Pseudomyrma capensis GERSTÄCKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 355; 1873, in 'v. d. Decken's Reisen in Ostafrika, Gliederthiere,' p. 357 (♀, ♀).

Sima natalensis EMERY, 1892, Zool. Anzeiger, XV, p. 237; 1895, Ann. Soc. Ent. France, LXIV, p. 22 (♀, ♀); 1899, Mem. Accad. Sc. Bologna, (5) VIII, p. 4 (larva), Pl. II, fig. 7. FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 26 (♀); 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 367 (♀). SANTSCI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 15 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 176 (♀, ♀).

Type locality: Port Natal, NATAL.

South Africa generally (G. Arnold). CAPE PROVINCE: (H. Brauns). NATAL: Durban (Marley); Weenen District (J. M. Hutchinson); Stamford Hill (I. Trägårdh). ZULULAND: Umfolosi (I. Trägårdh). TRANSVAAL: Makapan; Hamman's Kraal (E. Simon). GERMAN EAST AFRICA: Mto-ya-Kifaru (Katona); Ndara; Ugono Mts. (v. d. Decken). SOMALILAND: Obbia (Bricchetti-Robecchi).

15. Var. *obscurata* (EMERY).

Sima natalensis var. *obscurata* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 22 (♀). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 5 (♀); 1911, Sitzb. Bayer. Akad. Wiss., p. 274; 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 368 (♀); 1915, *ibid.*, (5) L, p. 336 (♀).

Sima natalensis subsp. *obscurata* ARNOLD, 1916, Ann. South African Mus., XIV, p. 177 (♀, ♀, ♂).

Type locality: Cape of Good Hope.

CAPE PROVINCE: Algoa Bay (H. Brauns). NATAL: Durban (C. B. Cooper). NORTHERN RHODESIA: shore of the Zambesi River (Fr. Steiner).

15. Var. *usambarensis* (FOREL).

Sima natalensis var. *usambarensis* FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 367 (♀). SANTSCI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 70 (♀, ♀).

Type locality: Monga, GERMAN EAST AFRICA (H. Prell).

GERMAN EAST AFRICA: Kilema, Mt. Kilimanjaro, 1440 m.; Kibosho (Alluaud and Jeannel). BRITISH EAST AFRICA: Moschi (Zimmer); Mombasa; Shimoni; Nairobi, Kikuyu; Voi, Taita; Maji-ya-chumvi, Wa-Nyika (Alluaud and Jeannel).

15a. Subsp. *caffra* (SANTSCI).

Sima natalensis subsp. *caffra* SANTSCI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 15 (♀, ♀, ♂). TRÄGÅRDH, *ibid.*, p. 43.

Sima natalensis subsp. *obscurata* var. *caffra* ARNOLD, 1916, Ann. South African Mus., XIV, p. 178 (♀, ♀, ♂).

Type locality: Dukudu, ZULULAND (I. Trägårdh).

ZULULAND: Lake Sibayi (I. Trägårdh).

15b. Subsp. *cuitensis* (FOREL).

Sima natalensis subsp. *cuitensis* FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 368 (♀).

Sima natalensis FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 311; in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 564.

Type locality: Between the Cubango River and Cuito River, MOSSAMEDES (Baum and Van der Kellen).

15b1. Var. *bulawayana* (FOREL).

Sima natalensis subsp. *cuitensis* var. *bulawayana* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 112 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 179 (♀, ♀).

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

SOUTHERN RHODESIA: Plumtree; Bembesi; Victoria Falls (G. Arnold).

16. *Tetraponera oberbecki* (FOREL).

Sima oberbecki FOREL, 1911, Rev. Zool. Afr., I, p. 275 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 350 (♀); 1916, Rev. Suisse Zool., XXIV, p. 403 (♀). SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 9.

Type locality: Congo da Lemba, BELGIAN CONGO (R. Mayné).

BELGIAN CONGO: Leopoldville (Kohl); Elisabethville (Leprieux). FRENCH CONGO: Samkita (F. Faure).

17. *Tetraponera ophthalmica* (EMERY). See p. 107.

Sima ophthalmica EMERY, 1912, Ann. Soc. Ent. Belgique, LVI, p. 98 (♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 403 (♀).

Type locality: CAMEROON (Conradt).

BELGIAN CONGO: Bengamisa; St. Gabriel; Stanleyville (H. Kohl); Thysville (J. Bequaert). CAMEROON: Batanga (G. Schwab).

18. *Tetraponera penzigi* (MAYR).

Sima penzigi MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 10 (♀, ♀, ♂). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 5 (♀).

Sima penzigi SJÖSTEDT, 1908, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, pp. 116-117.

Sima (Tetraponera) penzigi ARNOLD, 1916, Ann. South African Mus., XIV, p. 180 (♀, ♀, ♂).

Type locality: Mt. Kilimanjaro, Kahe, GERMAN EAST AFRICA (Sjöstedt).

GERMAN EAST AFRICA: Tanga, Usambara (Sjöstedt). BECHUANALAND: Lehututu (L. Schultze). ABYSSINIA: (Penzig).

18a. Subsp. *continua* (FOREL).

Sima penzigi subsp. *continua* FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 138 (♀).

Type locality: Gotta, southern ABYSSINIA (de Rothschild).

19. *Tetraponera prelli* (FOREL).

Sima prelli FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 365 (♀).

Type locality: Monga, GERMAN EAST AFRICA (H. Prell).

19a. Var. *odiosa* (FOREL).

Sima prelli var. *odiosa* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 403 (♀).

Type locality: BELGIAN CONGO (Kohl).

20. *Tetraponera schulthessi* (SANTSCHI).

Sima schulthessi SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 249 (♀).

Type locality: Rikatla, Delagoa Bay, PORTUGUESE EAST AFRICA (Junod).

21. *Tetraponera triangularis* (STITZ).

Sima triangularis STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 131 (♀).

Type locality: Alen, SPANISH GUINEA (Tessmann).

21a. Subsp. *illota* (SANTSCHI).

Sima triangularis subsp. *illota* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 334 (♀, ♀).

Type locality: Olokemeji, SOUTHERN NIGERIA (F. Silvestri).

Viticicola Wm. M. WHEELER

Viticicola Wm. M. WHEELER, 1920, Psyche, XXVII, p. 53.

Genotype: *Sima tessmanni* Stitz, 1910.

1. ***Viticicola tessmanni*** (STITZ). See p. 109 (♀, ♀, ♂).

Sima tessmanni STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 131, fig. 2 (♀). SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 84 (♀).

Type locality: Alen, SPANISH GUINEA (Tessmann).

Belgian Congo: Medje (Lang and Chapin).

1b. Var. ***castanea*** Wm. M. WHEELER. See p. 112 (♀, ♀).

Type locality: Avakubi, BELGIAN CONGO (Lang and Chapin).

Pachysima EMERY

Sima subg. *Pachysima* EMERY, 1912, Ann. Soc. Ent. Belgique, LVI, p. 97.

Genotype: *Tetraponera aethiops* F. Smith, 1877.

1. ***Pachysima aethiops*** (F. SMITH). See p. 114.

Tetraponera aethiops F. SMITH, 1877, Trans. Ent. Soc. London, p. 71 (♀).

Sima aethiops EMERY, in Dalla Torre, 1893, 'Cat. Hym.,' VII, p. 53. H. KOHL, 1909, Natur u. Offenbarung, LV, p. 167. SANTSCHI, 1911, Rev. Zool. Afr., I, p. 207 (♂). FOREL, 1913, *ibid.*, II, p. 352 (♀). LAMBORN, 1914, Trans. Ent. Soc. London, (1913), pp. 442 and 493 (♀). H. SCHMITZ, 1915, Deutsch. Ent. Zeitschr., p. 500.

Sima spininoda ERN. ANDRÉ, 1892, Rev. d'Ent. Caen, XI, p. 51 (♀, ♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 55. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 144. H. KOHL, 1909, Natur u. Offenbarung, LV, pp. 99, 103, 104, 106, 108, and 167. STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 131 (♀, ♀); 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exped. (1907-08),' III, p. 381 (♀, ♀).

Sima (Pachysima) aethiops EMERY, 1912, Ann. Soc. Ent. Belgique, LVI, p. 97, figs. 2 and 3 (♀, ♀, larva). FOREL, 1913, Rev. Suisse Zool., XXI, p. 668 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 350 (♀); 1916, Rev. Suisse Zool., XXIV, p. 403 (♀, ♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 174 (♀, ♀, ♂).

Sima (Pachysima) spininoda STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 374 (♀, ♀, ♂).

Type locality: South Africa.

SOUTHERN NIGERIA: Oni Camp, east of Lagos (Lamborn). FERNANDO PO: (Conradt). CAMEROON: Bipindi (Zenker); Mundame (Conradt); Bibundi (Tessmann); Metit (G. Schwab). SPANISH GUINEA: Eloby Island (H. Brauns); Alen (Tessmann). FRENCH CONGO: Samkita (F. Faure). BELGIAN CONGO: Kibombo (J. Bequaert); Watikala; Tshopo River; Candolo; Okiavo-Lindi River; St. Gabriel (Kohl); Mawambi to Avakubi; Duma; Libenge (Schubotz); Avakubi; Stanleyville; Ambelokudi; Isangi; Panga; Medje; Bafwabaka (Lang and Chapin).

2. ***Pachysima latifrons*** (EMERY). See p. 120.

Sima (Pachysima) latifrons EMERY, 1912, Ann. Soc. Ent. Belgique, LVI, p. 98, fig. 4 (♀). FOREL, 1913, *ibid.*, LVII, p. 350 (♀). SANTSCHI, 1914, Deutsch. Ent. Zeitschr., p. 288 (♀, ♀, ♂).

Sima æthiops FOREL, 1913, Rev. Zool. Afric., II, p. 315 (♀).

Type locality: Gaboon, FRENCH CONGO.

BELGIAN CONGO: Kondué (Luja); Niangara (Lang and Chapin). CAMEROON: Tiko near Victoria (F. Silvestri). FRENCH CONGO: Lobay (Ringensbach).

MYRMICINÆ Lepeletier

Myrmicini Emery

Cratomyrmex EMERY

Cratomyrmex EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 572.

Genotype: *Cratomyrmex regalis* Emery, 1892.

1. *Cratomyrmex regalis* EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 572 (♀), Pl. xv, fig. 16. SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 307, fig. 3 (♂, ♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 335 (♂).

Messor (*Cratomyrmex*) *regalis* SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 377 (♂).

Type locality: Benue, NIGERIA.

LOWER DAHOMY: Aguagua (Roubaud). NIGERIA: Lagos; Olokemeji (F. Silvestri). CONGO: (Cabra).

1. Var. *rubeus* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 308 (♂).

Type locality: LOWER DAHOMY (Le Moullet).

12. Var. *sculpturatus* (STITZ).

Cratomyrmex sculpturatus STITZ, 1916, 'Wiss. Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 378, fig. 2 (♂).

Type locality: Fort de Possel to Fort Crampel, FRENCH CONGO (Schubotz).

FRENCH CONGO: Falls of the Nana River near Fort Crampel (Haberer).

Pheidolini Emery

Messor FOREL

Aphænogaster subg. *Messor* FOREL, 1890, Ann. Soc. Ent. Belgique, XXXIV, C.R., p. lxxviii.

Messor EMERY, 1908, Deutsch. Ent. Zeitschr., p. 437.

Formica (part) FABRICIUS, Gmelin, Latreille, etc. *Myrmica* (part) FÖRSTER. *Atta* (part) SMITH, ROGER. *Aphænogaster* (part) ERN. ANDRÉ, DALLA TORRE, EMERY, MAYR, etc. *Stenamma* (part) EMERY, FOREL, MAYR, etc.

Genotype: *Formica barbara* Linnaeus.

1. *Messor barbarus* (LINNÆUS) EMERY, 1908, Deutsch. Ent. Zeitschr., p. 442 (♂, ♀, ♂).

Formica barbara LINNÆUS, 1767, 'Syst. Nat.,' Ed. 12, I, 2, p. 962 (♂).

Aphænogaster barbara MAGRETTI, 1877, Ann. Mus. Civ. Genova, IX, p. 373 (♂, ♀, ♂); 1884, *ibid.*, XXI, p. 541 (♂); 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 245 (♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 99.

Aphænogaster (*Messor*) *barbara* FOREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 352 (♂).

Stenamma (*Messor*) *barbarum* FOREL, 1904, Ann. Mus. Zool. Ac. Sc. St. Pétersbourg, VIII, (1903), p. 373 (♀).

Type locality: Barbary.

The following localities probably all refer to some of the subspecies or varieties mentioned below.

ANGLO-EGYPTIAN SUDAN: El Hefera, Settit; Metemma; Sebderat (Magretti); Godo Burka (Kachovskij). SOMALILAND: (C. Keller). ERITREA: Keren; Kor Lebka (Magretti).

1. Var. *punctatus* (FOREL).

Aphenogaster barbara var. *punctata* FOREL, 1886, Journ. Asiat. Soc. Bengal, LV, pt. 2, p. 248 (♀).

Aphenogaster (Messor) *barbara* var. *punctata* EMERY, 1891, 'Explor. Tunisie, Fourmis,' p. 12. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 89.

Messor barbarus subsp. *punctatus* REICHENSBERGER, 1913, Zool. Jahrb. Abt. Syst., XXXV, pp. 187, 198, 201, and 203. WASMANN, 1918, Tijdschr. v. Ent., LX, (1917), p. 396.

Type locality: Kashmir, INDIA.

ABYSSINIA: Abuker; Diré Dava; Harar (Kristensen); southern Abyssinia (Ilg).

1a. Subsp. *latinodus* SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, p. 93, fig. 2, 2 (♀).

Type locality: East Africa (?) (Reichensperger).

1b. Subsp. *lubberti* FOREL, in Schultze, 1910, 'Forschungereise in Südafrika,' IV, p. 13 (♀).

Messor barbarus subsp. *lubberti* ARNOLD, 1920, Ann. South African Mus., XIV, p. 411 (♀).

Type locality: Okahandja, GERMAN SOUTHWEST AFRICA (Peters).

GERMAN SOUTHWEST AFRICA: (Lübbert).

1c. Subsp. *ruginodis* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 374, fig. 1 (♀).

Type locality: Fort Crampel, FRENCH CONGO (Schubotz).

1d. Subsp. *semirufus* (ERN. ANDRÉ) EMERY, 1908, Deutsch. Ent. Zeitschr., pp. 445 and 447 (2, ♀, ♀).

Aphenogaster barbara var. *semirufa* ERN. ANDRÉ, 1883, 'Spec. Hym. Europ. Algérie,' II, pp. 355 and 368 (♀, ♀).

Type locality: not designated; described from the shores of the CASPIAN SEA, SYRIA, PERSIA, and ABYSSINIA.

1d. Var. *angularis* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 75 (♀).

Type locality: Naivasha, 1900 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

1d. Var. *galla* EMERY, 1908, Deutsch. Ent. Zeitschr., p. 447, footnote. FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 250 (♀). ESCHERICH, 1911, Biol. Centralbl., XXXI, p. 48, footnote. KARAWANAW, 1911, Rev. Russe Ent., XI, p. 3 (♀). EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 3 (♀, ♀, ♂).

Aphenogaster (Messor) *barbarus* EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 112.

Stenamma (Messor) *barbarum* subsp. *caduca* var. *galla* EMERY, 1895, *ibid.*, XXXV, p. 179 (♀).

Messor barbarus subsp. *caducus* var. *galla* EMERY, 1897, *ibid.*, XXXVIII, p. 597.

Stenamma (Messor) *barbarum* var. *galla* MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 5; in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 14 (♀, ♀).

Messor barbarus subsp. *galla* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 335 (♀); 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, p. 92, fig. 2, 1 (♀).

Type locality: Boran Galla, Upper Ganale, ABYSSINIA (V. Bottego).

SENEGAMBIA: Dakar; Thiès (F. Silvestri); Longa. ANGLO-EGYPTIAN SUDAN: Khartum (Karawaiew); White Nile Region (I. Trägårdh). ABYSSINIA: (Ilg); Webi; Ogaden (Bricchetti-Robecchi); Kaka, Schoa (Antinori). SOMALILAND: Milmil (Ruspoli); Mogadiscio; Errer-es-Saghir (Bricchetti-Robecchi); Obbia (Pavesi). ERITREA: Asmara; Keren; Nefasit (F. Silvestri); Bogos (Antinori). GERMAN EAST AFRICA: Mt. Meru; Kibonoto, Mt. Kilimanjaro (Sjöstedt).

1d. Var. *rufulus* FOREL, 1918, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 156.

Messor barbarus subsp. *semirufus* var. *rufa* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 250 (♀). ESCHERICH, 1911, Biol. Centralbl., XXXI, p. 48, footnote. KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 4 (♀). (*nec rufa* Karawaiew, 1909).

Type locality: Nefasit, ERITREA (K. Escherich).

ANGLO-EGYPTIAN SUDAN: Khartum (Karawaiew).

1d. Var. *triempressus* (SANTSCHI).

Messor barbarus subsp. *galla* var. *triempressa* SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, p. 92 (♀).

Type locality: Baguirini, Techeckna, Chari-Chad.

CASAMANCE: (Claveau). ABYSSINIA: Harar; Schoa.

2. *Messor braunsi* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 138 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 413 (♀).

Type locality: Willowmore, CAPE PROVINCE (H. Brauns).

3. *Messor capensis* (MAYR).

Atta capensis MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 743 (♀); 1863, ibid., XIII, p. 396.

Aphenogaster capensis ROGER, 1863, 'Verzeich. Formicid.,' p. 30. MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidæ, p. 87 (♀), Pl. III, fig. 25; 1866, Verh. Zool. Bot. Ges. Wien, XVI, p. 896 (♀, ♂), Pl. XX, fig. 9. EMERY, 1884, Ann. Mus. Civ. Genova, XXI, p. 383 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 101.

Aphenogaster barbara var. *capensis* PÉRINGUEY, 1886, Trans. Ent. Soc. London, Proc., p. xxxvii. WASMANN, 1894, 'Verzeichn. Myrmecoph. Termitoph. Arthrop.,' p. 113. DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, p. 340.

Stenamma (Messor) barbarum subsp. *capense* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 35.

Aphenogaster (Messor) barbara var. *capensis* WASMANN, 1896, Notes Leyden Mus., XVIII, p. 75.

Aphenogaster barbara subsp. *capensis* BINGHAM, 1906, Trans. Ent. Soc. London, Proc., p. xxv (♀, ♂).

Messor barbarus subsp. *capensis* ARNOLD, 1913, Proc. Rhodesia Sc. Assoc., XII, p. 14. FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 242 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 405 (♀, ♀, ♂).

Stenamma (Aphenogaster) capense REICHENSFELDER, 1915, Med. Göteborgs Mus. Zool. Afd., V, p. 16.

Type locality: Cape of Good Hope (Novara Expedition).

CAPE PROVINCE: Cape Town (E. Simon); Kimberley (Dixey and Longstaff); Willowmore (G. Arnold). TRANSVAAL: Makapan; Pretoria (E. Simon). NATAL: Durban (G. Arnold); Wessels Neck (C. B. Cooper).

31. Var. **probus** (FOREL).

Messor barbarus subsp. *capensis* var. *probus* FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 266 (♂). ARNOLD, 1920, Ann. South African Mus., XIV, p. 410 (♀).

Type locality: Bothaville, ORANGE FREE STATE (H. Brauns).

32. Var. **schenki** (FOREL).

Messor barbarus subsp. *capensis* var. *schenki* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 15 (♂).

Messor barbarus subsp. *capensis* var. *schencki* ARNOLD, 1920, Ann. South African Mus., XIV, p. 410 (♂).

Type locality: Bethanien, GERMAN SOUTHWEST AFRICA (Schenek).

33. Var. **tropicorum** (FOREL).

Messor barbarus subsp. *capensis* var. *tropicorum* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 444 (♂).

Type locality: MOSSAMEDES (Baum and Van der Kellen).

3a. Subsp. **decipiens** (FOREL) SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, p. 94.

Stenamma (Messor) barbarum subsp. *capense* var. *decipiens* FOREL, 1905, Ann. Soc. Ent. Belgique, XLIX, p. 177 (♂, ♀).

Messor barbarus subsp. *capensis* var. *decipiens* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 15 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 410 (♀).

Type locality: NATAL (R. C. Wroughton).

CAPE PROVINCE: Cape Flats near Cape Town (L. Schultze).

3b. Subsp. **pseudogegyptiacus** (EMERY).

Aphenogaster pseudogegyptiaca EMERY, 1884, Ann. Mus. Civ. Genova, XXI, p. 384 (♂).

Aphenogaster (Messor) barbarus subsp. *capensis* var. *pseudo-egyptiaca* EMERY, 1891, 'Explor. Tunisie, Fourmis,' p. 12.

Stenamma (Messor) barbarum subsp. *capense* var. *pseudo-egyptiacum* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 35.

Messor capensis subsp. *pseudo-egyptiaca* SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, p. 94.

Messor barbarus subsp. *capensis* var. *pseudogegyptiaca* ARNOLD, 1920, Ann. South African Mus., XIV, p. 408 (♂, ♀, ♂), Pl. v, figs. 59 and 59a-b.

Type locality: Cape of Good Hope (L. Péringuey).

TRANSVAAL: Makapan (E. Simon). RHODESIA: Bulawayo (G. Arnold).

4. **Messor cephalotes** EMERY, 1908, Deutsch. Ent. Zeitschr., p. 443.

Stenamma (Messor) barbarum subsp. *cephalotes* EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 179 (♂); 1897, *ibid.*, XXXVIII, p. 597 (♂). FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 61 (♂).

Stenamma (Messor) barbarum var. *cephalotes* MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 14.

Type locality: ARUSSI GALLA, Ganale Gudda, ABYSSINIA (V. Bottego).

ABYSSINIA: Giari Bule (Ruspoli). BRITISH EAST AFRICA: Leitokitok, Ngare Rongai (Sjöstedt).

- 4a. Subsp. *plinii* SANTSCHI, 1917, Bull. Soc. Hist. Nat. Afr. Nord, VIII, p. 94.
Messor plinii SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 165 (♀); 1914,
 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 75 (♀).

Type locality: Nakuru, Rift Valley, 1820 m., BRITISH EAST AFRICA (Ch. Alluaud).
 BRITISH EAST AFRICA: Tchania-Kamiti, Kikuyu (Alluaud and Jeannel).

5. *Messor denticornis* FOREL, in Schultze, 1910, 'Forschungsreise in Süd-
 afrika,' IV, p. 14 (♀, ♀, ♂), Pl. 1, figs. 1 and 3. ARNOLD, 1920, Ann. South African
 Mus., XIV, p. 411 (♀, ♀, ♂).

Type locality: Lüderitz Bay, GERMAN SOUTHWEST AFRICA (L. Schultze).

GERMAN SOUTHWEST AFRICA: Ababis (R. W. Tucker). CAPE PROVINCE:
 Kamaggas; Steinkopf (L. Schultze); Kimberley (G. Arnold).

- 5₁. Var. *brunni* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 444 (♀). ARNOLD,
 1920, Ann. South African Mus., XIV, p. 413 (♀).

Type locality: SOUTHWEST AFRICA (Brunn).

CAPE PROVINCE: Steekstown (Wartmann).

- 5₂. Var. *parvidens* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,'
 IV, p. 15 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 413 (♀).

Type locality: Kubub, GERMAN SOUTHWEST AFRICA (L. Schultze).

Pheidole WESTWOOD

Pheidole WESTWOOD, 1840, Ann. Mag. Nat. Hist., VI, p. 87.

Leptomyrma MOTSCHULSKY, 1863, Bull. Soc. Nat. Moscou, XXXVI, p. 17.

Pheidole BINGHAM, 1903, 'Fauna Brit. India, Hym.,' II, p. 220.

Oecophthora HEER, 1852, 'Hausameise Madeiras,' p. 15.

Aphenogaster (part) EMERY. *Atta* (part) SYKES, F. SMITH. *Formica* (part)
 FABRICIUS et auct. *Myrmica* (part) SMITH et auct. *Oecodoma* (part) JERDON, MCG-
 GRIDGE. *Pristomyrmex* MAYR.

Genotype: *Atta providens* Sykes, 1835.

All the Ethiopian species belong to the subgenus *Pheidole*, *sensu stricto*.

1. *Pheidole æberlii* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 91 (♀,
 ♀, ♂); 1904, Ann. Soc. Ent. Belgique, XLVIII, p. 172.

Pheidole æberlei MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 7 (♀).
 SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 337 (♀). EMERY, 1916,
 Rev. Zool. Afr., IV, p. 250.

Type locality: Upper Senegal, SENEGAMBIA.

SENEGAMBIA: Thiès (F. Silvestri). ANGLO-EGYPTIAN SUDAN: Ondurman (I.
 Trägårdh).

- 1a. Subsp. *erythræa* EMERY, 1916, Rev. Zool. Afr., IV, p. 250.

Pheidole æberlii var. *erythræa* EMERY, 1901, Bull. Soc. Ent. Italiana, XXXIII,
 p. 61 (♀, ♀).

Type locality: Massaua, ERITREA (Belli).

ERITREA: Ghinda (Belli).

2. *Pheidole akermani* ARNOLD, 1920, Ann. South African Mus., XIV, pp.
 419 and 471 (♀, ♀).

Type locality: Pietermaritzburg, NATAL (C. Akerman).

3. *Pheidole areniphila* FOREL, in Schultze, 1910, 'Forschungsreise in Süd-
 afrika,' IV, p. 30; 1910, Ann. Soc. Ent. Belgique, LIV, p. 14. ARNOLD, 1920, Ann.
 South African Mus., XIV, p. 479 (♀, ♀, ♂).

Pheidole arenicola FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 11. (♂, ♀, ♂).

Type locality: Khakhea, BECHUANALAND (L. Schultze).

4. *Pheidole arnoldi* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 130 (♂, ♀); 1913, Deutsch. Ent. Zeitschr., Beih., p. 213 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 418 and 445, fig. 25 (♂, ♀, ♂).

Type locality: Bulawayo, RHODESIA (G. Arnold).

SOUTHERN RHODESIA: common (G. Arnold).

4. Var. *ballaensis* ARNOLD, 1920, Ann. South African Mus., XIV, p. 448 (♂, ♀).

Type locality: Balla-Balla, SOUTHERN RHODESIA (G. Arnold).

4. Var. *rufescens* ARNOLD, 1920, Ann. South African Mus., XIV, p. 449 (♂, ♀).

Type locality: Sipapoma, SOUTHERN RHODESIA (G. Arnold).

SOUTHERN RHODESIA: Victoria Falls (G. Arnold).

5. *Pheidole aspera* MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 746 (♂, ♀); 1863, *ibid.*, XIII, p. 440. ROGER, 1863, 'Verzeich. Formicid.,' p. 31. MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidæ, p. 97 (♂, ♀, ♂), Pl. III, fig. 26. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 88. ARNOLD, 1920, Ann. South African Mus., XIV, p. 477 (♂, ♀).

Type locality: Cape of Good Hope (Novara Expedition).

6. *Pheidole aurivillii* MAYR, 1896, Ent. Tidskr., XVII, p. 238 (♂, ♀). SJÖSTEDT, 1904, 'I Västafrikas Urskogar,' p. 29. SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 369 (♂, ♀, ♂); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 337 (♂, ♀).

Type locality: CAMEROON (Sjöstedt).

FRENCH GUINEA: Kakoulima (F. Silvestri). FRENCH CONGO: Brazzaville (A. Weiss).

6. Var. *attenuata* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 370 (♂). See p. 129.

Type locality: M'Piaka, FRENCH CONGO (A. Weiss).

BELGIAN CONGO: Medje; Bafwabaka (Lang and Chapin); Walikale to Lubutu (J. Bequaert).

6. Var. *rubricalva* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 337 (♂, ♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

6a. Subsp. *kasalensis* FOREL, 1911, Rev. Zool. Afr., I, p. 279 (♂, ♀); 1913, *ibid.*, II, p. 353 (♂, ♀). LAMBORN, 1914, Trans. Ent. Soc. London, (1913), p. 443 (♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 415 (♂, ♀, ♂).

Type locality: Kondué, BELGIAN CONGO (Luja).

BELGIAN CONGO: St. Gabriel (Kohl). SOUTHERN NIGERIA: Lagos (Lamborn).

6a. Var. *amarilæa* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 336 (♂, ♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

7. *Pheidole batrachorum* WM. M. WHEELER. See p. 128 (♂, ♀).

Type locality: Akenge, BELGIAN CONGO (Lang and Chapin).

8. *Pheidole bequaerti* FOREL, 1913, Rev. Zool. Afr., II, p. 330 (♂, ♀). BEQUAERT, *ibid.*, p. 427.

Type locality: Lake Kabwe, BELGIAN CONGO (J. Bequaert).

9. *Pheidole buchholzi* MAYR, 1900, Ent. Tidskr., XXI, p. 276 (♂, ♀). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 436 (♀). SANTSCI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 336 (♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 413 (♂, ♀).

Type locality: Mungo River, CAMEROON (R. Buchholz).

CAMEROON: (L. v. Muralt); Victoria (F. Silvestri). BELGIAN CONGO: Makanga (Kohl).

10. *Pheidole caffra* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 33 (♂). SANTSCI, 1910, *ibid.*, LXXVIII, (1909), p. 361 (♂). ARNOLD, 1920, Ann. South African Mus., XIV, p. 478 (♂).

Type locality: Hamman's Kraal, TRANSVAAL (E. Simon).

FRENCH CONGO: Brazzaville (A. Weiss).

10i. Var. *amena* FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 365 (♂).

Type locality: ERITREA.

10a. Subsp. *abyssinica* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 254 (♂, ♀). REICHENSPERGER, 1913, *ibid.*, XXXV, p. 193.

Pheidole caffra subsp. *abyssinica* WASMANN, 1911, Tijdschr. v. Ent., LIV, p. 199; 1918, *ibid.*, LX, (1917), p. 395.

Type locality: Ghinda, ERITREA (K. Escherich).

ABYSSINIA: Bisa Timu near Harar; Harar (Kristensen). ERITREA: Nefasit (K. Escherich).

10b. Subsp. *bayeri* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 413 (♂, ♀).

Type locality: Kasindi, BELGIAN CONGO (Bayer).

10b. Var. *thysvillensis* Wm. M. WHEELER. See p. 150 (♂, ♀).

Type locality: Thysville, BELGIAN CONGO (Lang and Chapin; J. Bequaert).

10c. Subsp. *senilifrons* Wm. M. WHEELER. See p. 130 (♂, ♀).

Type locality: Yakuluku, BELGIAN CONGO (Lang and Chapin).

11. *Pheidole capensis* MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 746 (♂, ♀); 1863, *ibid.*, XIII, p. 440. ROGER, 1863, 'Verzeich. Formicid.,' p. 31. MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidae, p. 100 (♂, ♀), Pl. iv, fig. 29. PÉRINGUEY, 1886, Trans. Ent. Soc. London, Proc., p. xxxvi. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 89. WASMANN, 1894, 'Verzeich. Myrmecoph. Termitoph. Arthrop.,' pp. 115 and 118. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 134 (♂, ♀). WASMANN, 1896, Notes Leyden Mus., XVIII, p. 76. FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 130 (♂, ♀, ♀). REICHENSPERGER, 1913, Zool. Jahrb. Abt. Syst., XXXV, pp. 203 and 204; 1915, Med. Göteborgs Mus. Zool. Afd., V, p. 42. EMERY, 1916, Rev. Zool. Afr., IV, p. 246, figs. 9a-b (♂, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 419 and 441, fig. 23 (♂, ♀, ♀).

Pheidole megacephala subsp. *capensis* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 34. WASMANN, 1911, Tijdschr. v. Ent., LIV, p. 199. REICHENSPERGER, 1915, Med. Göteborgs Mus. Zool. Afd., V, p. 72.

Type locality: Cape of Good Hope (Novara Expedition).

CAPE PROVINCE: Willowmore (H. Brauns); Cape Town; Kimberley (E. Simon).

11i. Var. *dregei* EMERY, 1916, Rev. Zool. Afr., IV, p. 246, fig. 9a (♂). ARNOLD, 1920, Ann. South African Mus., XIV, p. 443 (♂).

Pheidole megacephala subsp. *dregei* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 34 (♀).

Type locality: Kimberley, CAPE PROVINCE (E. Simon).

TRANSVAAL: Makapan (E. Simon). ORANGE FREE STATE: Bloemfontein (E. Simon). PORTUGUESE EAST AFRICA: Delagoa Bay.

11. Var. *modestior* SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 337 (♀, ♂).

Type locality: Majuba Neck, CAPE PROVINCE (G. Arnold).

NATAL.

11a. Subsp. *redenburgensis* (FOREL) EMERY, 1916, Rev. Zool. Afr., IV, p. 246, fig. 9c (♀).

Pheidole (*Allophidole*) *cutensis* subsp. *redenburgensis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 135 (♀, ♀, ♀, ♂).

Pheidole cutensis subsp. *redenburgensis* FOREL, 1917, Bull. Soc. Vaudoise Sc. Nat., (5) LI, p. 721.

Pheidole (*Allophidole*) *cutensis* subsp. *reddersburgensis* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 418 and 450, fig. 26 (♀, ♀, ♀, ♂).

Type locality: Reddersburg, ORANGE FREE STATE (H. Brauns).

12. *Pheidole clavata* EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 114. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 89. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 134.

Aphenogaster clavata EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 372 (♀).

Type locality: Keren, Bogos, ERITREA (Beccari).

13. *Pheidole concinna* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXXVIII, (1909), p. 362 (♀, ♀, ♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

14. *Pheidole crassinoda* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 32 (♀, ♀). FOREL, 1894, Mitth. Schweiz. Ent. Ges., VIII, p. 94 (without description). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 134; in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 12 (♀, ♀, ♀, ♂). FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 253 (♀, ♀, ♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 9 (♀, ♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 77 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 420 and 472 (♀, ♀).

Type locality: Makapan, TRANSVAAL (E. Simon).

GERMAN SOUTHWEST AFRICA: Damaraland (Ganz); Okahandja (Casper); Herero. BRICHUANALAND: Lehututu (L. Schultze). NATAL: Verulam (Weitzacker). DELAGOA: (Liengme). GERMAN EAST AFRICA: Tanga (Alluaud and Jeannel); Kibonoto, Mt. Kilimanjaro; Ngare-na-Nyuki, Mt. Meru (Sjöstedt). WEST ABYSSINIA: (Ilg).

14i. Var. *pluto* ARNOLD, 1920, Ann. South African Mus., XIV, p. 473 (♀, ♀).

Type locality: Amatongas Forest, PORTUGUESE EAST AFRICA (G. Arnold).

14a. Subsp. *ruspolii* EMERY, 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 597 (♀, ♀). FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 138 (♀, ♂). STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exped. (1907-08),' III, p. 385 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 77 (♀, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 474 (♀, ♀, ♀, ♂), Pl. vi, figs. 63-66.

Pheidole crassinoda var. *ruspolii* FOREL, 1904, Ann. Mus. Zool. Acad. Sc. St. Pétersbourg, VIII, (1903), p. 373 (2).

Type locality: Giari Bule, ABYSSINIA (Ruspoli).

SOMALILAND: Arigalgala (Ruspoli). ABYSSINIA: Ogaden (Ruspoli); KARSSA (de Rothschild). ANGLO-EGYPTIAN SUDAN: Godo Burka (Dmitriev). BELGIAN CONGO: Kassenje, Lake Albert (Schubotz). BRITISH EAST AFRICA: Mbuyuni, Pori, 1100 m. (Alluaud and Jeannel). RHODESIA: Bulawayo (G. Arnold).

15. *Pheidole cuitensis* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 437 (2, ♀); 1917, Bull. Soc. Vaudoise Sc. Nat., (5) LI, p. 721. EMERY, 1916, Rev. Zool. Afr., IV, p. 246.

Type locality: Between the Cuito River and the Cubango River, MOSSAMEDES (Baum and Van der Kellen).

16. *Pheidole escherichi* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 254 (2, ♀, ♀).

Type locality: Ghinda, ERITREA (K. Escherich).

17. *Pheidole excellens* MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 744 (2); 1863, *ibid.*, XIII, p. 440. ROGER, 1863, 'Verzeich. Formicid.,' p. 30. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 90. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 134. SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 335 (♀).

Type locality: GOLD COAST.

FRENCH GUINEA: Kindia; Konakry (F. Silvestri). SLAVE COAST. TOGO: Bismarckburg.

17a. Subsp. *rhodesiana* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 131 (2, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 419 and 469 (2, ♀, ♂), Pl. VI, fig. 69.

Type locality: Bulawayo, RHODESIA (G. Arnold).

17b. Subsp. *weissi* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 361 (2); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 335 (2, ♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

NIGERIA: Lagos (F. Silvestri).

18. *Pheidole foreli* MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 8 (2, ♀). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 12 (♀). WASMANN, 1911, Tijdschr. v. Ent., LIV, p. 199. FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 241 (2, ♀). REICHENSPERGER, 1915, Med. Göteborgs Mus. Zool. Afd., V, pp. 34, 45, and 73. EMERY, 1916, Rev. Zool. Afr., IV, p. 245. ARNOLD, 1920, Ann. South African Mus., XIV, pp. 417 and 432 (2, ♀, ♀).

Type locality: Sunday River Mts. (2400 ft.), near Port Elizabeth, CAPE PROVINCE (H. Brauns).

CAPE PROVINCE: Deep River near Cape Town in the Cape Flats (L. Schultze); KNYSNA (H. Brauns). ORANGE FREE STATE: Bothaville; Reddersburg (H. Brauns). NATAL: Estcourt (R. C. Wroughton).

18a. Var. *pubens* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 12 (2, ♀); 1910, Ann. Soc. Ent. Belgique, LIV, p. 436 (♀, ♀, ♂). ARNOLD, 1920, Ann. South African Mus., XIV, p. 434 (2, ♀).

Type locality: NATAL (R. C. Wroughton; Haviland).

19. *Pheidole irritans* (SMITH) DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, pp. 331, 332, and 371.

Myrmica irritans F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 127 (♀).
ROGER, 1863, 'Verzeich. Formicid.,' p. 33. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 433. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 110.

Type locality: Port Natal, NATAL.

CAPE PROVINCE: Nahoon River (Dixey and Longstaff). NATAL: Colenso; Howick (Dixey and Longstaff).

20. *Pheidole jordanica* DE SAULCY, 1874, Bull. Soc. Hist. Nat. Metz, XIII, p. 17 (♂, ♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 91. EMERY, 1916, Rev. Zool. Afric., IV, p. 233, figs. 4a-c (♂, ♀).

Pheidole megacephala subsp. *jordanica* EMERY, 1889, Ann. Mus. Civ. Genova, XXVII, p. 442 (♂, ♀).

Pheidole sinaitica WASMANN, 1894, 'Verzeichn. Myrmecoph. Termitoph. Arthrop.,' p. 119 (nec Mayr).

Pheidole sinaitica subsp. *laticeps* MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 6 (♂, ♀) (not "*latifrons*" as quoted by Emery, 1916).

Pheidole schmitzi FOREL, 1911, Rev. Suisse Zool., XIX, p. 455 (♂, ♀, ♀).

Pheidole sinaitica var. *laticeps* KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 7 (♂, ♀).

Type locality: Jericho, PALESTINE (Piochard de la Brûlerie).

CYRENAICA, EGYPT. ANGLO-EGYPTIAN SUDAN: Khartum; Port Sudan (Karawaiew).

21. *Pheidole kitcheneri* FOREL, emend.

Pheidole kitschneri FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 436 (♂, ♀?). ARNOLD, 1920, Ann. South African Mus., XIV, p. 479 (♂, ♀?).

Pheidole kitcheneri SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicid.,' p. 76 (♂, ♀, ♀).

Type locality: Mountains of NATAL (R. C. Wroughton).

BRITISH EAST AFRICA: Cheteni; Fort Hall, Wa-Kikuyu, 1330 m.; Tehania River, 1520 m. (Alluaud and Jeannel).

22. *Pheidole kohli* MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 11, footnote (♂, ♀). See p. 131.

Type locality: Waboniland, BRITISH EAST AFRICA.

BELGIAN CONGO: Medje; Garamba (Lang and Chapin).

23. *Pheidole liengmei* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 93 (♂); 1913, Ann. Soc. Ent. Belgique, LVII, p. 134 (♂, ♀). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 23 (♀). REICHENSPERGER, 1915, *ibid.*, V, p. 18. LAMBORN, 1920, Trans. Ent. Soc. London, (1919), Proc., pp. liii and lv. ARNOLD, 1920, Ann. South African Mus., XIV, pp. 419 and 465, fig. 30 (♂, ♀, ♀), Pl. VI, fig. 71.

Type locality: Delagor, PORTUGUESE EAST AFRICA (Liengme).

ZULULAND: Mkosi (I. Trägårdh). RHODESIA: Bulawayo; Rhodesia Falls (G. Arnold). GERMAN EAST AFRICA: Lindi (Lamborn).

23i. Var. *malindana* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 135 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 468 (♀).

Type locality: Malindi, SOUTHERN RHODESIA (G. Arnold).

23. Var. *shinsendensis* FOREL, 1913, Rev. Zool. Afr., II, p. 327 (♀).

Pheidole liengmei var. *shinsendensis* BEQUAERT, 1913, *ibid.*, II, p. 426.

Pheidole liengmei var. *shinsendensis* ARNOLD, 1920, Ann. South African Mus., XIV, p. 468 (♂, ♀).

- Type locality: Shinsenda, BELGIAN CONGO (J. Bequaert).
 RHODESIA: Bembesi (G. Arnold).
 23a. Subsp. *micrartifex* FOREL, 1913, *Deutsch. Ent. Zeitschr.*, Beih., p. 215 (2, ♀). ARNOLD, 1920, *Ann. South African Mus.*, XIV, p. 468 (2, ♀).
 Type locality: Bulawayo, RHODESIA (G. Arnold).
 24. *Pheidole maufei* ARNOLD, 1920, *Ann. South African Mus.*, XIV, pp. 418 and 438, fig. 22a (2, ♀).
 Type locality: Belingwe, SOUTHERN RHODESIA (G. Arnold).
 25. *Pheidole mayri* FOREL, 1894, *Mitth. Schweiz. Ent. Ges.*, IX, p. 91 (2, ♀).
 Type locality: Senegal.
 26. *Pheidole megacephala* (FABRICIUS) ROGER, 1863, 'Verzeich. Formicid.,' pp. 30 and 49. MAYR, 1863, *Verh. Zool. Bot. Ges. Wien*, XIII, p. 441. MORRIS, 1880, *American Naturalist*, XIV, pp. 667-670. FOREL, 1881, *Mitth. München. Ent. Ver.*, V, p. 8 (♀). LUBBOCK, 1882, 'Ants, Bees and Wasps,' 5th Ed., pp. 20, 180, and 233 (♀), Pl. II, figs. 3 and 4. ERN. ANDRÉ, 1883, 'Spec. Hym. Europ. Algérie,' II, pp. 383, 385, and 386 (2, ♀, ♀, ♂). PÉRINGUEY, 1886, *Trans. Ent. Soc. London, Proc.*, p. xxxvi. BLACKBURN and CAMERON, 1886, *Mem. Manchester Lit. Phil. Soc.*, (3) X, p. 236. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 92. MAYR, 1893, *Jahrb. Hamburg. Wiss. Anst.*, X, 2, p. 201. WASMANN, 1894, 'Verzeichn. Myrmecoph. Termitoph. Arthrop.,' pp. 115, 116, 118, and 119. MAYR, 1896, *Ent. Tidskr.*, XVII, p. 238. WASMANN, 1896, *Notes Leyden Mus.*, XVIII, p. 76. EMERY, 1897, *Ann. Mus. Civ. Genova*, XXXVIII, p. 598 (♀). H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 40. WASMANN, 1899, *Deutsch. Ent. Zeitschr.*, p. 406. MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 7. WASMANN, 1904, *Notes Leyden Mus.*, XXV, p. 45. MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 13. DIXEY and LONGSTAFF, 1907, *Trans. Ent. Soc. London*, pp. 332 and 336. SANTSCHI, 1910, *Ann. Soc. Ent. France*, LXXVIII, (1909), p. 370. WASMANN, 1911, *Tijdschr. v. Ent.*, LIV, p. 198. REICHENSPERGER, 1913, *Zool. Jahrb. Abt. Syst.*, XXXV, pp. 203, 212, and 215. SANTSCHI, 1914, *Boll. Lab. Zool. Gen. Agrar. Portici*, VIII, p. 336 (♀, ♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 75 (2, ♀, ♀); 1914, *Med. Göteborgs Mus. Zool. Afd.*, III, p. 22 (2, ♀, ♀). TRÄGÅRDH, *ibid.*, p. 44. ALLUAUD and JEANNEL, 1914, *Arch. Zool. Gén. Exp.*, LIII, p. 366. EMERY, 1916, *Rev. Zool. Afr.*, IV, pp. 235 and 238. WASMANN, 1918, *Tijdschr. v. Ent.*, LX, (1917), p. 395. SANTSCHI, 1920, 'Études Maladies Parasites Cuckooer S. Thomé,' X, p. 2 (2, ♀, ♀). See pp. 131 and 1018.
?Formica edax FORSKÅL, 1775, 'Descr. Anim.,' p. 84.
Formica megacephala FABRICIUS, 1793, 'Ent. Syst.,' II, p. 361. COQUEBERT, 1799, 'Illustr. Iconogr. Ins.,' I, p. 26 (♀, ♀), Pl. vi, fig. 9. LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 232 (♀), Pl. x, fig. 67. FABRICIUS, 1804, 'Syst. Piez.,' p. 411.
Formica (Myrmica) trinodis LOSANA, 1834, *Mem. Accad. Sc. Torino*, XXXVII, p. 327, Pl. xxxvi, fig. 6.
Oecophthor pusilla HEER, 1852, 'Hausameise Madeiras,' p. 15 (2, ♀, ♀, ♂), Pl. I, figs. 1-4. BROWN, 1869, *Proc. Boston Soc. Nat. Hist.*, XII, p. 211.
Myrmica? levigata SMITH, 1855, *Trans. Ent. Soc. London*, (2) III, p. 130 (♀), Pl. ix, figs. 7 and 8.
Myrmica trinodis MAYR, 1855, *Verh. Zool. Bot. Ges. Wien*, V, p. 414, footnote (♀).

Myrmica (*Pheidole*) *lavigata* F. SMITH, 1858, 'Cat. Brit. Fossor. Hym.,' pp. 35 and 225 (♀).

Pheidole pusilla F. SMITH, *ibid.*, p. 173, Pl. IX, figs. 18-20. MAYR, 1861, 'Europ. Formicid.,' p. 70 (2, ♀, ♀); 1870, Verh. Zool. Bot. Ges. Wien, XX, pp. 981 and 984 (2, ♀); in Fedtschenko, 1877, 'Voy. Turkestan, Formicid.,' p. 18; 1878, Verh. Zool. Bot. Ges. Wien, XXVIII, pp. 674 and 676 (2, ♀). BLACKBURN and KIRBY, 1880, Ent. Monthly Mag., XVII, p. 89. MAYR, 1887, Verh. Zool. Bot. Ges. Wien, XXXVII, pp. 596 and 606 (2, ♀).

Pheidole janus F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 175 (♀), Pl. IX, figs. 13-17. MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 360.

?*Ecophthora perniciosus* GERSTAECKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 263 (♀); in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 516 (♀).

Myrmica lavigata F. SMITH, 1859, Zoologist, XVII, p. 6385; 1862, Ent. Annual, p. 70 (♀, ♀), Pl., figs. 4, 7, 8.

Pheidole lavigata MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 747 (2). WHITE, 1883, 'Ants and Their Ways,' p. 270 (♀).

?*Pheidole perniciosus* MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 441. ROGER, 1863, 'Verzeich. Formicid.,' p. 31. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 94.

Pheidole megacephala subsp. *pusilla* EMERY, 1916, Rev. Zool. Afr., IV, pp. 235 and 239, fig. 1a (2); 1919, *ibid.*, VI, p. 170, fig. 5a (2).

Pheidole megacephala var. *pusilla* SANTSCHI, 1920, 'Études Maladies Parasites Cacaoyer S. Thomé,' X, p. 2 (2, ♀).

Type locality: MAURITIUS (= Isle de France).

Cosmopolitan in tropical and subtropical regions.

SENEGAMBIA: Dakar (F. Silvestri). CAMEROON: Sjöstedt). SAN THOMÉ: (de Seabra). FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: (Kohl); Malela; Matadi; Thysville; Boma (J. Bequaert); Zambi; Banana; Niangara; Akenge; Stanleyville (Lang and Chapin). ANGOLA: St. Paul de Loanda (F. Silvestri). CAPE PROVINCE: Cape Town (Raffray). NATAL: Stamford Hill (I. Trägårdh); Colenso (Dixey and Longstaff). ZULULAND: Junction of the Umfolosi Rivers (I. Trägårdh). TRANSVAAL: Johannesburg (Dixey and Longstaff). PORTUGUESE EAST AFRICA: Tete (Peters). GERMAN EAST AFRICA: Tanga; New Moschi, Mt. Kilimanjaro, 800 m. (Alluaud and Jeannel); Kibonoto, Kilimanjaro; Usambara (Sjöstedt); Bagamoyo; Kihengo (Stuhlmann). ZANZIBAR: (Stuhlmann). BRITISH EAST AFRICA: Shimoni; Ramisi River; Naivasha, Rift Valley, 1900 m.; Nairobi; Port Florence, Kavirondo Bay (Alluaud and Jeannel). ABYSSINIA: Ogaden (Ruspoli); Harar (Kristensen). ANGLO-EGYPTIAN SUDAN: Abba Island, White Nile (I. Trägårdh).

25a. Subsp. *ilgi* (FOREL) EMERY, 1916, Rev. Zool. Afr., IV, pp. 236 and 244. See p. 132.

Pheidole rotundata subsp. *ilgi* FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 82 (2, ♀); 1907, Rev. d'Ent. Caen, XXVI, p. 139 (2, ♀, ♀, ♂); 1913, Ann. Soc. Ent. Belgique, LVII, p. 128 (2, ♀). REICHENSPERGER, 1913, Zool. Jahrb. Abt. Syst., XXXV, pp. 191 and 195; 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, p. 123. H. SCHMITZ, 1916, Zoolog. Meded. Mus. Leiden, II, p. 28, footnote. WASMANN, 1918, Tijdschr. v. Ent., LX, (1917), p. 395.

Pheidole rotundata subsp. *impressifrons* var. *ilgi* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 417 and 427 (2, ♀, ♀, ♂).

Type locality: Harar, ABYSSINIA (Ilg; Kristensen).

ABYSSINIA: Hiéka-Bourka; Karssa; Tehafianani; Kounhi; Adis-Abeba (de Rothschild); Diré Daus (Kristensen). RHODESIA: Bulawayo (G. Arnold). BRITISH EAST AFRICA: Fundu Island near Pemba (Voeltzkow). BELGIAN CONGO: Lesse (J. Bequaert).

26b. Subsp. *impressifrons* WASMANN, 1905, Notes Leyden Mus., XXV, p. 110; 1911, Tijdschr. v. Ent., LIV, p. 199. EMERY, 1916, Rev. Zool. Afr., IV, pp. 236 and 243, figs. 7b and c (♂, ♀).

Pheidole rotundata subsp. *impressifrons* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 213 (♂, ♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 337 (♂, ♀). FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., 55, L, p. 239 (♂, ♀); 1916, Rev. Suisse Zool., XXIV, p. 416 (♂, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 426 (♂, ♀, ♀).

Pheidole megacephala subsp. *impressiceps* WASMANN, 1904, Notes Leyden Mus., XXV, pp. 38, 41, 46, and 72, footnote (♂, ♀, ♀) (nec Mayr).

Pheidole punctulata subsp. *impressifrons* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 127 (♂, ♀, ♀); 1913, Rev. Zool. Afr., II, pp. 328 and 352 (♂). J. BEQUAERT, 1913, *ibid.*, II, p. 427. REICHENSBERGER, 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, p. 123. WASMANN, 1918, Tijdschr. v. Ent., LX, (1917), p. 396.

Pheidole rotundata var. *impressifrons* LAMBORN, 1914, Trans. Ent. Soc. London, (1913), p. 443.

Pheidole punctatissima subsp. *impressifrons* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 22 (♂, ♀). TRÄGÅRDH, *ibid.*, p. 45. REICHENSBERGER, 1915, *ibid.*, V, pp. 10, 14, and 18.

Pheidole megacephala subsp. *punctulata* var. *impressiceps* EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 3 (♂, ♀).

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauds).

SOUTHERN NIGERIA: Lagos (Lamborn). BELGIAN CONGO: (Kohl); Welgelegen (J. Bequaert). RHODESIA: Bulawayo (G. Arnold). TRANSVAAL: Pretoria (F. Silvestri). CAPE PROVINCE: Willowmore (H. Brauds). NATAL: Mountains at 4000 ft. (Haviland). ZULULAND: Junction of Umfolozi (I. Trägårdh). ERITREA: Asmara (F. Silvestri). ABYSSINIA: Harar (Kristensen).

26b. Var. *atrociior* SANTSCHI *in litt.* EMERY, 1916, Rev. Zool. Afr., IV, pp. 236 and 243, fig. 7d (♂).

Type locality: SIERRA LEONE.

26c. Subsp. *melancholica* (SANTSCHI) WASMANN, 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, p. 280. EMERY, 1916, Rev. Zool. Afr., IV, pp. 236 and 242, fig. 7a (♂, ♀). See p. 132.

Pheidole punctulata subsp. *melancholica* SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 164 (♂, ♀).

Pheidole punctulata var. *melancholica* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 337 (♂, ♀, and mermithergate).

Type locality: Jacqueville, IVORY COAST (Lohier).

GOLD COAST: Aburi (F. Silvestri). BELGIAN CONGO: Garamba (Lang and Chapin); St. Gabriel (Kohl).

26c. Var. *angulata* (STITZ) EMERY, 1916, Rev. Zool. Afr., IV, pp. 236 and 243 (♂).

Pheidole megacephala subsp. *punctulata* var. *angulata* STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08)', III, p. 385 (♂, ♀).

Type locality: Lake Mohasi, east of Lake Kivu, GERMAN EAST AFRICA (Schubotz).

26c. Var. *costauriensis* (SANTSCHI) EMERY, 1916, Rev. Zool. Afr., IV, pp. 236 and 243, fig. 7a (♂).

Pheidole rotundata subsp. *costauriensis* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 433 (♂); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 337 (♂, ♀); 1915, Ann. Soc. Ent. France, LXXXIV, p. 250 (♀).

Type locality: Winnebah, GOLD COAST.

FRENCH GUINEA: Kindia; Kakoulima (F. Silvestri). SOUTHERN NIGERIA: Ibadan (F. Silvestri).

26d. Subsp. *nkomoana* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 415 (♂, ♀, ♂).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

26e. Subsp. *punctulata* (MAYR) FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 94. WASMANN, 1894, 'Verzeichn. Myrmecoph. Termitoph. Arthrop.', p. 117. FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 311. EMERY, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 62 (♂). FOREL, in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 564. WASMANN, 1911, Tijdschr. v. Ent., LIV, p. 199. EMERY, 1916, Rev. Zool. Afr., IV, pp. 235 and 241, figs. 6a and 6c (♂, ♀). See pp. 132 and 1019.

Pheidole punctulata MAYR, 1866, Verh. Zool. Bot. Ges. Wien, XVI, p. 899 (♂). EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 270; 1892, Ann. Soc. Ent. France, LX, (1891), p. 563. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 95. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 4 (♂). WASMANN, 1896, Notes Leyden Mus., XVIII, p. 76. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 40. WASMANN, 1907, Deutsch. Ent. Zeitschr., p. 149. FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 81; 1907, Mitt. Naturh. Mus. Hamburg, XXIV, p. 15 (♂, ♀). F. C. WELLMAN, 1908, Ent. News, XIX, p. 230. FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, pp. 53, 59, and 65 (♂, ♀); 1910, *ibid.*, LIV, p. 441; 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 362 (♂, ♀, ♀); 1912, Mitt. Naturh. Mus. Hamburg, XXIX, p. 180; 1913, Rev. Zool. Afr., II, p. 328 (♂, ♀). BEQUAERT, *ibid.*, p. 426. FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 353 (♂, ♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 337 (♂, ♀). FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 240 (♂, ♀, ♀, ♂). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicides,' p. 76 (♂, ♀). REICHENSPERGER, 1915, Med. Göteborgs Mus. Zool. Afd., V, p. 20. WASMANN, 1920, Tijdschr. v. Ent., LXII, (1919), p. 128. ARNOLD, 1920, Ann. South African Mus., XIV, pp. 417 and 420 (♂, ♀, ♀, ♂), Pl. vi, fig. 67. J. HEWITT, 1920, South African Journ. Nat. Hist., II, p. 109.

Pheidole talpa GERSTAECKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 356 (♂); 1873, 'in v. d. Decken's Reisen in Ostafrika, Gliederthiere,' p. 360 (♂), Pl. XIV, fig. 11.

Pheidole megacephala var. *punctulata* WASMANN, 1898, Wien. Ent. Zeitg., XVII, p. 103; 1899, Notes Leyden Mus., XXI, p. 50. MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 14. REICHENSPERGER, 1915, 'Med. Göteborgs Mus. Zool. Afd., V, pp. 37, 38, 41, 46, and 53.

Type locality: Caffraria.

FRENCH GUINEA: Mamou (Silvestri). IVORY COAST: Assinie (C. Alluaud). FRENCH CONGO: Ogowe (Mocquety). PORTUGUESE CONGO: Conde; Landana (Petit); Shiloango (Tschoffen). BELGIAN CONGO: Lower Congo (Solon); Kasai,

Kondué (Luja); Leopoldville; Elisabethville; Kasenga; Bukama; Lukonzolwa (J. Bequaert); Banana (Busschodts); Kinshasa (Waelbroeck); Banana to Boma (Tschoffen); Kisantu (Goossens); Mayombe (de Bréy; Luja); Kiniati; Congo da Lemba (Luja); Boma; Zambi; Banana; Bolobo; Stanleyville; Ngayu; Avakubi; Niapu; Faradje; Caramba (Lang and Chapin); Tua (J. Maes). ANGOLA: St. Paul de Loanda (F. Silvestri). BENGUELA: (C. Wellman). MOSSAMEDES: between the Cubango and Cuito Rivers (Baum and Van der Kellen). RHODESIA: Bulawayo (G. Arnold). TRANSVAAL: Pretoria. CAPE PROVINCE: Lower Albany; Grahamstown (Hewitt); Port Elizabeth (H. Brauns). NATAL: Durban (G. Arnold and L. Bevis). PORTUGUESE EAST AFRICA: Delagoa Bay (Liengme); Mozambique. ZANZIBAR: (Ville). GERMAN EAST AFRICA: Kibonoto, Mt. Kilimanjaro (Sjöstedt); Marangu, Mt. Kilimanjaro, 1800 m. (Alluaud and Jeannel); Amani (H. Prell); Mbaramu (v. d. Decken); Moschi (Zimmer). BRITISH EAST AFRICA: Chake Chake, Pemba Island (Voeltzkow); Nairobi, 1660 m. (Alluaud and Jeannel). UGANDA: Entebbe (Schultze). ABYSSINIA: Let Marefa, Schoa (Antinori). ERITREA: Ghinda (Belli).

26e. Var. **atrox** (FOREL) EMERY, 1916, Rev. Zool. Afr., IV, pp. 236 and 241, figs. 7b and d (2).

Pheidole punctulata subsp. *atrox* FOREL, 1913, *ibid.*, II, p. 328 (2, ♀, ♀). BEQUAERT, *ibid.*, p. 426. FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 126 (2, ♀, ♀). WASMANN, 1918, Tijdschr. v. Ent., LX, (1917), p. 395.

Pheidole punctatissima subsp. *atrox* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afld., III, p. 22 (2, ♀, ♀). TRÄGÄRDH, *ibid.*, p. 44. REICHENSPERGER, 1915, *ibid.*, V, p. 18.

Pheidole punctulata var. *atrox* ARNOLD, 1920, Ann. South African Mus., XIV, p. 424 (2, ♀, ♀, ♂).

Pheidole inquilina? FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 240 (mermithergate!). ARNOLD, 1920, Ann. South African Mus., XIV, 425 (recognizes its true nature as a mermithergate and found it in the nests of *P. megacephala punctulata* var. *atrox*).

Type locality: Elisabethville, BELGIAN CONGO (J. Bequaert).

BELGIAN CONGO: Bukama (J. Bequaert). RHODESIA: Bulawayo (G. Arnold). ZULULAND: Junction of the Umfolosi Rivers (I. Trägårdh). NATAL: Durban, Amanzimtoti (I. Trägårdh).

26e. Var. **speculifrons** (STITZ) EMERY, 1916, Rev. Zool. Afr., IV, pp. 236 and 242 (2).

Pheidole megacephala var. *speculifrons* STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08)', III, p. 386 (2, ♀) (*nec Pheidole speculifrons* Dalla Torre, 1893).

Type locality: Bukoba, Victoria Nyanza, GERMAN EAST AFRICA (Schubotz).

26f. Subsp. **rotundata** (FOREL) EMERY, 1916, Rev. Zool. Afr., VI, pp. 236 and 244 (2, ♀).

Pheidole rotundata FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 92 (2, ♀). EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 34. FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 311; in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 564; 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 239 (2, ♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 337 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 76 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 417 and 425, fig. 21 (2, ♀, ♂).

Type locality: DELAGOA (Liengme).

MOSSAMEDES: between the Cubango and Cuito Rivers (Baum and Van der Kellen). TRANSVAAL: Pretoria (E. Simon; F. Silvestri). NATAL: Durban (G. Arnold). RHODESIA: Livingstone; Chirinda Forest (G. Arnold). GERMAN EAST AFRICA: New Moschi, Mt. Kilimanjaro, 800 m. (Alluaud and Jeannel). BRITISH EAST AFRICA: Nairobi, Kikuyu (Alluaud).

27. *Pheidole mentita* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 338 (♂, ♀).

Type locality: Kindia, FRENCH GUINEA (F. Silvestri).

27i. Var. *pullata* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 339 (♂, ♀).

Type locality: Camayenne, FRENCH GUINEA (F. Silvestri).

28. *Pheidole minima* MAYR, 1900, Ent. Tidskr., XXI, p. 275 (♂, ♀).

Type locality: Mungo River, CAMEROON (R. Buchholz).

28i. Var. *catella* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 339 (♂, ♀).

Type locality: Olokemeji, SOUTHERN NIGERIA (F. Silvestri).

GOLD COAST: Aburi (F. Silvestri).

28a. Subsp. *corticicola* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 366 (♂, ♀, ♂, ♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

FRENCH CONGO: Mindouli; Gomba (A. Weiss).

28b. Subsp. *faurei* SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 15 (♂, ♀).

Type locality: Samkita, FRENCH CONGO (F. Faure).

28c. Subsp. *malelana* Wm. M. WHEELER. See p. 133 (♂, ♀).

Type locality: Malela, BELGIAN CONGO (Lang and Chapin).

29. *Pheidole mylognatha* Wm. M. WHEELER. See p. 134 (♂, ♀).

Type locality: Banana, BELGIAN CONGO (Lang and Chapin).

30. *Pheidole niapuana* Wm. M. WHEELER. See p. 136 (♂, ♀).

Type locality: Niapu, BELGIAN CONGO (Lang and Chapin).

31. *Pheidole nigeriensis* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 340 (♂).

Type locality: Olokemeji, SOUTHERN NIGERIA (F. Silvestri).

32. *Pheidole njassæ* VIEHMAYER, 1914, Arch. f. Naturg., LXXIX, Abt. A, Heft 12, (1913), p. 35 (♂, ♀).

Type locality: Marow, Langenburg, GERMAN EAST AFRICA.

32i. Var. *legitima* (SANTSCHI).

Pheidole njassæ var. *sculptor* VIEHMAYER, 1914, Arch. f. Naturg., LXXIX, Abt. A, Heft 12, (1913), p. 36 (♂, ♀). (nec *P. flavens* var. *sculptor* Forel, 1893).

Pheidole njassæ var. *legitima* SANTSCHI, 1916, Bull. Soc. Ent. France, p. 242.

Type locality: Khutu Steppe, Morogoro, GERMAN EAST AFRICA (K. Schwartz).

33. *Pheidole occipitalis* ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 321 (♂, ♀). DALLA TORRE, 1893, 'Cat. Hym.' VII, p. 93.

Type locality: SIERRA LEONE (Mocquerys).

33a. Subsp. *neutalis* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 336 (♂, ♀).

Type locality: Camayenne, FRENCH GUINEA (F. Silvestri).

34. *Pheidole* ? *pallidelutea* (LATREILLE) ROGER, 1863, 'Verzeich. Formicid.,' p. 30. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 94.

Formica pallidelutea LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 241 (♀). MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 418.

Type locality: Senegal.

35. *Pheidole philippi* EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 8 (2, ♀).

Type locality: Ghinda, ERITREA (F. Silvestri).

36. *Pheidole picata* (FOREL) EMERY, 1916, Rev. Zool. Afr., IV, p. 245, fig. 8b (2, ♀). See p. 1019.

Pheidole punctulata var. *picata* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 370.

FRENCH CONGO: Brazzaville (A. Weiss).

37. *Pheidole prelli* FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 362 (2, ♀, ♀, ♂).

Type locality: Mombasa, BRITISH EAST AFRICA (H. Prell).

37a. Subsp. *redbankensis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 128 (2, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 418 and 434, fig. 22 (2, ♀).

Type locality: Bulawayo, RHODESIA (G. Arnold). This is the locality given by Forel; Arnold (1920) corrects it to Redbank, SOUTHERN RHODESIA.

37a. Var. *politocciptus* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXXIV, (1915), p. 503 (2, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 418 and 436 (2, ♀, ♀, ♂).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

RHODESIA: Bulawayo (G. Arnold).

38. *Pheidole pulchella* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 360 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

39. *Pheidole rugaticeps* EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 375 (2, ♀, ♀). MAGRETTI, 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 245 (♀); 1884, Ann. Mus. Civ. Genova, XXI, p. 542 (♀). EMERY, 1892, *ibid.*, XXXII, p. 112 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 96. FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 253 (2, ♀, ♀). EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 3 (2, ♀).

Type locality: Sciotel, ERITREA (Beccari).

ERITREA: Ghinda (K. Escherich; F. Silvestri). SOMALILAND: Obbia (Bricchetti-Robecchi); Sinadogo; Erdal (Pavesi). ANGLO-EGYPTIAN SUDAN: El Hefera, Settit (Magretti). ARABIA: Yemen.

39a. Subsp. *arabs* (EMERY) FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 253 (2). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 336.

Pheidole rugaticeps var. *arabs* EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 535 (2, ♀, ♀).

Type locality: Tes, southern ARABIA (Doria and Beccari).

ERITREA: Ghinda (K. Escherich). SENEGAMBIA: Dakar; Thiès (F. Silvestri).

40. *Pheidole saxicola* WM. M. WHEELER. See p. 138 (2, ♀).

Type locality: Zambi, BELGIAN CONGO (Lang, Chapin and Bequaert).

BELGIAN CONGO: Boma (J. Bequaert).

41. *Pheidole scabriuscula* GERSTÄCKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 356 (♂); 1873, in 'v. d. Decken's Reisen in Ostafrika, Gliederthiere,' p. 360 (♂).
 c DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 96. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 40.

Type locality: Ndara, BRITISH EAST AFRICA (v. d. Decken).

42. *Pheidole schoutedeni* FOREL, 1913, Rev. Zool. Afr., II, p. 329 (2, ♀).
 BEQUAERT, *ibid.*, p. 427.

Type locality: Welgelegen, BELGIAN CONGO (J. Bequaert).

- 42a. Var. *platycephala* (STITZ) ARNOLD, 1920, Ann. South African Mus., XIV, p. 450.

Pheidole platycephala STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp., 1910-11,' I, p. 380, fig. 3 (2, ♀).

Type locality: Libenge, BELGIAN CONGO (Schubotz).

43. *Pheidole schultzei* FOREL, in Schultze, 1910, 'Forschungsreise in Süd-afrika,' IV, p. 9, fig. 1 (2, ♀, ♀, ♂). ARNOLD, 1920, Ann. South African Mus., XIV, p. 457 (2, ♀, ♀, ♂).

Pheidole schultzei FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 436 (2, ♀).

Type locality: Between Koca and Sekgoma, BECHUANALAND (L. Schultze).

BECHUANALAND: Between Kgokong and Kang (L. Schultze). RHODESIA: Victoria Falls (R. C. Wroughton).

- 43a. Var. *gwaalensis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 129 (2, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 419 and 458, fig. 28 (2, ♀, ♀, ♂).

Type locality: Gwari, RHODESIA (G. Arnold).

44. *Pheidole sculpturata* MAYR, 1866, Verh. Bot. Ges. Wien, XVI, p. 897 (2). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 96. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 31 (2, ♀); 1900, Bull. Soc. Ent. Italiana, XXXII, p. 114, fig. 8 (2). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 361 (2, ♀). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 9 (2); 1910, Ann. Soc. Ent. Belgique, LIV, p. 436 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 419 and 461, fig. 29 (2, ♀).

Type locality: Caffraria.

FRENCH CONGO: Gomba; Mandouga; Mbamu (A. Weiss). GERMAN SOUTH-WEST AFRICA: (Lübbert); Okahandja (Casper). TRANSVAAL: Pretoria (E. Simon; Lounsbury). ORANGE FREE STATE: Bloemfontein (E. Simon); Bothaville (H. Brauns). NATAL: (R. C. Wroughton; Haviland).

- 44a. Var. *areolata* FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 269 (2, ♀).
 ARNOLD, 1920, Ann. South African Mus., XIV, p. 463 (2, ♀).

Type locality: Bothaville, ORANGE FREE STATE (H. Brauns).

- 44a. Subsp. *berthoudi* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 89 (2, ♀); 1910, Ann. Soc. Ent. Belgique, LIII, p. 65 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 361 (2, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 463 (2, ♀).

Type locality: Valdezia, TRANSVAAL (P. Berthoud).

FRENCH CONGO: Gomba; Mandouga; Mbamu (A. Weiss). BENGUELA: (Wellman).

- 44b. Subsp. *dignata* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 250 (2, ♀).

Type locality: Gomba, FRENCH CONGO (A. Weiss).

FRENCH CONGO: Mandouga (A. Weiss).

44c. Subsp. *katonæ* FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 21 (♂).

Type locality: Boma Gombe, GERMAN EAST AFRICA (Katona).

44d. Subsp. *welgelegenensis* FOREL, 1913, Rev. Zool. Afr., II, p. 327 (2, ♀). BEQUAERT, *ibid.*, p. 426. FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 354 (♀).

Type locality: Welgelegen, BELGIAN CONGO (J. Bequaert).

BELGIAN CONGO: Nieuwdorp (Leplae).

44e. Subsp. *zambesiana* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 133 (2, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 464 (2, ♀).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

RHODESIA: Shiloh (G. Arnold).

45. *Pheidole sinaitica* MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 745 (2); 1863, *ibid.*, XIII, p. 441. ROGER, 1863, 'Verzeich. Formicid.,' p. 31. MAYR, 1878, Verh. Zool. Bot. Ges. Wien, XXVIII, p. 675 (2). ERN. ANDRÉ, 1881, Ann. Soc. Ent. France, (6) I, p. 73; 1883, 'Spec. Hym. Europ. Algérie,' II, pp. 383 and 384 (2, ♀). MAGRETTI, 1884, Ann. Mus. Civ. Genova, XXI, p. 542 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 96. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 94. MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicidæ,' p. 6 (2, ♀). FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 253 (2, ♀). EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 3 (2, ♀); 1916, Rev. Zool. Afric., IV, p. 234, figs. 4d and e. *Pheidole synaitica* MAGRETTI, 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 245 (♀).

Type locality: Sinai Peninsula (v. Frauenfeld).

ERITREA: Ghinda (K. Escherich); Nefasit (K. Escherich; F. Silvestri). SOUTHERN ABYSSINIA: (Ilg). ANGLO-EGYPTIAN SUDAN: Athara River; Sebderat (Magretti).

46. *Pheidole speculifera* EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 373 (2, ♀). MAGRETTI, 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 245 (♀); 1884, Ann. Mus. Civ. Genova, XXI, p. 542 (♀). EMERY, 1895, *ibid.*, XXXV, p. 178 (2). FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 253 (2, ♀); 1913, Rev. Suisse Zool., XXI, p. 668 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 77 (♀). See p. 140.

Pheidole speculifrons DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 96.

Type locality: Anseba, ERITREA (Beccari).

BELGIAN CONGO: Faradje; Garamba (Lang and Chapin). CAMEROON: Johann-Albrechtshöhe (Conradt). BRITISH EAST AFRICA: Mt. Kenia, Amboni River, 1800 m. (Alluaud and Jeannel). ABYSSINIA: Arussi Galla, Ganale Gudda (V. Bottego). ERITREA: Ghinda; Melelia (K. Escherich). ANGLO-EGYPTIAN SUDAN: Bahr-el-Salaam (Magretti).

46i. Var. *ascarus* EMERY, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 62 (2). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 77 (2, ♀).

Type locality: Ghinda, ERITREA (Belli).

BRITISH EAST AFRICA: Nairobi, 1600 m. (Alluaud and Jeannel).

46j. Var. *cubangensis* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 413 (2).

Pheidole crassinoda var. *cubangensis* FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 306 (2).

Pheidole crassinoda var. *kubangensis* FOREL, in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 561 (♂).

Type locality: between the Cubango and Cuito Rivers, MOSSAMEDES (Baum and Van der Kellen).

BELGIAN CONGO: near Lisala, above Nouvelle Anvers (Kohl).

47. *Pheidole spinulosa* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 438 (♂, ♀, ♀, ♂). EMERY, 1916, Rev. Zool. Afr., IV, p. 250 (♂). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 419 and 453 (♂, ♀, ♀, ♂).

Type locality: BASUTOLAND (R. C. Wroughton).

47a. Subsp. *conigera* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 440 (♂).

Type locality: South Africa (Wood Mason).

47b. Subsp. *messalina* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 440 (♂, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 454 (♂, ♀).

Type locality: BASUTOLAND (R. C. Wroughton).

47b1. Var. *nexa* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 129 (♂, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 455, fig. 27, (♂, ♀, ♀), Pl. VI, fig. 70.

Type locality: RHODESIA (G. Arnold).

RHODESIA: Bulawayo; Plumtree (G. Arnold).

47b2. Var. *poweri* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 241 (♂, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 457 (♂, ♀).

Type locality: Kimberley, CAPE PROVINCE (Power).

48. *Pheidole squalida* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 364 (♂, ♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

49. *Pheidole strator* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 256 (♂, ♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 76 (♂, ♀). ALLUAUD and JEANNEL, 1914, Arch. Zool. Gén. Exp., LIII, p. 366.

Type locality: Ghinda, ERITREA (K. Escherich).

BRITISH EAST AFRICA: Shimoni; Blue Post Hotel, Kikuyu, 1520 m. (Alluaud and Jeannel).

49a. Subsp. *fugax* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 418 and 449, fig. 25a (♂, ♀).

Type locality: Amatongas Forest, PORTUGUESE EAST AFRICA (G. Arnold).

50. *Pheidole teneriffana* FOREL, 1893, Ann. Soc. Ent. Belgique, XXXVII, p. 465 (♂, ♀). EMERY, 1916, Rev. Zool. Afr., IV, p. 250. SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 378.

Pheidole teneriffana EMERY, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 61 (♂).

Type locality: Laguna, Teneriffe, CANARY ISLANDS (Medina).

BRITISH EAST AFRICA: Mombasa (G. Arnold). ERITREA: Massaua (Belli).

51. *Pheidole tenuinodis* MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 10 (♂, ♀). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 12 (♂, ♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 127 (♂, ♀); 1913, Deutsch. Ent. Zeitschr., Beih., p. 215 (♂, ♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 337 (♂, ♀); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 22 (♂). EMERY, 1916, Rev. Zool. Afr., IV, p. 245. ARNOLD, 1920, Ann. South African Mus., XIV, pp. 417 and 428 (♂, ♀, ♀).

Type locality: Bothaville, ORANGE FREE STATE (H. Brauns).

CAPE PROVINCE: Ladismith (H. Brauns); Kirstenbosch near the Cape (F. Silvestri). NATAL: Pinetown (I. Trägårdh). RHODESIA: Matoppo Hills (G. Arnold). GERMAN SOUTHWEST AFRICA: Okahandja (Casper); Grootfontein (von Erffer); Rooibank (L. Schultze). BECHUANALAND: Between Kooa and Thopane (L. Schultze).

51. Var. *sipapomæ* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 417 and 430 (2, ♀).

Type locality: Sipapoma, Ungusa River, SOUTHERN RHODESIA (G. Arnold).

SOUTHERN RHODESIA: Victoria Falls (G. Arnold).

51a. Subsp. *bothæ* (FOREL) SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 22 (2, ♀, ♀).

Pheidole foreli subsp. *bothæ* FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 62 (2, ♀).

Type locality: Bothaville, ORANGE FREE STATE (H. Brauns).

NATAL: Van Reenen; Balgown (F. Silvestri).

52. *Pheidole termitophila* FOREL, in Wasmann, 1904, 'Termitophilen aus dem Sudan, Res. Swed. Zool. Exped. Jägerskiöld,' XIII, p. 13 (2, ♀, ♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 339 (2).

Type locality: Kaka on the Ghrab el Aish, White Nile, ANGLO-EGYPTIAN SUDAN (I. Trägårdh).

SENEGAMBIA: Dakar (F. Silvestri).

52a. Subsp. *liberiensis* FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 270 (2, ♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 339 (2, ♀).

Type locality: Golah, LIBERIA.

FRENCH GUINEA: Mamou; Kakoulima (F. Silvestri).

53. *Pheidole tricarinata* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 340 (2).

Type locality: Camayenne, FRENCH GUINEA (F. Silvestri).

54. *Pheidole vanderveldi* FOREL, 1913, Rev. Zool. Afr., II, p. 326 (2, ♀). BEQUAERT, *ibid.*, p. 426. SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 23 (2, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 481 (2, ♀).

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

NATAL: Durban (I. Trägårdh).

55. *Pheidole ? variolosa* EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 113 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 97.

Type locality: Mogadiscio, SOMALILAND (Bricchetti-Robecchi).

SOMALILAND: Obbia (Pavesi).

56. *Pheidole victoris* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 127 (2, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 417 and 430 (2, ♀).

Type locality: Malindi, SOUTHERN RHODESIA (G. Arnold).

57. *Pheidole xocensis* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 213 (2, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 419 and 443, fig. 24 (2, ♀), Pl. vi, fig. 68.

Type locality: Xoce River, SOUTHERN RHODESIA (G. Arnold).

57a. Var. *bulawayensis* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 215 (2, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 445 (2, ♀).

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

Melissotarsini Emery**Melissotarsus EMERY**

Melissotarsus EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 378.

Genotype: *Melissotarsus beccarii* Emery, 1877.

1. **Melissotarsus beccarii** EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 379, fig. (2, ♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 74. FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 243 (♀, ♀, ♂). ARNOLD, 1916, Ann. South African Mus., XIV, p. 188 (♀, ♀, ♂), Pl. v, figs. 51, 51a, 51b.

Type locality: Keren, ERITREA (Beccari).

NATAL: Durban (Marley; C. B. Cooper).

2. **Melissotarsus emeryi** FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 133 (2).

Type locality: near the Colba River, ABYSSINIA (de Rothschild).

21. Var. **philipes** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 71 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 342 (♀).

Type locality: Taveta, BRITISH EAST AFRICA (Alluaud and Jeannel).

SENEGAMBIA: Thiès (F. Silvestri). GERMAN EAST AFRICA: Bismarckhügel, Mt. Kilimanjaro (Alluaud and Jeannel).

3. **Melissotarsus major** SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 85 (♀).

Type locality: Penge, BELGIAN CONGO (J. Bequaert).

4. **Melissotarsus weissi** SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 356, fig. 3 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

Myrmicariini Emery**Myrmicaria W. SAUNDERS**

Myrmicaria W. SAUNDERS, 1841, Trans. Ent. Soc. London, III, p. 57.

Physatta F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 171.

Heptacondylus F. SMITH, *ibid.*, p. 141.

Genotype: *Myrmicaria brunnea* W. Saunders, 1841.

1. **Myrmicaria baumi** FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 307 (♀); in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 562 (♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 5 (♀, ♂). STITZ, 1911, Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08), III, p. 383, fig. 5 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 266 (♀).

Type locality: between the Cubango and Cuito Rivers, MOSSAMEDES (Baum and Van der Kellen).

GERMAN SOUTHWEST AFRICA: (Lübbert). BECHUANALAND: Kooa; Severelela to Khakhea (L. Schultze).

2. **Myrmicaria eumenoides** (GERSTÄCKER) MAYR, 1866, Verh. Zool. Bot. Ges. Wien, XVI, p. 905 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 155. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 83. MAYR, 1896, Ent. Tidskr., XVII, p. 249. FOREL, 1897, Abhandl. Senckenberg. Naturf. Ges., XXI, p. 188 (♀). H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 42. BINGHAM, 1903, Ann. Mag. Nat. Hist., (7) XII, p. 62 (♀). SJÖSTEDT, 1904, 'I Västafrikas Urskogar,' p. 507. WASMANN, 1907, Deutsch. Ent. Zeitschr., p. 151. MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 17. FOREL, 1910, Ann. Soc. Ent. Belgique, LIV,

p. 446 (♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 5 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 133 (♀, ♂). WASMANN, 1911, Tijdschr. v. Ent., LIV, p. 201. STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 385, fig. 5 (♀, ♂). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 354 (♀, ♂). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 116 (♀); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 34 (♀, ♂). TRÄGÅRDH, *ibid.*, p. 45. ARNOLD, 1914, Proc. Rhodesia Sc. Assoc., XIII, p. 25. H. BRAUNS, *ibid.*, pp. 34, 36, 38, and 39. FOREL, 1916, Rev. Suisse Zool., XXIV, p. 404 (♀, ♂). ARNOLD, 1916, Ann. South African Mus., XIV, p. 263 (♀, ♂), Pl. VII, figs. 87, 87a, 89a, and 89b. WASMANN, 1918, Tijdschr. v. Ent., LX, (1917) p. 394.

Heptacondylus eumenoides GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 263 (♀); in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 514 (♀). MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 423. ROGER, 1863, 'Verzeich. Formicid.,' p. 28. GERSTÄCKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 356; 1873, in 'v. d. Decken's Reisen in Ost Afrika, Gliederthiere,' p. 358 (♀). RADOSZKOWSKY, 1881, Journ. Sci. Ac. Lisboa, VIII, No. 31, p. 198.

Physatta natalensis F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 172 (♀).

Heptacondylus sulcatus MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 756 (♀); 1863, *ibid.*, XIII, p. 423. ROGER, 1863, 'Verzeich. Formicid.,' p. 28.

Heptacondylus natalensis MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 423. ROGER, 1863, 'Verzeich. Formicid.,' p. 27.

Myrmicaria fodiens subsp. *eumenoides* EMERY, 1893, Rev. Suisse Zool., I, p. 219; 1895, Ann. Soc. Ent. France, LXIV, p. 42.

Myrmicaria fodiens var. *eumenoides* ZAVATTARI, 1909, 'Il Ruwenzori, Parte Scientif.,' I, p. 214 (♀).

Type locality: Tete, PORTUGUESE EAST AFRICA (Peters).

CAMEROON: Bibundi (Tessmann). SPANISH GUINEA: Alen (Tessmann). BELGIAN CONGO: Congo da Lemba (Luja); Romée near Stanleyville (Kohl); Kasindi; primary forest W. of Mt. Ruwenzori; Ituri; Aruwimi; Kwidjwi Island (Schubotz). SOUTHERN RHODESIA: (G. Arnold). ANGOLA: (Welwitsch). GERMAN SOUTHWEST AFRICA: (Lübbert). BECHUANALAND: Lehututu to Kang; Kooa (L. Schultze). BASUTOLAND. TRANSVAAL: Barberton (Rendall); Valdezia (P. Berthoud); Makapuu; Pretoria; Hamman's Kraal (E. Simon). NATAL: Stamford Hill; Amanzimtoti (I. Trägårdh); Durban (G. Arnold). DELAGOA: (Liengme; A. Müller). GERMAN EAST AFRICA: Mombo, Usambara (Sjöstedt); Bondei (C. W. Schmidt). BRITISH EAST AFRICA: Mombasa (v. d. Decken). UGANDA: Ibanda; Mitiana; Butiti; Nakitawa (Duke of Abruzzi). ZANZIBAR: (A. Voeltzkow).

2. Var. *fusca* STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 385 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 116 (♀).

Type locality: Boga, north of Mt. Ruwenzori, BELGIAN CONGO (Schubotz).

2a. Subsp. *occidentalis* SANTSCHI, 1920, 'Études Maladies Parasites Cacoyer S. Thomé,' X, p. 3, footnote (♀).

Type locality: Dimbroko, IVORY COAST (Le Moutl).

2b. Subsp. *opaciventris* (EMERY) FOREL, 1911, Rev. Zool. Afr., I, p. 276; 1913, Ann. Soc. Ent. Belgique, LVII, p. 354 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 116 (♀). See p. 143.

Myrmicaria opaciventris EMERY, 1893, Rev. Suisse Zool., I, p. 221 (♂, ♀, ♂).
ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 4 (♂, ♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXXVIII, (1909), p. 391 (♂, ♀, ♂). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 54 (♀); 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 263 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 133 (♀).

Myrmicaria eumenoides var. *opaciventris* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 393 (♂, ♀).

Type locality: BENGUELA (Buchner).

SIERRA LEONE: Samlia Falls, River N'Gamie (Mocquerys). CAMEROON: Mundame (Conradt); Yaunde (Zenker). FRENCH CONGO: Ogowé (Mocquerys); Brazzaville; Madingu (A. Weiss). BELGIAN CONGO: Leopoldville (Dubois); Duma; Libenge (Schubotz); Kimuenza (Schultze); Kimpoko (Büttner); Dungu to Niangara, village of Denge (Hutereau); Mälela; Thysville; Stanleyville; Avakubi; Medje; Akenge; Bafwabaka; Ngayu; Faradje (Lang and Chapin); Walikale to Labutu (J. Bequaert); Yakuluku (J. Rodhain). UGANDA: (Benoit). WEST AFRICA: (Ilg).

2b1. Var. *congolensis* (FOREL). See p. 146.

Myrmicaria eumenoides var. *congolensis* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 59 (♂); 1910, *ibid.*, LIV, p. 421; 1911, Rev. Zool. Afr., I, p. 276 (♂, ♀); 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 362 (♂); 1913, Ann. Soc. Ent. Belgique, LVII, p. 354 (♂, ♂); 1913, Rev. Zool. Afr., II, p. 337; 1916, Rev. Suisse Zool., XXIV, p. 404. WASMANN, 1918, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VII, p. 135; 1918, Tijdschr. v. Ent., LXI, p. 80.

Myrmicaria eumenoides subsp. *congolensis* J. BEQUAERT, 1913, Rev. Zool. Afr., II, p. 429. SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 342 (♂); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' pp. 115 and 116 (♂, ♀).

Type locality: Lower Congo (in stomach of *Manis temminckii*; Solon).

CAMEROON: Victoria (F. Silvestri); Soppo (v. Rothkirch). BELGIAN CONGO: Kondué (Lujá); Welgelegen; Kisantu (J. Bequaert); Lukula (Daniel); Mafungu; Congo da Lemba; Kiniati; Banza Masola (R. Mayné); Mayombe (de Briey). GERMAN EAST AFRICA: Amani (Zimmer); Kigali, near Issawe (Zimmer). UGANDA: Chacansengula to Kasengui (Bayer); Buzubizi; Unyoro Province, Region of Mubende, Lake Albert; Mt. Ruwenzori, Ibanda, 1400 m. (Alluaud). BRITISH EAST AFRICA: Mombasa (H. Prell).

2b2. Var. *consanguinea* (SANTSCHI).

Myrmicaria eumenoides subsp. *congolensis* var. *consanguinea* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' pp. 115, 116 (♂, ♀).

Type locality: Likoni, BRITISH EAST AFRICA (Alluaud and Jeannel).

BRITISH EAST AFRICA: Tiwi; Gazi; River Ramisi; Blue Post Hotel, Kikuyu (Alluaud and Jeannel).

2b3. Var. *crucheti* (SANTSCHI). See p. 146.

Myrmicaria eumenoides subsp. *congolensis* var. *crucheti* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 311 (♂); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 116 (♂).

Type locality: Cucala, BENGUELA (J. Cruchet).

GERMAN EAST AFRICA. BELGIAN CONGO: Leopoldville; Stanleyville; Avakubi; Ngayu (Lang and Chapin). CAMEROON: Metit (Schwab).

3. *Myrmicaria exigua* ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 320 (♂, ♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 155. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 134. FOREL, 1916, Rev. Suisse Zool., XXIV, p. 404.

Type locality: SIERRA LEONE (Mocquerys).

CAMEROON: (H. Brauns). BELGIAN CONGO: (Kohl). FRENCH CONGO: Gaboon.

3a. Var. *gracilis* (STITZ) SANTSCHI, 1920, Rev. Zool. Afr., VIII, p. 120.

Myrmicaria gracilis STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 133 (♀). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 445 (♀).

Type locality: Bibundi, CAMEROON (Tessmann).

CAMEROON: (L. v. Muralt).

3c. Var. *obscura* SANTSCHI, 1920, Rev. Zool. Afr., VIII, p. 120 (♀).

Type locality: Luali, BELGIAN CONGO (J. Bequaert).

3c. Var. *pulla* SANTSCHI, 1920, Rev. Zool. Afr., VIII, p. 119 (♀).

Type locality: Yambuya, BELGIAN CONGO (J. Bequaert).

3a. Var. *rufiventris* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 345 (♀); 1916, Rev. Suisse Zool., XXIV, p. 404 (♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

BELGIAN CONGO: Lumaliza; Makanga; Batiamponde (Kohl).

3d. Subsp. *kisangani* WM. M. WHEELER. See p. 148 (♀).

Type locality: Stanleyville, BELGIAN CONGO (Lang and Chapin).

4. *Myrmicaria laevior* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 445 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 267 (♀).

Type locality: NATAL (Haviland).

5. *Myrmicaria nigerrima* ARNOLD, 1916, Ann. South African Mus., XIV, p. 270 (♀), Pl. VII, fig. 92.

Type locality: Matoppo Hills, SOUTHERN RHODESIA (G. Arnold).

6. *Myrmicaria nigra* (MAYR) DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 155. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 42. FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 446 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 269 (♀, ♂), Pl. VII, fig. 91.

Heptacondylus niger MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 755 (♀); 1863, ibid., XIII, p. 423. ROGER, 1863, 'Verzeich. Formicid.,' p. 27. MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidae, p. 110 (♀).

Type locality: Cape of Good Hope (Novara Exped.).

CAPE PROVINCE: Cape Town (E. Simon); Montagu Pass (H. Brauns). NATAL: at 5500 ft. (R. C. Wroughton).

7. *Myrmicaria nitida* STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 133 (♀).

Type locality: Duala, CAMEROON (Schäfer).

SPANISH GUINEA: Nkolentanga (Tessmann).

According to Forel, 1910, Ann. Soc. Ent. Belgique, LIV, p. 421, this cannot be separated from *M. eumenoides* subsp. *opaciventris* var. *congolensis* (Forel).

7a. Var. *fumata* SANTSCHI, 1916, Bull. Soc. Ent. France, p. 242.

Myrmicaria nitida var. *brunnea* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 254 (♀) (nec *Myrmicaria brunnea* Saunders).

Type locality: Near Dimbroko, IVORY COAST (Posth).

8. *Myrmicaria salambo* WM. M. WHEELER. See p. 147 (♀).

Type locality: Garamba, BELGIAN CONGO (Lang and Chapin).

9. *Myrmicaria striata* STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 382, fig. 5 (♀). FOREL, 1913, Rev. Zool. Afr., II, p. 337 (♀). BEQUAERT, *ibid.*, p. 429. FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 212 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 267 (♀, ♀, ♂), Pl. VII, figs. 90a, b, c.

Myrmicaria striatula SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 116 (♀).

Type locality: Bukoba, Victoria Nyanza, GERMAN EAST AFRICA (Schubotz).

BELGIAN CONGO: Elisabethville (J. Bequaert). SOUTHERN RHODESIA: Bembesi; Plumtree; Bulawayo (G. Arnold). GERMAN EAST AFRICA: Kibosho, Mt. Kilimanjaro (C. Alluaud).

9i. Var. *verticalis* SANTSCHI, 1920, 'Études Maladies Parasites Cacaoyer S. Thomé,' p. 3, footnote (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

9a. Subsp. *buttgenbachi* FOREL, 1913, Rev. Zool. Afr., II, p. 337 (♀).

Type locality: not given; probably Katanga Region, BELGIAN CONGO (Buttgenbach).

9b. Subsp. *insularis* SANTSCHI, 1920, 'Études Maladies Parasites Cacaoyer S. Thomé,' p. 2 (♀, ? ♂).

Type locality: SAN THOMÉ (de Seabra).

10. *Myrmicaria tigrænsis* (GUÉRIN) F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 141. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 436. ROGER, 1863, 'Verzeich. Formicid.,' pp. 28 and 48. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 156.

Myrmica tigrænsis GUÉRIN, in Lefebvre, 1848, 'Voy. Abyssinie,' IV, Zool., pt. 6, p. 351 (♂), Pl. VII, fig. 7.

Type locality: ABYSSINIA.

Cardiocondyliini Emery

Cardiocondyla EMERY

Cardiocondyla EMERY, 1869, Ann. Accad. Natural. Napoli, (2) II, p. 20.

Emeryia FOREL, 1890, Ann. Soc. Ent. Belgique, XXXIV, C. R., p. cx (♂).

Genotype: *Cardiocondyla elegans* Emery, 1869.

1. *Cardiocondyla emeryi* FOREL, 1881, Mitth. München. Ent. Ver., V, p. 5 (♀). ERN. ANDRÉ, 1881, Ann. Soc. Ent. France, (6) I, p. 69 (♀), Pl. III, figs. 10, 11; 1883, 'Spec. Hym. Europ. Algérie,' II, pp. 328 and 329 (♀, ♂), Pl. XXI, figs. 9-12, and 14-15. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 71. FOREL, 1904, Ann. Soc. Ent. Belgique, XLVIII, p. 422 (♀, ♀, ♂); 1907, Ann. Mus. Nat. Hungarici, V, p. 17 (♂); 1913, Ann. Soc. Ent. Belgique, LVII, pp. 138 and 351 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 200 (♀, ♀, ♂), Pl. v, figs. 57, 57a. See pp. 150 and 1021.

Type locality: St. Thomas, WEST INDIES.

Tropicopolitan. MADEIRA: (E. Schmitz). BELGIAN CONGO: Congo da Lemba (R. Mayné); Thysville (J. Bequaert). RHODESIA: Bulawayo (G. Arnold). GERMAN EAST AFRICA: Arusha-chini (Katona). NATAL: Durban (C. B. Cooper).

1a. Subsp. *mahdii* KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 8 (♀).

Type locality: Khartum, ANGLO-EGYPTIAN SUDAN (Karawaiew).

Crematogastrini Emery

Crematogaster LUND¹

Crematogaster LUND, 1831, Ann. Sc. Nat., XXIII, p. 132. MAYR, 1855, Verh. Zool. Bot. Ges. Wien, V, p. 468.

Acrocalia MAYR, 1852, Verh. Zool. Bot. Ges. Wien, II, p. 147.

Formica (part) FABRICIUS, OLIVIER. *Atta* (part) BUCKLEY. *Crematogaster* MAYR, FOREL, SANTSCHI, etc. *Monomorium* (part) BUCKLEY.

Genotype: *Formica scutellaris* Olivier, 1791.

Subgenus 1. *Crematogaster* LUND, *sensu stricto*

Subgenotype: same as genotype.

1. *Crematogaster aegyptiaca* (MAYR). See p. 1022.

Crematogaster aegyptiaca MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 765 (♂); 1863, *ibid.*, XIII, p. 404. ROGER, 1863, 'Verzeich. Formicid.', p. 36. EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 535. ERN. ANDRÉ, 1883, 'Spec. Hym. Europ. Algérie,' II, p. 392 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 79. EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 10.

Type locality: EGYPT.

SOUTHERN ARABIA: ADEN. ERITREA: Assab (Doria and Beccari).

1a. Subsp. *senegalensis* (RÖGER) SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 344. See p. 1022.

Crematogaster senegalensis RÖGER, 1863, Berlin. Ent. Zeitschr., VII, p. 206 (♀); 1863, 'Verzeich. Formicid.', p. 36. EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 379 (♀). MAGRETTI, 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 245; 1884, Ann. Mus. Civ. Genova, XXI, p. 343 (♀, ♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 86. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 144; in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 7.

Type locality: Senegal (Sichel).

GOLD COAST. ANGLO-EGYPTIAN SUDAN: Suakin; Metemma; Sebderat; Ain (Magretti); Ondurman (I. Trägårdh). ERITREA: Bogos; Keren (Beccari).

1a₁. Var. *devincta* (SANTSCHI).

Crematogaster aegyptiaca subsp. *senegalensis* var. *devincta* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 343 (♀).

Type locality: Konakry, FRENCH CONGO (F. Silvestri).

1a₂. Var. *robusta* EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 10 (♀, ♀, ♂).

¹*Crematogaster nitidus* "Smith," recorded from Angola (Welwitsch) by Radoszkowsky, 1881, Journ. Sci. Ac., Lisboa, VIII, No. 31, p. 197, has apparently never been described.

Crematogaster ancipitula ? Forel, 1917, Bull. Soc. Vaudoise Sc. Nat., (5) LI, p. 252 (♀), was originally described as the ♀ of *C. wellmani* Forel. It has not been included in the Catalogue, since it is probably the ♀ of another form.

Crematogaster scutellaris (Olivier) subspecies *schmidtii* (Mayr) (*Acrocalia schmidtii* Mayr, 1852, Verh. Zool. Bot. Ges. Wien, II, p. 149, ♀, ♀; type locality: Wipbach, Carniola) is, according to Emery (1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 10), introduced in some of the harbors of the Red Sea: Port Sudan (*Crematogaster aegyptiaca* Karawaisew, 1911, Rev. Russe Ent., XI, p. 8, ♀); Massaua (*C. castaneae* Emery, 1877, Ann. Mus. Civ. Genova, IX, p. 379, ♀; 1881, *ibid.*, XVI, p. 535).

Sjöstedt (1908, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 116) mentions *Crematogaster schmidtii* "Mayr in litt." as occurring in swollen thorns of *Acacia drepanolobium*, near Kahé, Usambura. I have not found the description of a form bearing that name.

Crematogaster robusta EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 379 (♀, ♂).
DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 85.

Crematogaster aegyptiaca FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 257 (♀, ♂) (nec Mayr).

Type locality: Keren, ERITREA (Beccari).

ERITREA: Nefasit (K. Escherich; F. Silvestri).

2. *Crematogaster alulai* EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 11.

Crematogaster menileki var. *alulai* EMERY, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 61 (♀).

Crematogaster menileki var. *alulai* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 412 (♀).

Crematogaster menileki var. *alulai* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 93 (♀).

Type locality: Ghinda, ERITREA (Belli).

BRITISH EAST AFRICA: Shimoni (Alluaud and Jeannel).

2a. Subsp. *scrutans* (FOREL) EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 11 (♀).

Crematogaster scrutans FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 258 (♀).
ARNOLD, 1920, Ann. South African Mus., XIV, p. 518 (♀).

Type locality: Nefasit, ERITREA (K. Escherich; F. Silvestri).

3. *Crematogaster amabilis* (SANTSCHI).

Crematogaster amabilis SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 355 (♀); 1912, Bull. Soc. Ent. France, p. 412 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 91 (♀, ♀, ♂).

Type locality: Kilema, Mt. Kilimanjaro, 1440 m., GERMAN EAST AFRICA (Alluaud).

BRITISH EAST AFRICA: Crater of Longonot, Rift Valley, 2450 m.; Kijabe, Kikuyu Escarpment, 2100 m.; Molo, Mau Escarpment, 2420 m. (Alluaud and Jeannel).
GERMAN EAST AFRICA: Mt. Kilimanjaro, near the Bismarckhügel, 2700-2800 m. (Alluaud and Jeannel).

4. *Crematogaster brunneipennis* (ERN. ANDRÉ).

Crematogaster brunneipennis ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 323 (♀, ♀). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 144.

Crematogaster brunneipennis DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 80.

Type locality: SIERRA LEONE (Mocquerys).

4a. Subsp. *acaciæ* (FOREL).

Crematogaster acaciæ FOREL, 1892, Zool. Anzeiger, XV, p. 141 (♀). KELLER, ibid., p. 139. H. KOHL, 1909, Natur u. Offenbarung, LV, p. 152.

Crematogaster acaciæ FOREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 353 (♀).
DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 79.

Type locality: Faf Plain, Webi River, ABYSSINIA (C. Keller).

4a. Var. *generosa* (SANTSCHI).

Crematogaster acaciæ var. *generosa* SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 343 (♀, ♀).

Type locality: Durban, NATAL (G. Arnold).

4a₂. Var. **gloriosa** (SANTSCHI).

Crematogaster acaciæ var. *gloriosa* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 346 (♀).

Type locality: Boma, BELGIAN CONGO (F. Silvestri).

4a₃. Var. **victoriosa** (SANTSCHI). See p. 152.

Crematogaster acaciæ subsp. *victoriosa* SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 497 (♀).

Crematogaster acaciæ subsp. *victoriosa* SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 15. ARNOLD, 1920, Ann. South African Mus., XIV, pp. 488 and 500, fig. 35 (♀).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

BELGIAN CONGO: Zambi (J. Bequaert). NATAL: Durban (G. Arnold).

4b. Subsp. **omniparens** (FOREL), teste SANTSCHI in litt.

Crematogaster bruneipennis subsp. *omniparens* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 237 (♀).

Crematogaster bruneipennis subsp. *omniparens* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 488 and 499, fig. 34 (♀).

Type locality: Durban, NATAL (C. B. Cooper; G. Arnold).

5. **Crematogaster capensis** (MAYR).

Crematogaster capensis MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 764 (♀, ♀); 1863, *ibid.*, XIII, p. 404. ROGER, 1863, 'Verzeich. Formicid.,' p. 37. MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidæ, p. 103 (♀, ♀), Pl. IV, fig. 30. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 80. ARNOLD, 1920, Ann. South African Mus., XIV, p. 512 (♀, ♀).

Type locality: Cape of Good Hope (Novara Expedition).

5i. Var. **calens** (FOREL).

Crematogaster capensis var. *calens* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 435 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 513 (♀).

Type locality: Delagoa, PORTUGUESE EAST AFRICA (Liengme).

5a. Subsp. **tropicorum** (FOREL).

Crematogaster capensis subsp. *tropicorum* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 435 (♀).

Type locality: Ibo, PORTUGUESE EAST AFRICA.

6. **Crematogaster castanea** F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 136 (♀, ♀), Pl. IX, fig. 2. See p. 1022.

Crematogaster tricolor GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 263 (♀); in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 515 (♀), Pl. XXXII, fig. 10.

Crematogaster arboreus F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 138 (♀), Pl. XIV (nest) (*nec* Mayr, teste Arnold).

Crematogaster castanea ROGER, 1863, 'Verzeich. Formicid.,' p. 37. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 404. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 80. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 99. LAMBORN, 1920, Trans. Ent. Soc. London, (1919), Proc., p. liii. ARNOLD, 1920, Ann. South African Mus., XIV, p. 488 (♀, ♀).

Crematogaster arborea ROGER, 1863, 'Verzeich. Formicid.,' p. 37. ARNOLD, 1920, Ann. South African Mus., XIV, p. 506 (♀).

Crematogaster tricolor ROGER, 1863, 'Verzeich. Formicid.,' p. 37. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 405; 1866, *ibid.*, XXXVI, p. 365. FOREL,

1887, Mitth. Schweiz. Ent. Ges., VII, p. 388 (♀, ♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 87. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 99. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 139. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 40. MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 16. SjöSTEDT, 1908, *ibid.*, II, 8, pp. 99, 100, 117. SANTSCHI, 1912, Bull. Soc. Ent. France, p. 412 (♀). FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 213 (♀). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 21 (♀, ♀). TRÄGÄRDH, *ibid.*, p. 44. REICHENSPERGER, 1915, *ibid.*, V, pp. 8 and 20.

Cremastogaster tricolor var. *castanea* FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 80; 1911, Rev. Zool. Afr., III, p. 279 (♀). SANTSCHI, 1912, Bull. Soc. Ent. France, p. 412 (♀); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 22 (♀).

Cremastogaster castanea subsp. *tricolor* ARNOLD, 1920, Ann. South African Mus., XIV, p. 489 (♀, ♀, ♂), Pl. VI, figs. 72 and 72a.

Type locality: Port Natal, NATAL. Originally given by error as "Albania," but this was corrected by Roger in 1863.

BELGIAN CONGO: KONDUC (Luja). ORANGE FREE STATE. TRANSVAAL: Valdezia (P. Berthoud). NATAL: Durban; Stamford Hill (I. Trägårdh). CAPE PROVINCE: Cape of Good Hope; Port Alfred (C. B. Cooper; Hewitt); Grahamstown (G. Arnold). PORTUGUESE EAST AFRICA: Delagoa Bay (Liengme); Tete (Peters). ZULULAND: Dukudu (I. Trägårdh). GERMAN EAST AFRICA: Lindi (Lamborn); Tanga (H. Brauns); Mt. Kilimanjaro (Sjöstedt). BRITISH EAST AFRICA: Lamu Island; Fundu Island; W. Pemba (Voeltzkow).

61. Var. *simia* (FOREL).

Cremastogaster tricolor var. *simia* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 6 (♀, ♀, ♂). SANTSCHI, 1912, Bull. Soc. Ent. France, p. 412 (♀); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 22 (♀).

Cremastogaster castanea subsp. *tricolor* var. *simia* ARNOLD, 1920, Ann. South African Mus., XIV, p. 491 (♀, ♀, ♂).

Type locality: Chakamakue, between Lehututu and Letlake, BECHUANALAND (L. Schultze).

GERMAN SOUTHWEST AFRICA: (Peters). NATAL: Stamford Hill (I. Trägårdh).

6a. Subsp. *ferruginea* (FOREL) EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 12.

Cremastogaster ferruginea FOREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 353 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 81. FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 258 (♀).

Cremastogaster ferruginea EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 178 (♀); 1897, *ibid.*, XXXVIII, p. 598.

Cremastogaster tricolor subsp. *ferruginea* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♀).

Cremastogaster castanea subsp. *ferruginea* ARNOLD, 1920, Ann. South African Mus., XIV, p. 495 (♀).

Type locality: SOMALILAND (C. Keller).

ABYSSINIA: Lower Ganale (V. Bottego); Webi; Ganale; Ogaden; Daua (Ruspoli). ERITREA: Ghinda; Nefasit (K. Escherich).

6a₁. Var. *aquila* (FOREL).

Cremastogaster ferruginea var. *aquila* FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 139 (♀).

Cremastogaster tricolor subsp. *ferruginea* var. *aquila* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♂); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 87 (♂, ♂).

Type locality: South of Lake Rudolf, BRITISH EAST AFRICA (de Rothschild).

BRITISH EAST AFRICA: Naivasha, Rift Valley, 1900 m.; Gilgil, 1980 m.; Nairobi; Voi, Taita, 600 m. (Alluaud and Jeannel).

622. Var. *bruta* (SANTSCHI).

Cremastogaster tricolor subsp. *ferruginea* var. *bruta* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♂); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 22 (♂); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 87 (♂).

Cremastogaster castanea subsp. *rufonigra* var. *bruta* ARNOLD, 1920, Ann. South African Mus., XIV, p. 495 (♂).

Type locality: NATAL.

NATAL: Stamford Hill (I. Trägårdh). BRITISH EAST AFRICA: Naivasha, Rift Valley, 1400 m. (Alluaud and Jeannel). SOUTHERN RHODESIA: Springvale (G. Arnold).

623. Var. *durbanensis* (FOREL).

Cremastogaster ferruginea var. *durbanensis* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 234 (♂).

Cremastogaster castanea subsp. *tricolor* var. *durbanensis* ARNOLD, 1920, Ann. South African Mus., XIV, p. 491, fig. 32 (♂, ♀).

Type locality: Durban, NATAL (G. Arnold).

624. Var. *hararica* (FOREL).

Cremastogaster ferruginea var. *hararica* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 97 (♂).

Cremastogaster tricolor subsp. *ferruginea* var. *hararica* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♂).

Cremastogaster castanea subsp. *ferruginea* var. *hararica* EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 4 (♂).

Type locality: Harar, southern ABYSSINIA (Ilg).

ERITREA: Nefasit (F. Silvestri).

625. Var. *ulugurensis* (FOREL).

Cremastogaster ferruginea var. *ulugurensis* FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 369 (♂).

Cremastogaster tricolor subsp. *ferruginea* var. *ulugurensis* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♂) (*ulugurensis*); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 86 (♂, ♀, ♂).

Cremastogaster ferruginea var. *hararica* MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 15, Pl. II (nest).

Type locality: Uluguru Mts., GERMAN EAST AFRICA (Zimmer).

GERMAN EAST AFRICA: Morogoro (Zimmer); Kibonoto, Mt. Kilimanjaro (Sjöstedt). BRITISH EAST AFRICA: Nairobi, Kikuyu, 1700 m.; Blue Post Hotel, Tchania River, 1520 m. (Alluaud and Jeannel).

626. Var. *yambatensis* (FOREL).

Cremastogaster ferruginea var. *yambatensis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 353 (♂).

Type locality: Yambata, BELGIAN CONGO (R. Mayné).

6b. Subsp. **insidiosus** (SANTSCHI).

Cremastogaster castanea subsp. *insidiosus* SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 378 (♀).

Type locality: Balli Neck Pass, GERMAN SOUTHWEST AFRICA (G. Arnold).

6c. Subsp. **inversa** (FOREL).

Cremastogaster tricolor var. *inversa* FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 81 (♀); 1909, Ann. Soc. Ent. Belgique, LIII, p. 53 (♀); 1911, Rev. Zool. Afr., I, p. 279 (♀).

Cremastogaster tricolor var. *inversa* EMERY, 1912, Annuar. Mus. Zool. Univ. Napoli, N. S., III, No. 26, p. 2 (♀, ♀, ♂).

Cremastogaster tricolor subsp. *inversa* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 86 (♀).

Type locality: Patta Island, BRITISH EAST AFRICA (Voeltzkow).

BELGIAN CONGO: Kondué (Luja). SOUTHERN RHODESIA: Ciuma (Elena d'Aosta). GERMAN EAST AFRICA: between Lake Kivu and Victoria Nyanza (Elena d'Aosta). ZANZIBAR: (Deville). BRITISH EAST AFRICA: Kisumu, Kavirondo Bay (C. Alluaud); Manda Island (Voeltzkow).

6c. Var. **analis** (SANTSCHI). See p. 152.

Cremastogaster flaviventris var. *analis* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXXVIII, (1909), p. 372 (♀). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 433 (♀, ♀, ♂).

Cremastogaster tricolor subsp. *inversa* var. *analis* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

BELGIAN CONGO: Bafwasende to Avakubi (Lang and Chapin); Thysville (J. Bequaert).

6c. Var. **flaviventris** (SANTSCHI). See p. 152.

Cremastogaster flaviventris SANTSCHI, 1910, Ann. Soc. Ent. France, LXXXVIII, (1909), p. 370 (♀, ♀, ♂).

Cremastogaster tricolor subsp. *inversa* var. *flaviventris* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 86 (♀).

Type locality: Sankuru, BELGIAN CONGO (Luja).

CENTRAL UGANDA: (C. Alluaud). BELGIAN CONGO: Garamba (Lang and Chapin).

6d. Subsp. **museisapientiae** FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 272 (♀, ♀, ♂).

Cremastogaster tricolor subsp. *museisapientiae* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♀).

Type locality: Tropical Africa; exact locality unknown.

6e. Subsp. **rufimembrum** (SANTSCHI).

Cremastogaster tricolor subsp. *rufimembrum* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 431 (♀).

Type locality: Khutu Steppe, GERMAN EAST AFRICA (K. Schwartze).

6f. Subsp. **rufonigra** (EMERY).

Cremastogaster tricolor subsp. *rufonigra* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 27 (♀).

Cremastogaster tricolor var. *rufonigra* FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 22 (♀); in Schultz, 1910, 'Forschungsreise in Südafrika,' IV, p. 6 (♀).

Cremastogaster tricolor subsp. *rufoniger* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, pp. 124 and 352 (♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 342 (♀).

Cremastogaster castanea subsp. *rufonigra* ARNOLD, 1920, Ann. South African Mus., XIV, p. 492 (♀, ♀, ♂).

Type locality: Hebron, TRANSVAAL (E. Simon).

TRANSVAAL: HAMMAN'S KRAAL (E. Simon); Pretoria (F. Silvestri). SOUTHERN RHODESIA: common (G. Arnold). GERMAN EAST AFRICA: Kibosho (H. Brauns). BELGIAN CONGO: Elisabethville (Leplae). GERMAN SOUTHWEST AFRICA: (Lübbert). KALAHARI: Kakir (L. Schultze).

6f. Var. *busschodtsi* (EMERY).

Cremastogaster tricolor var. *busschodtsi* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 487 (♀). MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 80. FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 80.

Cremastogaster tricolor subsp. *rufoniger* var. *busschodtsi* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♀).

Type locality: Banana, BELGIAN CONGO (Busschodts).

BRITISH EAST AFRICA: Fundu Island, W. Pemba (Voeltzkow).

6f. Var. *cacodæmon* (FOREL).

Cremastogaster peringueyi subsp. *cacodæmon* (part) FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 234 (♀, ♀, ♂).

Cremastogaster castanea subsp. *rufonigra* var. *cacodæmon* ARNOLD, 1920, Ann. South African Mus., XIV, p. 494 (♀, ♀, ♂).

Type locality: Willowmore, CAPE PROVINCE (G. Arnold).

6f. Var. *mediorufa* (FOREL).

Cremastogaster tricolor var. *mediorufa* FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 81, footnote (♀, ♂). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 372 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 124 (♀, ♀, ♂).

Cremastogaster tricolor subsp. *rufonigra* var. *mediorufa* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 86 (♀).

Cremastogaster castanea subsp. *rufonigra* var. *medio-rufa* ARNOLD, 1920, Ann. South African Mus., XIV, p. 493 (♀, ♀, ♂), Pl. vi, figs. 77, 77a, and 77b.

Type locality: Shilouvane, TRANSVAAL (Junod).

BELGIAN CONGO: Bami Island (A. Weiss). BRITISH EAST AFRICA: Tiwi (Alluaud and Jeannel). CAPE PROVINCE: Aberdeen (H. Brauns).

6f. Var. *quisquilia* SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 502 (♀).

Cremastogaster castanea subsp. *rufonigra* var. *quisquilia* ARNOLD, 1920, Ann. South African Mus., XIV, p. 495 (♀).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

7. *Crematogaster censor* (FOREL).

Cremastogaster senegalensis subsp. *censor* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 434 (♀).

Type locality: Senegal (Eberli).

7a. Subsp. **junodi** (FOREL).

Cremastogaster censor subsp. *junodi* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 406 (♀).

Type locality: Shilouvane, TRANSVAAL (Junod).

8. ***Cremastogaster chiarinii*** EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 117; 1897, *ibid.*, XXXVIII, p. 598; 1899, *ibid.*, XXXIX, p. 500; 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 61. H. KOHL, 1909, Natur u. Offenbarung, LV, pp. 152 and 168.

Cremastogaster chiarinii EMERY, 1881, Ann. Mus. Civ. Genova, XVI, pp. 271, fig. (♂) and 535. DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 80. WASMANN, 1894, 'Verzeichn. Myrmecoph. Termitoph. Arthrop.', p. 120. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 99. WASMANN, 1896, Notes Leyden Mus., XVIII, p. 76. F. KOHL, 1907, Denkschr. Ak. Wiss. Wien, LXXI, p. 282 (♀, ♀). MAYR, 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. 1, p. 388; in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.', II, 8, p. 16 (♀, ♀, ♂). SJÖSTEDT, 1908, *ibid.*, II, 8, pp. 99 and 115. WASMANN, 1911, Tijdschr. v. Ent., LIV, p. 201. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicide', p. 85 (♀). REICHENSBERGER, 1915, Med. Göteborgs Mus. Zool. Afd., V, p. 35.

Type locality: Kaka, Schoa, ABYSSINIA (Antinori).

SOUTHERN ARABIA: Tes (Doria and Beccari); Ras Fartak (O. Simony); Gischin (W. Hein). ERITREA: Saati (Belli). SOMALILAND: Obbia (Bricchetti-Robecchi). ABYSSINIA: Dimé to Bass Narok (V. Bottego); southern Abyssinia (A. Ilg); Ogaden (Ruspoli). ANGLO-EGYPTIAN SUDAN: Khor Attar; Taufkia near Fashoda (F. Werner). BRITISH EAST AFRICA: Lusinga Island, Victoria Nyanza (C. Alluaud). GERMAN EAST AFRICA: Kibonoto, Mt. Kilimandjaro; Ngare na Nyuki R., Mt. Meru (Sjöstedt).

8₁. Var. **æthiops** (FOREL).

Cremastogaster chiarinii var. *æthiops* FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 142, footnote (♀).

Type locality: SOMALILAND (Keller).

8₂. Var. **affabilis** (FOREL).

Cremastogaster chiarinii var. *affabilis* FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 142 (♀).

Type locality: Daouele Region, SOMALILAND (de Rothschild).

ABYSSINIA: Harar (Ilg).

8₃. Var. **cincta** EMERY, 1896, Ann. Mus. Civ. Genova, XXXVII, p. 157 (♂); 1897, *ibid.*, XXXVIII, p. 598 (♀); 1899, *ibid.*, XXXIX, p. 500.

Cremastogaster chiarinii FOREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 353 (♂, ♀, ♂) (*nec* Emery). WASMANN, *ibid.*, p. 355.

Cremastogaster chiarinii FOREL, 1892, Zool. Anzeiger, XV, p. 140 (♂, ♀, ♂). KELLER, *ibid.*, p. 139. (*nec* Emery).

Cremastogaster chiarinii var. *cincta* FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 142, footnote.

Type locality: Lugh, SOMALILAND (V. Bottego).

ABYSSINIA: near the Webi River (C. Keller); Canale; Ogaden; Haucio; Leboi (Ruspoli).

84. Var. **v-nigrum** (FOREL).

Cremastogaster chiarinii var. *v-nigrum* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 434 (♀).

Type locality: CONGO.

8a. Subsp. **sellula** SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 431 (♀).

Type locality: Khutu Steppe, GERMAN EAST AFRICA (K. Schwartz).

8b. Subsp. **subsulcata** SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 432 (♀).

Cremastogaster chiarini KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 7 (♀).

Type locality: Khartum, ANGLO-EGYPTIAN SUDAN (Karawaiew).

8c. Subsp. **tædiosa** (FOREL) EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 12.

Cremastogaster tædiosa FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 98 (♀).

Type locality: Southern ABYSSINIA (Ifg).

ERITREA: Gomod, Saati (Belli); Algota (A. Fiori).

9. **Cremastogaster cicatriculosa** (ROGER).

Cremastogaster cicatriculosa ROGER, 1863, Berlin. Ent. Zeitschr., VII, p. 207 (♀); 1863, 'Verzeich. Formicid.,' p. 37. MAYR, 1866, Verh. Zool. Bot. Ges. Wien, XVI, p. 901 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 80.

Type locality: South Africa.

SIERRA LEONE.

10. **Cremastogaster constructor** EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 29 (♀), Pl. II, figs. 12, 13.

Cremastogaster constructor EMERY, in Mayr, 1895, Ann. Naturh. Hofmus. Wien, X, p. 135, footnote, fig. 3 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 487 and 535, fig. 49 (♀).

Type locality: Hamman's Kraal, TRANSVAAL (E. Simon).

10a. Var. **kirbyi** (MAYR).

Cremastogaster constructor var. *kirbyi* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, pp. 135 and 137 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 535 (♀).

Type locality: South Africa.

11. **Cremastogaster excisa** (MAYR) H. KOHL, 1909, Natur u. Offenbarung, LV, pp. 109 and 167. See p. 153.

Cremastogaster excisa MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 139 (♀). SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 501 (♀).

Type locality: LORINGO, FRENCH CONGO (H. Brauns).

SIERRA LEONE. BELGIAN CONGO: Zambi; Thysville (J. Bequaert); Faradje; near Lie (Lang and Chapin).

11a. Subsp. **andrei** (FOREL). See p. 153.

Cremastogaster excisa subsp. *andrei* FOREL, 1911, Rev. Zool. Afr., I, p. 277 (♀; according to Arnold in litt., with 11-jointed antennae).

Cremastogaster impressa subsp. *andrei* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 412 (♀).

Type locality: Congo da Lemba, BELGIAN CONGO (R. Mayné).

BELGIAN CONGO: Oso River; Sitaweza between Walikale and Lubutu (J. Bequaert).

11b. Subsp. *cavinota* (STITZ).

Cremastogaster excisa subsp. *cavinota* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 385, fig. 5 (♀).

Type locality: between Fort de Porsel and Fort Cranipel, FRENCH CONGO (Schubotz).

11c. Subsp. *caelestis* (SANTSCHI).

Cremastogaster caelestis SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 353 (♀). FOREL, 1913, Rev. Zool. Afr., II, p. 324 (♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 346 (♀).

Cremastogaster excisa subsp. *caelestis* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 342.

Type locality: CASAMANCE (Claveau).

BELGIAN CONGO: Kondué (Luja). FRENCH GUINEA: Camayenne (F. Silvestri).

11c. Var. *dirce* (FOREL).

Cremastogaster excisa subsp. *caelestis* var. *dirce* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 342 (♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

11d. Subsp. *impressa* (EMERY) SANTSCHI *in litt.*

Cremastogaster impressa EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 486 (♀, ♀). H. KOHL, 1909, Natur u. Offenbarung, LV, pp. 103 and 167.

Cremastogaster impressa FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, pp. 59 and 69 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 373 (♀). FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 270 (♀). SANTSCHI, 1912, Bull. Soc. Ent. France, p. 412 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 353 (♀); 1916, Rev. Suisse Zool., XXIV, p. 408 (♀, ♀, ♂).

Type locality: CAMEROON (Conradt).

IVORY COAST: Bessim (Bonhouse). BELGIAN CONGO: Lower Congo (Solon); Kilongalanga (Kohl); Stanleyville (Kohl); Isangi (Laurent); Congo da Lemba (R. Mayné). FRENCH CONGO: Brazzaville (A. Weiss). CAPE PROVINCE: Algoa Bay (H. Brauns).

11d. Var. *aglæa* SANTSCHI. See p. 153 (♀).

Type locality: Dimbroko, IVORY COAST (Le Mout).

11d. Var. *euphrosyne* SANTSCHI. See p. 153 (♀).

Type locality: Faradjé, BELGIAN CONGO (Lang and Chapin).

BELGIAN CONGO: Thysville (Lang and Chapin).

11d. Var. *sapora* (FOREL). See p. 154.

Cremastogaster impressa var. *sapora* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 408 (♀, ♀).

Type locality: BELGIAN CONGO (Kohl).

BELGIAN CONGO: Yakuluku (Lang and Chapin).

11e. Subsp. *lacustris* (SANTSCHI).

Cremastogaster excisa subsp. *lacustris* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 99 (♀).

Type locality: Lusinga Island, Victoria Nyanza, BRITISH EAST AFRICA (C. Alluaud).

11f. Subsp. **maynei** (FOREL).

Cremastogaster impressa subsp. *maynei* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 352 (♀).

Type locality: Congo da Lemba, BELGIAN CONGO (R. Mayné).

12. **Cremastogaster foraminiceps** (SANTSCHI).

Cremastogaster foraminiceps SANTSCHI, 1912, Bull. Soc. Ent. France, p. 412 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 89 (♀, ♀, ♂). RAF-FRAY, 1913, *ibid.*, Pselaphidae, p. 55, fig. (nest; without description).

Type locality: Nairobi, Kikuyu, 1700 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

BRITISH EAST AFRICA: River Burguret, Mt. Kenia, 2200 m.; Mt. Kenia, 2400 m.; River Amboni, 1800 m.; Voi (Alluaud and Jeannel).

12a. Subsp. **nigrans** (FOREL).

Cremastogaster foraminiceps subsp. *nigrans* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 338 (♀, ♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

12b. Subsp. **staitchi** (FOREL).

Cremastogaster foraminiceps subsp. *staitchi* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 339 (♀, ♀).

Type locality: Lumaliza, BELGIAN CONGO (Kohl).

13. **Cremastogaster gallicola** (FOREL).

Cremastogaster gallicola FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 95 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 487 and 531, fig. 47 (♀, ♀).

Type locality: DELAGOA (Liangme).

SOUTHERN RHODESIA: Somabula (G. Arnold).

13i. Var. **rauana** (FOREL).

Cremastogaster gallicola var. *rauana* FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 22 (♀).

Type locality: Moschi near the Rau River, Arushi-chini, GERMAN EAST AFRICA (Katona).

13a. Subsp. **latro** (FOREL).

Cremastogaster gallicola subsp. *latro* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 6 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 533 (♀).

Type locality: between Kgokong and Kang, BECHUANALAND (L. Schultze).

13b. Subsp. **spuria** (FOREL).

Cremastogaster gallicola subsp. *spuria* FOREL, 1913, Rev. Zool. Afr., II, p. 323 (♀). BEQUAERT, *ibid.*, p. 425.

Type locality: Elisabethville, BELGIAN CONGO (J. Bequaert).

14. **Cremastogaster gerstaeckeri** (DALLA TORRE).

Cremastogaster cephalotes GERSTAECKER, 1871, Arch. f. Naturg., XXXVII, I, p. 356 (♂, ♀) (nec F. Smith); 1873, in 'v. d. Decken's Reisen in Ostafrika, Gliederthiere,' p. 358 (♂, ♀).

Cremastogaster cephalotes KELLER, 1892, Zool. Anzeiger, XV, p. 137. H. KOHL, 1909, Natur u. Offenbarung, LV, p. 152.

Cremastogaster gerstaeckeri DALLA TORRE, 1892, Wien. Ent. Zeitg., XI, March, p. 90; 1893, 'Cat. Hym.,' VII, p. 82.

- Crematogaster gerstäckeri* FOREL, 1892, Zool. Anzeiger, XV, April, p. 141.
ZAVATTARI, 1909, 'Il Ruwenzori, Parte Scientif.', I, p. 214 (♀, ♀).
- Crematogaster gerstäckeri* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 137 (♀).
H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 40.
Type locality: Mombasa, BRITISH EAST AFRICA (v. d. Decken).
UGANDA: Butiti (Duke of Abruzzi).
15. *Crematogaster godefreyi* (FOREL).
Crematogaster foraminiceps subsp. *godefreyi* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 237 (♀).
Crematogaster vulcania subsp. *godefreyi* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 406.
Crematogaster newillei subsp. *carininotum* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), pp. 279 and 280 (♀).
Crematogaster sjøstedti subsp. *godefreyi* SANTSCHI, 1918, Bull. Soc. Ent. France, p. 184.
Crematogaster godefreyi ARNOLD, 1920, Ann. South African Mus., XIV, p. 503, fig. 37 (♀).
Crematogaster newillei subsp. *carininotum* ARNOLD, 1920, Ann. South African Mus., XIV, p. 499 (♀).
Type locality: King William's Town, CAPE PROVINCE (R. Godfrey).
RHODESIA: Bulawayo (G. Arnold).
15. Var. *arnoldi* (FOREL).
Crematogaster arnoldi FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 238 (♀).
Crematogaster godefreyi var. *arnoldi* ARNOLD, 1920, Ann. South African Mus., XIV, p. 504, fig. 38 (♀).
Type locality: Durban, NATAL (Marley).
16. *Crematogaster gutenbergi* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 433 (♀).
Type locality: GERMAN SOUTHWEST AFRICA.
17. *Crematogaster ilgi* (FOREL) SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 502.
Crematogaster scrutans var. *ilgi* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 259 (♀).
ARNOLD, 1920, Ann. South African Mus., XIV, p. 519 (♀).
Type locality: Western ABYSSINIA (Ilg).
TRANSVAAL: Zoutpansberg.
18. *Crematogaster impressiceps* (MAYR). See p. 154.
Crematogaster impressiceps MAYR, 1902, Verh. Zool. Bot. Ges. Wien, LII, p. 294 (♀).
Type locality: FERNANDO PO (R. Buchholz).
CAMEROON: Abo (R. Buchholz). BELGIAN CONGO: Panga; Faradje (Lang and Chapin).
18. Var. *frontalis* SANTSCHI. See p. 154 (♀).
Type locality: Malela, BELGIAN CONGO (Lang and Chapin).
BELGIAN CONGO: Kunga near Malela (Lang, Chapin and J. Bequaert).
18. Var. *longiscapa* (STITZ).
Crematogaster impressiceps var. *longiscapa* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 384 (♀), Pl. xx (nest).
Type locality: Duma, BELGIAN CONGO (Schubotz).

18. Var. *lujana* (FOREL).

Crematogaster impressiceps var. *lujana* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 340 (♂).

Type locality: KONDUE, BELGIAN CONGO (LUIJA).

19. *Crematogaster inconspicua* (MAYR).

Crematogaster inconspicua MAYR, 1896, Ent. Tidskr., XVII, p. 246 (♂, ♀).
AURIVILLIUS, *ibid.*, p. 253, Pl. IV, fig. 1 (nest). SJÖSTEDT, 1904, I Västafrikas Urskogar, p. 504, fig. 1 (nest).

Type locality: CAMEROON (Sjöstedt).

19a. Subsp. *incorrecta* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV (1916), p. 280 (♂).

Type locality: GABOON, FRENCH CONGO (F. Faure).

20. *Crematogaster inermis* (MAYR).

Crematogaster inermis MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 766 (♂); 1895, Ann. Naturh. Hofmus. Wien, X, p. 137 (♂). KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 8 (♂). ARNOLD, 1920, Ann. South African Mus., XIV, p. 487 (♂).

Type locality: Sinai Peninsula (v. Frauenfeld).

ANGLO-EGYPTIAN SUDAN: Port Sudan (Karawaiew).

20a. Subsp. *delagoensis* (FOREL).

Crematogaster inermis subsp. *delagoensis* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 99 (♂). ARNOLD, 1920, Ann. South African Mus., XIV, p. 539, fig. 51 (♂).

Type locality: DELAGOA (Liengme).

21. *Crematogaster melanogaster* (EMERY).

Crematogaster arborea MAYR, 1866, Verh. Zool. Bot. Ges. Wien, XVI, p. 900 (♂, ♀), Pl. xx, fig. 10; 1886, *ibid.*, XXXVI, p. 360; 1896, Ent. Tidskr., XVII, p. 248, footnote (♂). ARNOLD, 1920, Ann. South African Mus., XIV, p. 507 (♂). (*nec C. arborea* F. Smith, *teste* Mayr and Arnold).

Crematogaster arborea subsp. *melanogaster* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 29 (♂, ♀).

Crematogaster arborea subsp. *melanogaster* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 7 (♂, ♀); 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 234 (♂). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 486 and 508, fig. 39 (♂, ♀).

Type locality: Port Natal, NATAL (Gueinzus).

CAPE PROVINCE: De Aar (E. Simon); Conway (Hewitt); Willowmore (H. Brauns); Little Namaland (L. Péringuez); Kamagga (L. Schultze).

22. *Crematogaster menileki* (FOREL) EMERY, 1899, Ann. Mus. Civ. Genova, XXXIX, p. 500.

Crematogaster menileki FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 97 (♂).

Crematogaster menileki SANTSCHI, 1912, Bull. Soc. Ent. France, p. 412 (♂).

Type locality: Schoa, southern ABYSSINIA (Ilg).

ABYSSINIA: Buditu to Dimé (V. Bottego).

22a. Subsp. *occidentalis* (MAYR).

Crematogaster menileki subsp. *occidentalis* MAYR, 1902, Verh. Zool. Bot. Ges. Wien, LII, p. 293 (♂). SANTSCHI, 1912, Bull. Soc. Ent. France, p. 412 (♂).

Crematogaster menileki subsp. *occidentalis* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 252 (♂).

Type locality: FERNANDO PO (R. Buchholz).

GOLD COAST. DAHOMEY: Kouande (Desanti).

- 22a₁. Var. **atrigaster** (FOREL).
Crematogaster menileki subsp. *occidentalis* var. *atrigaster* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 353 (♀).
Type locality: Congo da Lemba, BELGIAN CONGO (R. Mayné).
- 22a₂. Var. **brazzai** (SANTSCHI).
Crematogaster impressa subsp. *brazzai* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 373 (♂, ♀).
Crematogaster menileki subsp. *occidentalis* var. *brazzai* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 412 (♂).
Type locality: Brazzaville, FRENCH CONGO (A. Weiss).
- 22b. Subsp. **proserpina** SANTSCHI. See p. 154 (♂).
Type locality: Malela, BELGIAN CONGO (Lang and Chapin).
- 22b₁. Var. **pluton** (SANTSCHI).
Crematogaster menileki subsp. *proserpina* var. *pluton* SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 87 (♀).
Type locality: Zambi, BELGIAN CONGO (J. Bequaert).
- 22c. Subsp. **satan** (FOREL).
Crematogaster menileki subsp. *satan* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 407 (♀).
Type locality: BELGIAN CONGO (Kohl).
- 22c₁. Var. **satanula** (FOREL).
Crematogaster menileki subsp. *satan* var. *satanula* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 407 (♀).
Type locality: St. Gabriel, BELGIAN CONGO (Kohl).
23. **Crematogaster mimosæ** (SANTSCHI).
Crematogaster mimosæ SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 87, fig. 11 (♀).
Type locality: Mt. Kenia, 2000 m., BRITISH EAST AFRICA (C. Alluaud).
24. **Crematogaster misella** (ARNOLD).
Crematogaster gallicola subsp. *sjöstedti* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 237 (♀) (*nec* Mayr).
Crematogaster misella ARNOLD, 1920, Ann. South African Mus., XIV, p. 513, fig. 40a (♀).
Type locality: Bulawayo, RHODESIA (G. Arnold).
25. **Crematogaster monticola** (ARNOLD).
Crematogaster monticola ARNOLD, 1920, Ann. South African Mus., XIV, p. 517, fig. 40b (♀).
Type locality: Mountains of NATAL (Haviland).
26. **Crematogaster newillei** (FOREL) SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 279 (♀).
Crematogaster newillei FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 140 (♀).
SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 497 (♀).
Type locality: Uomber, southern ABYSSINIA (de Rothschild).
- 26a. Subsp. **cooperi** (FOREL) SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 280 (♀).

Crematogaster newillei subsp. *cooperi* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 233 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 488 and 497, fig. 33 (♀, ♀), Pl. VI, figs. 76 and 76a.

Type locality: Durban, NATAL (C. B. Cooper).

26a. Var. *ingravis* (FOREL) SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 280 (♀).

Crematogaster newillei subsp. *cooperi* var. *ingravis* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 341 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 498 (♀).

Type locality: Durban, NATAL (C. B. Cooper).

27. *Crematogaster nigriceps* EMERY, 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 600, fig. (♀).

Type locality: BANAS, SOMALILAND (Ruspoli).

ABYSSINIA: HAUACIO (Ruspoli).

27a. Subsp. *prelli* (FOREL).

Crematogaster nigriceps subsp. *prelli* FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 368 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 98 (♀).

Type locality: Kabe Steppe, GERMAN EAST AFRICA (H. Prell).

UGANDA: UNYORO Province, region of Lake Albert (C. Alluaud).

28. *Crematogaster nigronitens* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 281 (♀).

Crematogaster nigronitens ARNOLD, 1920, Ann. South African Mus., XIV, pp. 486 and 541, fig. 53 (♀).

Type locality: Matoppos Hills, SOUTHERN RHODESIA (G. Arnold).

29. *Crematogaster ochraceiventris* (STITZ).

Crematogaster ochraceiventris STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 391, fig. 8 (♀).

Type locality: Fort Crampel, FRENCH CONGO (Schubotz).

30. *Crematogaster opaciceps* (MAYR).

Crematogaster opaciceps MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 12 (♀), Pl. I, fig. 1; Pl. II, fig. 3 (nest). ARNOLD, 1920, Ann. South African Mus., XIV, p. 514 (♀).

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

30a. Var. *clepens* (FOREL).

Crematogaster opaciceps var. *clepens* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 353 (♀).

Type locality: Congo da Lemba, BELGIAN CONGO (R. Mayné).

30a. Subsp. *defleta* (FOREL).

Crematogaster opaciceps subsp. *defleta* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 434 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 515 (♀).

Type locality: TRANSVAAL (C. Keller).

31. *Crematogaster orobia* SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 343 (♀).

Type locality: Matroosberg, 4500 ft., CAPE PROVINCE (R. W. Tucker).

32. *Crematogaster peringueyi* EMERY, 1895, Ann. Soc. Ent. France, LXIII, p. 27 (♀, ♀, ♂), Pl. II, figs. 16 and 17. RAFFRAY, 1910, Ann. South African Mus., V, 8, pp. 422 and 423.

Cremastogaster peringueyi WASMANN, 1897, Wien. Ent. Zeitg., XVI, p. 201; 1898, *ibid.*, XVII, p. 97; 1899, Deutsch. Ent. Zeitschr., p. 170. MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 14, Pl. II, fig. 4 (nest). FÖREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 7 (♂, ♀, ♂); 1913, Deutsch. Ent. Zeitschr., Beih., p. 213 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 509 (♂, ♀, ♂), Pl. VI, fig. 83.

Type locality: CAPE TOWN, CAPE PROVINCE (E. Simon; L. Péringuey).

CAPE PROVINCE: Port Elizabeth (H. Brauns); Port Alfred (Hewitt); Port Nolloth; Steinkopf (L. Schultze); Pirie Forest (G. Rogers).

324. Var. *angustior* (ARNOLD).

Cremastogaster peringueyi subsp. *cacodæmon* (part) FÖREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 234 (♂, ♀, ♂).

Cremastogaster peringueyi var. *angustior* ARNOLD, 1920, Ann. South African Mus., XIV, p. 510, fig. 40 (♂, ♀, ♂).

Type locality: Durban, NATAL (C. B. Cooper).

325. Var. *cacochyma* (FÖREL).

Cremastogaster peringueyi subsp. *cacochyma* FÖREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 235 (♂, ♀).

Cremastogaster peringueyi var. *cacochyma* ARNOLD, 1920, Ann. South African Mus., XIV, p. 512 (♂, ♀).

Type locality: not given; probably CAPE PROVINCE.

326. Var. *gedeon* (FÖREL).

Cremastogaster peringueyi subsp. *cacodæmon* var. *gedeon* FÖREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 341 (♂).

Cremastogaster peringueyi var. *gedeon* ARNOLD, 1920, Ann. South African Mus., XIV, p. 512 (♂).

Type locality: Pietermaritzburg, NATAL (R. Bayer).

NATAL: New Hanover Rail. (C. B. Hardenburg).

33. *Crematogaster rivai* EMERY, 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 598, fig. (♂).

Type locality: Webi, ABYSSINIA (Ruspoli).

ABYSSINIA: Ganale (Ruspoli).

34. *Crematogaster ruspolii* FÖREL, 1892, Zool. Anzeiger, XV, p. 142 (♂, ♂). KELLER, *ibid.*, p. 139. H. KOHL, 1909, Natur u. Offenbarung, LV, pp. 152 and 168.

Cremastogaster ruspolii FÖREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 353 (♂, ♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 85.

Type locality: Faf, ABYSSINIA (C. Keller).

ABYSSINIA: Webi River (C. Keller).

341. Var. *atriscapis* (FÖREL).

Cremastogaster ruspolii var. *atriscapis* FÖREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 338 (♂); 1916, Rev. Suisse Zool., XXIV, p. 405 (♂).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

35. *Crematogaster schultzei* (FÖREL).

Cremastogaster schultzei FÖREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 6 (♂, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 486 and 519, fig. 41 (♂, ♀).

Type locality: Steinkopf, CAPE PROVINCE (L. Schultze).

CAPE PROVINCE: Kamaggas; Prince of Wales Bay (L. Schultze).

36. *Crematogaster sewellii* (FOREL). See p. 1023.

36a. Subsp. *acis* (FOREL).

Crematogaster sewellii subsp. *acis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 353 (♀).

Type locality: Congo da Lemba, BELGIAN CONGO (R. Mayné).

36b. Subsp. *marnoi* (MAYR). See p. 1024.

Crematogaster sewellii var. *marnoi* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 138 (♀); 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. I, p. 388.

Crematogaster swelli subsp. *marnoi* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 376 (♀).

Type locality: Sudan (E. Marno).

FRENCH CONGO: Gomba (A. Weiss). ANGLO-EGYPTIAN SUDAN: Mongalla (F. Werner).

37. *Crematogaster sjöstedti* (MAYR).

Crematogaster sjöstedti MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 17 (♀). Sjöstedt, 1908, *ibid.*, II, 8, p. 116. SANTSCHI, 1918, Bull. Soc. Ent. France, p. 184.

Crematogaster gallicola subsp. *sjöstedti* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 124. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 94 (♀).

Type locality: Usambara, near Tanda, GERMAN EAST AFRICA (Sjöstedt).

GERMAN EAST AFRICA: New Moschi, Mt. Kilimanjaro, 800 m. (Alluaud and Jeannel).

37i. Var. *maledicta* (FOREL).

Crematogaster excisa var. *maledicta* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 236 (♀, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 502, fig. 36 (♀, ♀).

Crematogaster sjöstedti var. *maledicta* SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 15.

Type locality: Bulawayo, RHODESIA (G. Arnold).

37i. Var. *pulla* (SANTSCHI).

Crematogaster gallicola subsp. *sjöstedti* var. *pulla* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 94 (♀).

Type locality: Mbuyuni, Pori, 1110 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

BRITISH EAST AFRICA: Taveta, 750 m. (Alluaud and Jeannel).

37s. Var. *rufescens* (SANTSCHI).

Crematogaster sjöstedti subsp. *bulawayensis* var. *rufescens* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 236 (♀).

Type locality: NATAL (Haviland).

37a. Subsp. *kohliella* (FOREL).

Crematogaster excisa subsp. *andrei* var. *kohliella* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 409 (♀).

Crematogaster sjöstedti subsp. *kohliella* SANTSCHI, 1918, Bull. Soc. Ent. France, p. 185.

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

38. *Crematogaster solers* (FOREL).

Crematogaster solers FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 7 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 516 (♀).

Type locality: Arasab River, near Kubub, GERMAN SOUTHWEST AFRICA (L. Schultze).

39. *Crematogaster stigmata* (SANTSCHI).

Crematogaster stigmata SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 344, fig. 15 (♀, ♂).

Type locality: Olokemeji, NIGERIA (F. Silvestri).

40. *Crematogaster vulcania* (SANTSCHI).

Crematogaster vulcania SANTSCHI, 1912, Bull. Soc. Ent. France, p. 413 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 94, fig. 13 (♀, ♀, ♂). LECERF, 1914, *ibid.*, 'Chenilles Galles Formicides,' pp. 5, 11, 13, 17, 18, 23, and 25.

Type locality: Longonot Neck, 2140 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

41. *Crematogaster werneri* (MAYR).

Crematogaster werneri MAYR, 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. 1, p. 388 (♀).

Type locality: Gondokoro, UGANDA (F. Werner).

41. Var. *cacozela* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 253.

Crematogaster cacozela SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar., VIII, p. 342 (♀).

Type locality: Camayenne, FRENCH GUINEA (F. Silvestri).

41. Var. *pasithea* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 253 (♀).

Type locality: LOWER DAHOMEY (Desanti).

Subgenus 2. *Sphaerocrema* SANTSCHI

Crematogaster subg. *Sphaerocrema* SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.

Subgenotype: *Crematogaster kneri* Mayr, 1862.

42. *Crematogaster (Sphaerocrema) bequaerti* (FOREL).

Crematogaster bequaerti FOREL, 1913, Rev. Zool. Afr., II, p. 319 (♀, ♀). BEQUAERT, *ibid.*, p. 425. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 93, fig. 12a (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 348, fig. 17a (♀).

Crematogaster (Sphaerocrema) bequaerti SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

42. Var. *atraplex* SANTSCHI. See p. 155 (♀).

Type locality: Yakuluku, BELGIAN CONGO (Lang and Chapin).

42. Var. *gerardi* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 252 (♀).

Type locality: Katakai, BELGIAN CONGO (Gérard).

42. Var. *mutabilis* (SANTSCHI).

Crematogaster bequaerti var. *mutabilis* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 93, fig. 12b (♀).

Type locality: Lumbwa, Mau Escarpment, 1900 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

42a. Subsp. *ludia* (FOREL).

Crematogaster bequaerti subsp. *ludia* FOREL, 1913, Rev. Zool. Afr., II, p. 321 (♀).

Type locality: Lake Kabwe, BELGIAN CONGO (J. Bequaert).

43. **Crematogaster (Sphaerocrema) chlorotica** EMERY.

Crematogaster chlorotica EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 483 (♂, ♀).

Crematogaster chlorotica SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 374 (♀). FOREL, 1913, Rev. Suisse Zool., XXI, p. 669 (♀).

Crematogaster (Sphaerocrema) chlorotica SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.

Type locality: CAMEROON (Conradt).

SIERRA LEONE. CAMEROON: Johann-Albrechtshöhe (Conradt). FRENCH CONGO: Brazzaville (A. Weiss).

44. **Crematogaster (Sphaerocrema) concava** EMERY. See p. 155.

Crematogaster concava EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 480 (♂; nec ♀).

Crematogaster concava SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 375 (♂, ♀, ♂). FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 338 (♀); 1916, Rev. Suisse Zool., XXIV, p. 404 (♂, ♀, ♂). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 384 (♀).

Crematogaster (Sphaerocrema) concava SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.

Type locality: Popokabaka, BELGIAN CONGO.

CAMEROON. FRENCH CONGO: Brazzaville (A. Weiss); Fort de Possel to Fort Crampel (Schubotz). BELGIAN CONGO: St. Gabriel (Kohl); Lukolela to Basoko; Stanleyville; Akenge (Lang and Chapin).

45. **Crematogaster (Sphaerocrema) gambiensis** (ERN. ANDRÉ).

Crematogaster gambiensis ERN. ANDRÉ, 1889, Rev. d'Ent. Caen, VIII, p. 228 (♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 82. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 137; 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. 1, p. 387. FOREL, 1913, Rev. Zool. Afr., II, p. 324 (♀, ♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 346 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 529 (♀).

Crematogaster gambiensis var. *longiruga* FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 140 (♀, ♀, ♂).

Type locality: GAMBIA (Mocquerys).

FRENCH GUINEA: Konakry (F. Silvestri). SIERRA LEONE: Samlia Falls, River N'Gamie. GOLD COAST. SLAVE COAST. BELGIAN CONGO: Kwezi to Kilo (Bayer). ANGLO-EGYPTIAN SUDAN: Mongalla (F. Werner). UGANDA: Gondokoro (F. Werner). BRITISH EAST AFRICA: Rendilé, Mt. Karoli; Mt. Nyiro (de Rothschild).

45. Var. **krantziana** (FOREL).

Crematogaster gambiensis var. *krantziana* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 234 (♂, ♂). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 487 and 530 (♂, ♀, ♂).

Type locality: Krantz Kloof near Durban, NATAL (Marley).

45. Var. **transversiruga** (SANTSCHI).

Crematogaster gambiensis var. *transversiruga* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 346 (♀).

Type locality: Lado, Redjaf, ANGLO-EGYPTIAN SUDAN (Reichensperger).

46. **Crematogaster (Sphaerocrema) kneri** (MAYR).

Crematogaster kneri MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 764 (♀); 1863, ibid., XIII, p. 404. ROGER, 1863, 'Verzeich. Formicid.,' p. 36. DALLA TORRE,

1893, 'Cat. Hym.,' VII, p. 82. FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 433 (♂, ♀, ♂); 1913, *ibid.*, LVII, p. 125 (♂, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 487 and 525, fig. 45 (♂, ♀).

Crematogaster kneri EMERY, 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 598 (♂); 1899, Ann. Soc. Ent. Belgique, XLIII, p. 483.

Crematogaster (Sphaerocrema) kneri SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.

Type locality: Akwapim Mts., GOLD COAST.

ABYSSINIA: Ganale (Ruspoli). ORANGE FREE STATE: Bothaville (H. Brauns).

NATAL: (Haviland). BELGIAN CONGO: (Seeldrayers).

46. Var. *amita* (FOREL).

Crematogaster kneri var. *amita* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 125 (♂). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 487 and 527 (♂), Pl. VI, fig. 75.

Type locality: Bulawayo, RHODESIA (G. Arnold).

46a. Var. *matabele* (ARNOLD).

Crematogaster kneri var. *matabele* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 487 and 528, fig. 46 (♂).

Type locality: Hillside, Bulawayo, RHODESIA (G. Arnold).

46a. Subsp. *hottentota* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 482 (♂).

Crematogaster kneri EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 27 (♂) (*nec* Mayr).

Crematogaster kneri subsp. *hottentota* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 125 (♂). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 487 and 528 (♂).

Type locality: Vrijburg, CAPE PROVINCE (E. Simon).

SOUTHERN RHODESIA: Bedza, Matoppo Hills; Springvale; Sipapoma (G. Arnold).

ORANGE FREE STATE: Bloemfontein (E. Simon).

47. *Crematogaster (Sphaerocrema) libengensis* (STITZ).

Crematogaster libengensis STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 389, fig. 7 (♂).

Type locality: Libenge, BELGIAN CONGO (Schubotz).

48. *Crematogaster (Sphaerocrema) luctans* (FOREL).

Crematogaster luctans FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 22 (♂, ♀).

Crematogaster luctans EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 12.

Crematogaster (Sphaerocrema) luctans SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.

Type locality: Mto-ya-Kifaru, GERMAN EAST AFRICA (Katona).

ERITREA: Nefasit (F. Silvestri).

49. *Crematogaster (Sphaerocrema) nigeriensis* (SANTSCHI).

Crematogaster nigeriensis SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 346, fig. 16 (♂).

Crematogaster (Sphaerocrema) nigeriensis SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.

Type locality: Lagos, SOUTHERN NIGERIA (F. Silvestri).

49a. Var. *wilniger* (FOREL).

Crematogaster nigeriensis var. *wilniger* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 405 (♂, ♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

50. **Crematogaster (Sphærocrema) pronotalis** SANTSCHI.
Crematogaster pronotalis SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 347, fig. 17b (♀).
Crematogaster pronotalis SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 252 (♀).
Crematogaster (Sphærocrema) pronotalis SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.
 Type locality: Olokemeji, NIGERIA (F. Silvestri).
 50_a. Var. **behanzini** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 251 (♀).
 Type locality: Kotonou, DAHOMEY.
 50_b. Var. **dakarensis** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 252 (♀).
Crematogaster pronotalis var. *dakarensis* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 349 (♀).
 Type locality: Dakar, SENEGAMBIA (F. Silvestri).
 50_c. Var. **funerea** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 251 (♀).
 Type locality: Molundu, CAMEROON (Reichensperger).
 50_d. Var. **liebknechti** (FOREL). See p. 156.
Crematogaster pronotalis var. *liebknechti* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 340 (♀).
 Type locality: St. Gabriel, BELGIAN CONGO (Kohl).
 BELGIAN CONGO: Yakuluku; Garamba (Lang and Chapin).
 51. **Crematogaster (Sphærocrema) rugosa** (ERN. ANDRÉ) SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 870.
Crematogaster rugosa ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 3 (♀).
 SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 348, fig. 17c (♀).
Crematogaster rugosa EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 484 (♀, ♀).
 Type locality: Ogowe, FRENCH CONGO (Mocquérays).
 IVORY COAST: Bassam. CAMEROON. BELGIAN CONGO: Avakubi (J. Bequaert).
 51_a. Var. **nigriventris** SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 87 (♀).
 Type locality: Dimbroko, IVORY COAST (Le Mout).
 51_b. Var. **rugaticeps** SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 87 (♀).
 Type locality: Molundu, CAMEROON (Reichensperger).
 52. **Crematogaster (Sphærocrema) rugosior** (SANTSCHI). See p. 156 (♀).
Crematogaster luctans subsp. *rugosior* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 375 (♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 405 (♀).
 Type locality: Brazzaville, FRENCH CONGO (A. Weiss).
 BELGIAN CONGO: (Kohl); Stanleyville (Lang, Chapin and Bequaert).
 53. **Crematogaster (Sphærocrema) sejuncta** (STITZ).
Crematogaster sejuncta STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 387, fig. 6 (♀).
 Type locality: Koloka near Angu, Belgian Congo (Schubotz).
 54. **Crematogaster (Sphærocrema) similis** (STITZ).
Crematogaster similis STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 388, fig. 7 (♀).

Crematogaster (Sphaerocrema) similis SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.

Type locality: Mt. Karisimbi, north of Lake Kivu, BELGIAN CONGO (Schubotz).

55. *Crematogaster (Sphaerocrema) striatula* EMERY.

Crematogaster striatulus EMERY, 1892, Ann. Soc. Ent. France, LXI, Bull., p. liii (♀); 1899, Ann. Soc. Ent. Belgique, XLIII, p. 482 (♀). H. KOHL, 1909, Natur u. Offenbarung, LV, pp. 103 and 167.

Crematogaster striatula DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 86. SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 374 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 141 (♀). SANTSCHI, 1911, Ann. Soc. Ent. Belgique, LV, p. 282; 1914, Deutsch. Ent. Zeitschr., p. 288 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 342 (♀).

Crematogaster (Sphaerocrema) striatula SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.

Type locality: Assinie, IVORY COAST (C. Alluaud).

SIERRA LEONE: IVORY COAST: (Lohier). GOLD COAST: Aburi (F. Silvestri). CAMEROON: Duala (v. Rothkirch). SPANISH GUINEA: Alen (Tessmann). FRENCH CONGO: Gomba (A. Weiss).

55i. Var. *benitensis* (SANTSCHI).

Crematogaster striatula var. *benitensis* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 374 (♀).

Type locality: River Benito, SPANISH GUINEA (de Brazza).

55i. Var. *obstinata* (SANTSCHI). See p. 157.

Crematogaster striatula var. *obstinata* SANTSCHI, 1911, Rev. Zool. Afr., I, p. 207 (♀).

Type locality: Gomba, FRENCH CONGO (A. Weiss).

BELGIAN CONGO: Leopoldville (J. Bequaert).

56. *Crematogaster (Sphaerocrema) wilwerthi* (SANTSCHI).

Crematogaster wilwerthi SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 372, fig. 6 (♀).

Type locality: Luki, BELGIAN CONGO (Wilwerth).

56i. Var. *confusa* (SANTSCHI).

Crematogaster wilwerthi var. *confusa* SANTSCHI, 1911, Rev. Zool. Afr., I, p. 208 (♀).

Type locality: Gomba, FRENCH CONGO (A. Weiss).

56i. Var. *fauconneti* (FOREL).

Crematogaster wilwerthi var. *fauconneti* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 434 (♀).

Type locality: Nyangwe to Stanleyville, BELGIAN CONGO (Fauconnet).

Subgenus 3. *Orthocrema* SANTSCHI

Crematogaster subg. *Orthocrema* SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.

Subgenotype: *Myrmica sordidula* Nylander, 1849.

57. *Crematogaster (Orthocrema) jeanneli* (SANTSCHI).

Crematogaster jeanneli SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 100 (♀).

Type locality: River Tchania, 1520 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

58. *Crematogaster* (*Orthocrema*) *muralti* (FOREL).*Crematogaster muralti* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 432 (♀).

Type locality: CAMEROON (L. v. Muralt).

58a. Subsp. *livingstonei* SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 344 (♀).*Crematogaster muralti* subsp. *livingstonei* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 486 and 536, fig. 50 (♀).

Type locality: Livingstone, NORTHERN RHODESIA (G. Arnold).

58b. Subsp. *ugandensis* (SANTSCHI).*Crematogaster muralti* subsp. *ugandensis* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 99 (♀).

Type locality: UNYORO PROVINCE, near HOIMA, UGANDA (C. Alluaud).

59. *Crematogaster* (*Orthocrema*) *pauciseta* EMERY.*Crematogaster pauciseta* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 485, fig. (♀, ♀).*Crematogaster pauciseta* FOREL, 1913, Rev. Suisse Zool., XXI, p. 669 (♀, ♀, ♂); 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 340 (♀).*Crematogaster* (*Orthocrema*) *pauciseta* SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.

Type locality: CAMEROON (Conradt).

BELGIAN CONGO: Kilongalanga near St. Gabriel (Kohl).

59a. Subsp. *dolens* (FOREL).*Crematogaster dolens* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 433 (♀).

Type locality: ZANZIBAR (Voeltzkow).

59b. Subsp. *grossulior* (FOREL).*Crematogaster pauciseta* subsp. *grossulior* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 404 (♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

60. *Crematogaster* (*Orthocrema*) *sordidula* (NYLANDER). See p. 1024.*Myrmica sordidula* NYLANDER, 1849, Acta Soc. Sc. Fennicæ, III, 1, p. 44 (♀).*Crematogaster sordidula* MAYR, 1855, Verh. Zool. Bot. Ges. Wien, V, p. 472 (♀, ♀).*Crematogaster sordidula* DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, pp. 336 and 371. ARNOLD, 1920, Ann. South African Mus., XIV, p. 537.*Crematogaster* (*Orthocrema*) *sordidula* SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.

Type locality: not designated.

SOUTHERN EUROPE, NORTH AFRICA, SYRIA, TURKESTAN.

The following records probably refer to some of the forms listed below: TRANSVAAL: Johannesburg (Dixey and Longstaff). CAPE PROVINCE: Nahoon River (Dixey and Longstaff).

60a. Subsp. *natalensis* (FOREL).*Crematogaster sordidula* subsp. *natalensis* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 431 (♀, ♀, ♂); 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 239 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 539 (♀, ♀, ♂).*Crematogaster sordidula* var. *natalensis* "Forel" MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 14 (without description).

Type locality: Mountains of NATAL (R. C. Wroughton).

BASUTOLAND: (R. C. Wroughton). NATAL: Estcourt (R. C. Wroughton);

Krantz Kloof (Marley). ORANGE FREE STATE: Bothaville (H. Brauns).

60a. Var. **braunsi** (FOREL).

Crematogaster sordidula subsp. *natalensis* var. *braunsi* FOREL, 1911, Rev. Zool. Afr., I, p. 277 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 539 (♀).

Crematogaster braunsi "Mayr" FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 432 (♀) (without description; this was never described by Mayr).

Type locality: Cape of Good Hope.

NATAL: in the mountains; Estcourt (R. C. Wroughton).

60b. Subsp. **rectinota** (FOREL).

Crematogaster sordidula subsp. *rectinota* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 126 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 93 (♀, ♀).

Crematogaster sordidula var. *rectinota* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 486 and 537 (♀, ♀), Pl. VI, fig. 73.

Type locality: Bulawayo, RHODESIA (G. Arnold).

BRITISH EAST AFRICA: River Ramisi (Alluaud and Jeannel). GERMAN EAST AFRICA: Tanga (Alluaud and Jeannel).

61. **Crematogaster (Orthocrema) transvaalensis** (FOREL).

Crematogaster transvaalensis FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 96 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 486 and 533 (♀).

Crematogaster transvaalensis EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 30 (♀), Pl. II, fig. 18.

Type locality: TRANSVAAL (P. Berthoud).

CAPE PROVINCE: Cape Town; Vrijburg (E. Simon).

61a. Var. **hammi** (ARNOLD).

Crematogaster transvaalensis var. *hammi* ARNOLD, 1920, Ann. South African Mus., XIV, p. 534, fig. 48 (♀).

Type locality: not indicated; probably in RHODESIA (G. Arnold).

Subgenus 4. **Atopogyne** FOREL

Crematogaster subg. *Atopogyne* FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 343.

Crematogaster subg. *Atopogynes* SANTSCHI, 1918, Bull. Soc. Ent. France, p. 183.

Subgenotype: *Formica depressa* Latreille, 1802.

62. **Crematogaster (Atopogyne) africana** (MAYR).

Crematogaster africana MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 142 (♀); 1902, Verh. Zool. Bot. Ges. Wien, LII, p. 294 (♀). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, pp. 53 and 69 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 378. FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 433. WASMANN, 1918, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VII, p. 145.

Crematogaster africana EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 480, fig. (♀). H. KOHL, 1909, Natur u. Offenbarung, LV, pp. 164 and 168.

Crematogaster buchneri subsp. *africana* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 352 (♀).

Crematogaster (Atopogynes) africana SANTSCHI, 1918, Bull. Soc. Ent. France, p. 183.

Type locality: Warship harbor in CAMEROON (H. Brauns).

GOLD COAST: Aburi (R. Buchholz). NIGERIA: Old Calabar. CAMEROON: Duala (v. Rothkirch). FRENCH CONGO: Brazzaville; Mindouli (A. Weiss). BELGIAN CONGO: (Kohl); Mayombe (Cabra; de Briey); Kondué (Luja).

62a. Var. **biemarginata** (FOREL).

Cremastogaster africana var. *biemarginata* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 433 (♀).

Type locality: CAMEROON.

62a. Var. **camena** (FOREL).

Cremastogaster buchneri subsp. *africana* var. *camena* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 410 (♀).

Type locality: Olombo, BELGIAN CONGO (Kohl).

62a. Var. **schumanni** (MAYR). See p. 157.

Cremastogaster buchneri subsp. *schumanni* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 144 (♀).

Type locality: CAMEROON (C. Schumann).

BELGIAN CONGO: Leopoldville (J. Bequaert).

62a. Var. **stanleyi** (SANTSCHI).

Cremastogaster buchneri subsp. *africana* var. *stanleyi* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 253 (♀).

Cremastogaster africana var. *stanleyi* FOREL, 1911, Rev. Zool. Afr., I, p. 278 (♀; without description).

Type locality: Mindouli, FRENCH CONGO (A. Weiss).

BELGIAN CONGO: Kondué (Luja).

62a. Var. **variegata** (MAYR).

Cremastogaster africana var. *variegata* MAYR, 1902, Verh. Zool. Bot. Ges. Wien, LII, p. 294 (♀).

Type locality: Victoria, CAMEROON (R. Buchholz).

62a. Subsp. **alligatrix** (FOREL).

Cremastogaster buchneri subsp. *alligatrix* FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 271 (♀); 1911, Rev. Zool. Afr., I, p. 278 (♀); 1913, *ibid.*, II, p. 352 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 351 (♀); 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 338 (♀). LAMBORN, 1914, Trans. Ent. Soc. London, (1913), p. 442 (♀). FARQUHARSON, 1919, *ibid.*, (1918), Proc., p. xl.

Type locality: Old Calabar, NIGERIA (Bates).

SOUTHERN NIGERIA: Oni Camp, east of Lagos (Lamborn). BELGIAN CONGO: St. Gabriel (Kohl); Kondué (Luja); Mayombe (De Briey).

62b. Subsp. **laurenti** (FOREL) SANTSCHI *in litt.* See p. 157.

Cremastogaster africana subsp. *laurenti* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 69 (♀).

Cremastogaster africana subsp. *laurenti* H. KOHL, 1909, Natur u. Offenbarung, LV, pp. 160, 164, and 168.

Cremastogaster buchneri subsp. *laurenti* FOREL, 1913, Rev. Zool. Afr., II, p. 352 (♀). LAMBORN, 1914, Trans. Ent. Soc. London, (1913), p. 442 (♀).

Type locality: Bokala, BELGIAN CONGO (Laurent).

SOUTHERN NIGERIA: Oni Camp east of Lagos (Lamborn). BELGIAN CONGO: Isangi; Stanleyville (Laurent); Tshopo River near Stanleyville (J. Bequaert).

62b. Var. **zeta** (FOREL). See p. 157.

Cremastogaster africana subsp. *laurenti* var. *zeta* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 70 (♀).

Type locality: BELGIAN CONGO (Laurent; Kohl).

BELGIAN CONGO: Pale (Niembo) between Walikale and Lubutu; Leopoldville (J. Bequaert); Stanleyville (Lang and Chapin).

62c. Subsp. *tibialis* SANTSCHI. See p. 157 (♀).

Type locality: Mosekowa between Walikale and Lubutu, BELGIAN CONGO (J. Bequaert).

62d. Subsp. *winkleri* (FOREL).

Crematogaster africana var. *winkleri* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 69 (♀); 1911, Sitzb. Bayer. Akad. Wiss., p. 272 (♀); 1911, Rev. Zool. Afr., I, p. 278 (♀). AULMANN and LA BAUME, 1912, 'Fauna d. Deutsch. Kolon.', V, 3, p. 63, fig. 36 (nest). AULMANN, 1913, *ibid.*, V, 5, p. 93, fig. 79 (nest).

Crematogaster buchneri subsp. *winkleri* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 124 (♀, ♀); 1913, Rev. Zool. Afr., II, pp. 324 and 352 (♀); 1913, Rev. Suisse Zool., XXI, p. 670. LAMBORN, 1914, Trans. Ent. Soc. London, (1913), p. 442 (♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 409 (♀).

Crematogaster africana var. *winkleri* L. REH, 1905, Zeitschr. f. Pflanzenkrankh., XV, p. 134 (without description).

Type locality: Victoria, CAMEROON (H. Winkler).

NIGERIA: Oni Camp east of Lagos (Lamborn); Old Calabar (Bates). CAMEROON: Johann-Albrechtshöhe (Conradt). BELGIAN CONGO: Olombo (Kohl); Eala; Bokala (Laurent); Kondué (Luja). RHODESIA: (G. Arnold).

62d. Var. *brieyi* (FOREL).

Crematogaster buchneri subsp. *winkleri* var. *brieyi* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 352 (♀).

Type locality: Mayombe, BELGIAN CONGO (de Briey).

62d. Var. *fickendeyi* (FOREL). See p. 158.

Crematogaster buchneri subsp. *winkleri* var. *fickendeyi* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, pp. 125 and 351 (♀); 1913, Rev. Zool. Afr., II, p. 324; 1916, Rev. Suisse Zool., XXIV, p. 409 (♀).

Type locality: Victoria, CAMEROON (Fickendey).

BELGIAN CONGO: Kondué (Luja); Leopoldville; St. Gabriel (Kohl); Masongo between Walikale and Lubutu (J. Bequaert).

62d. Var. *transversiruginota* (FOREL).

Crematogaster buchneri subsp. *winkleri* var. *transversiruginota* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 410 (♀).

Type locality: Motombe, Okiavo River, BELGIAN CONGO (Kohl).

63. *Crematogaster* (*Atopogyne*) *angusticeps* (SANTSCHI).

Crematogaster angusticeps SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 354, fig. (♀).

Type locality: Sikasso, FRENCH SUDAN (Chevalier).

64. *Crematogaster* (*Atopogyne*) *batesi* (FOREL).

Crematogaster batesi FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 270 (♀).

Type locality: Old Calabar, NIGERIA (Bates).

65. *Crematogaster* (*Atopogyne*) *buchneri* (FOREL).

Crematogaster buchneri FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 94 (♀). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 138 (♀). FOREL, 1913, Rev. Suisse Zool., XXI, p. 669 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 487 and 520 (♀).

Crematogaster buchneri EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 479, fig. (♂, ♀).

Crematogaster (Atopogynes) buchneri SANTSCHI, 1918, Bull. Soc. Ent. France, p. 183.

Type locality: BENGUELA (M. Buchner).

CAMEROON: (Conradt). ANGOLA.

651. Var. *biimpressa* (MAYR).

Crematogaster buchneri subsp. *biimpressa* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 139 (♂).

Type locality: Kuilu River, FRENCH CONGO.

652. Var. *grateri* (FOREL).

Crematogaster buchneri var. *grateri* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 409 (♂).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

653a. Subsp. *clariventris* (MAYR).

Crematogaster buchneri subsp. *clariventris* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 139 (♂); 1896, Ent. Tidskr., XVII, p. 242 (♂). FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 271 (♀); 1912, Mitt. Naturh. Mus. Hamburg, XXIX, p. 180; 1913, Ann. Soc. Ent. Belgique, LVII, p. 351 (♂); 1913, Rev. Zool. Afr., II, p. 352 (♂). LAMBORN, 1914, Trans. Ent. Soc. London, (1913), p. 442 (♂).

Type locality: Loango Coast, FRENCH CONGO (H. Brauns).

LIBERIA: (Scherer). SOUTHERN NIGERIA: Oni Camp east of Lagos (Lamborn). CAMEROON: (Sjöstedt). BELGIAN CONGO: Kondué (Luja). UGANDA: Entebbe (Schultze).

66. *Crematogaster (Atopogynes) bulawayensis* (FOREL).

Crematogaster buchneri subsp. *africana* var. *bulawayensis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 125 (♂).

Crematogaster gallicola var. *oraculum* FOREL, 1913, Rev. Zool. Afr., II, p. 323 (♂). BEQUAERT, *ibid.*, p. 425. FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 124 (♂); 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 237 (♂). (Synonym of *bulawayensis*, *teste* Arnold).

Crematogaster bulawayensis ARNOLD, 1920, Ann. South African Mus., XIV, p. 521, fig. 42 (♂), Pl. vi, figs. 74 and 74a.

Type locality: Bulawayo, RHODESIA (G. Arnold).

BELGIAN CONGO: Sankisia (J. Bequaert).

661. Var. *desperans* (FOREL).

Crematogaster gabonensis var. *desperans* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 233 (♂).

Crematogaster desperans SANTSCHI, 1918, Bull. Soc. Ent. France, p. 185.

Crematogaster bulawayensis var. *desperans* ARNOLD, 1920, Ann. South African Mus., XIV, p. 523, fig. 44 (♂).

Type locality: Durban, NATAL (G. Arnold).

662. Var. *rhodesiana* (FOREL).

Crematogaster inermis subsp. *delagoensis* var. *rhodesiana* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 126 (♂).

Crematogaster bulawayensis var. *rhodesiana* ARNOLD, 1920, Ann. South African Mus., XIV, p. 523, fig. 43 (♂), Pl. vi, figs. 82 and 82a.

Type locality: Bulawayo, RHODESIA (G. Arnold).

66a. Var. **zulu** (SANTSCHI).

Cremastogaster sjostedti subsp. *bulawayensis* var. *zulu* SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 15 (♀, ♀).

Type locality: Mfongosi, ZULULAND (W. E. Tones).

66a. Subsp. **infaceta** SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 498 (♀).

Cremastogaster vulcanica var. *FOREL*, 1913, Ann. Soc. Ent. Belgique, LVII, p. 124 (♀).

Cremastogaster vulcania subsp. *godefreyi* var. *foraminicipoides* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 406 (♀).

Cremastogaster sjostedti subsp. *infaceta* SANTSCHI, 1918, Bull. Soc. Ent. France, p. 184.

Cremastogaster godefreyi var. *foraminicipoides* ARNOLD, 1920, Ann. South African Mus., XIV, p. 505 (♀).

Cremastogaster bulawayensis subsp. *infaceta* ARNOLD, *ibid.*, 1920, XIV, p. 524 (♀).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

RHODESIA: Bulawayo (G. Arnold).

66a. Var. **pudica** SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 499 (♀).

Cremastogaster bulawayensis subsp. *infaceta* var. *pudica* ARNOLD, 1920, Ann. South African Mus., XIV, p. 525 (♀).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

67. **Crematogaster (Atopogyne) depressa** (LATREILLE).

Formica depressa LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 268 (♀), Pl. XI, fig. 73. ROGER, 1863, 'Verzeich. Formicid.,' p. 33. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 414.

Formica platygnatha ROGER, 1863, Berlin. Ent. Zeitschr., VII, p. 168 (♀); 1863, 'Verzeich. Formicid.,' p. 15. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 419.

Cremastogaster mandibularis ERN. ANDRÉ, 1889, Rev. d'Ent. Caen, VIII, p. 229 (♀).

Cremastogaster platygnatha ERN. ANDRÉ, 1890, *ibid.*, IX, p. 323 (♀).

Crematogaster depressa EMERY, 1892, Bull. Soc. Ent. Italiana, XXIII, p. 164; 1899, Ann. Soc. Ent. Belgique, XLIII, p. 478, fig. (♀).

Cremastogaster depressa DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 81. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 144. FOREL, in Schultze, 1910, 'Forchungsreise in Südafrika,' IV, p. 5 (♀, ♂).

Cremastogaster buchneri subsp. *foreli* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 138 (♀). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 53 (♀).

Cremastogaster (Atopogynes) depressa SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 351 (♀); 1918, Bull. Soc. Ent. France, p. 183.

Cremastogaster (Atopogynes) depressa subsp. *foreli* SANTSCHI, 1914, Boll. Lab. Gen. Agrar. Portici, VIII, p. 351 (♀).

Type locality: Coast of GUINEA (Palisot de Beauvois).

WEST AFRICA: (Fülleborn). SENEGAMBIA: LOS ISLANDS. SIERRA LEONE: (Mocquerys). GOLD COAST. SLAVE COAST. SOUTHERN NIGERIA: Lagos; Olokemeji; Ibadan (F. Silvestri); Old Calabar (H. Brauns). CAMEROON: Victoria. BELGIAN CONGO: Luki (A. Jullien).

671. **Var. adultera** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 252 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

672. **Var. fuscipennis** EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 479 (♀). See p. 159.

Crematogaster depressa var. *fuscipennis* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 380, figs. 8b-c and 9a-c (♀, ♀, ♀ ergatoid.). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 53 (♀, ♂); 1911, Rev. Zool. Afr., I, p. 279 (♀); 1913, ibid., II, p. 325 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 353 (♀).

Type locality: CAMEROON.

FRENCH CONGO: Gomba region (A. Weiss). BELGIAN CONGO: Kinshasa; Boyengue; Ikelemba (Waelbroeck); Leopoldville (Dubois; J. Bequaert); Sankisia (J. Bequaert); Mayombe (Deleval); Stanleyville; Medje; Niapu; Ambelokudi; Niangara (Lang and Chapin).

68. **Crematogaster (Atopogyne) gabonensis** EMERY.

- Crematogaster gabonensis* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 484 (♀).

Crematogaster gabonensis SANTSCHI, 1911, ibid., LV, p. 281 (♂). FOREL, 1913, ibid., LVII, p. 126 (♀).

Type locality: Gaboon, FRENCH CONGO.

IVORY COAST: (Lohier). CAMEROON: Victoria (Fickendey).

681. **Var. fuscitatis** (FOREL).

Crematogaster gabonensis var. *fuscitatis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 351 (♀).

Type locality: Kondué, BELGIAN CONGO (Luja).

BELGIAN CONGO: Congo da Lemba (R. Mayné).

69. **Crematogaster (Atopogyne) homeri** (FOREL).

Crematogaster (Atopogyne) homeri FOREL, 1913, Rev. Suisse Zool., XXI, p. 668 (♀).

Type locality: Johann-Albrechtshöhe, CAMEROON (Conradt).

70. **Crematogaster (Atopogyne) jullieni** (SANTSCHI).

Crematogaster jullieni SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 378, figs. 8a and 8d (♀, ♀). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 5 (♀).

Crematogaster (Atopogyne) jullieni FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 343.

Crematogaster (Atopogynes) jullieni SANTSCHI, 1918, Bull. Soc. Ent. France, p. 183.

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

WEST AFRICA: (Fülleborn). FRENCH CONGO: Gaboon.

71. **Crematogaster (Atopogyne) kasalensis** (FOREL).

Crematogaster kasalensis FOREL, 1913, Rev. Zool. Afr., II, p. 321 (♀, ♀); 1916, Rev. Suisse Zool., XXIV, p. 407 (♀, ♀).

Type locality: Kondué, BELGIAN CONGO (Luja).

BELGIAN CONGO: St. Gabriel (Kohl).

72. **Crematogaster (Atopogyne) kohli** (FOREL).

Crematogaster kohli FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, pp. 70 and 71 (♀); 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 340 (♀, ♀).

Type locality: Stanleyville, BELGIAN CONGO (Kohl).

BELGIAN CONGO: Romée near Stanleyville (Luja); St. Gabriel (Kohl).

73. *Crematogaster (Atopogyne) theta* (FOREL) SANTSCHI *in litt.* See p. 159.

Crematogaster africana var. *theta* FOREL, 1911, Rev. Zool. Afr., I, p. 278 (♀).

Crematogaster buchneri subsp. *laurenti* var. *theta* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, pp. 125 and 352 (♂); 1913, Rev. Zool. Afr., II, p. 325; 1916, Rev. Suisse Zool., XXIV, p. 410 (♀, ♀).

Type locality: Kondué, BELGIAN CONGO (Luja).

BELGIAN CONGO: St. Gabriel (Kohl); Medje; Avakubi; Stanleyville (Lang and Chapin).

74. *Crematogaster (Atopogyne) transiens* (FOREL) SANTSCHI *in litt.* See p. 160.

Crematogaster buchneri subsp. *transiens* FOREL, 1913, Rev. Zool. Afr., II, p. 324 (♀). BEQUAERT, *ibid.*, p. 426. SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 253.

Type locality: Kaba near Kikondja, BELGIAN CONGO (J. Bequaert).

BELGIAN CONGO: Kataké (Gérard); Avakubi; Stanleyville (Lang and Chapin).

75. *Crematogaster (Atopogyne) wasmanni* (SANTSCHI).

Crematogaster wasmanni SANTSCHI, 1910, Ann. Soc. Ent. France, LXXXVIII, (1909), p. 376, fig. 7 (♀).

Crematogaster (Atopogynes) wasmanni SANTSCHI, 1918, Bull. Soc. Ent. France, p. 183.

Type locality: Sankuru, BELGIAN CONGO (Luja).

76. *Crematogaster (Atopogyne) wellmani* (FOREL).

Crematogaster wellmani FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 64 (♂; nec ♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 376 (♀). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 433 (♀).

Crematogaster (Atopogynes?) wellmani SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 500 (♀).

Type locality: BENGUELA (C. Wellman).

NIGERIA: Old Calabar. FRENCH CONGO: Brazzaville (A. Weiss); Libreville.

76a. Var. *luciae* (FOREL).

Crematogaster concava EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 480 (♀) (nec ♀; nec *concava* André).

Crematogaster (Atopogyne) luciae FOREL, 1913, Rev. Zool. Afr., II, p. 325 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 353 (♀); 1913, Rev. Suisse Zool., XXI, p. 668 (♀); 1916, *ibid.*, XXIV, p. 411 (♀).

Crematogaster (Atopogynes?) wellmani var. *luciae* SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 500 (♀).

Type locality: SIERRA LEONE.

CAMEROON: (Conradt). BELGIAN CONGO: (Kohl); Kwesi to Kilo (Bayer).

76b. Var. *weissi* SANTSCHI.

Crematogaster wellmani var. *weissi* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXXVIII, (1909), p. 376 (♀).

Crematogaster wellmani var. *weissi* SANTSCHI, 1916, *ibid.*, LXXXIV, (1915), p. 500 (♀).

Type locality: Gomba, FRENCH CONGO (A. Weiss).

76a. Subsp. **retusa** SANTSCHI.

Crematogaster welmani subsp. *retusa* SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 500 (♀, ♂).

Crematogaster (*Atopogyne*) *welmani* subsp. *retusa* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 486 and 545, fig. 54 (♀, ♂).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

Subgenus 5. **Oxygyne** FOREL

Crematogaster subg. *Oxygyne* FOREL, 1901, Ann. Soc. Ent. Belgique, XLV, p. 376.

Crematogaster subg. *Oxygyne* SANTSCHI, 1918, Bull. Soc. Ent. France, p. 183.

Subgenotype: *Crematogaster* (*Oxygyne*) *daisyi* FOREL, 1901.

77. ***Crematogaster* (*Oxygyne*) *magitæ*** (FOREL).

Crematogaster (*Oxygyne*?) *magitæ* FOREL, in Schultz, 1910, 'Forschungsreise in Südafrika,' IV, p. 9 (♀); 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 369.

Type locality: West Africa (Fülleborn).

78. ***Crematogaster* (*Oxygyne*) *margaritæ*** EMERY.

Crematogaster margaritæ EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 31 (♀).

Crematogaster margaritæ MAYR, 1896, Ent. Tidskr., XVII, p. 243 (♀, ♂).

AURIVILLIUS, *ibid.*, p. 254, Pl. IV, fig. 2 (nest). SJÖSTEDT, 1904, 'I Västafriskas Urskogar,' p. 504, fig. 2 (nest).

Crematogaster (*Oxygyne*) *margaritæ* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 381, fig. 8c (♀).

Type locality: Kniulu, FRENCH CONGO.

CAMEROON: (Sjöstedt). FRENCH CONGO: Brazzaville (A. Weiss).

78a. Subsp. **lujae** (FOREL).

Crematogaster (*Oxygyne*) *margaritæ* subsp. *lujae* FOREL, 1913, Rev. Zool. Afr., II, p. 325 (♀).

Type locality: Kondué, BELGIAN CONGO (Luja).

79. ***Crematogaster* (*Oxygyne*) *oscaris*** (FOREL).

Crematogaster (*Oxygyne*) *oscaris* FOREL, in Schultz, 1910, 'Forschungsreise in Südafrika,' IV, p. 7 (♀, ♂), Pl. I, figs. 5 and 6. ARNOLD, 1920, Ann. South African Mus., XIV, p. 543 (♀, ♂).

Type locality: Kamaggas, CAPE PROVINCE (L. Schultz).

80. ***Crematogaster* (*Oxygyne*) *santschii*** (FOREL).

Crematogaster santschii, FOREL, 1913, Rev. Zool. Afr., II, p. 322 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 352 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 487 (♀).

Type locality: Kondué, BELGIAN CONGO (Luja).

BELGIAN CONGO: Congo da Lemba (R. Mayné).

80₁. Var. **brevarmata** (FOREL).

Crematogaster santschii var. *brevarmata* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 337 (♀, ♂).

Type locality: Kondué, BELGIAN CONGO (Luja).

80₂. Var. **clymene** (FOREL).

Crematogaster santschii var. *clymene* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 337 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 540, fig. 52 (♀).

Type locality: Durban, NATAL (C. B. Cooper).

81. **Crematogaster (Oxygyne) trautweini** (VIEHMEYER).
Crematogaster (Oxygyne) trautweini VIEHMEYER, 1914, Arch. f. Naturg., LXXIX, Abt. A, Heft 12, (1913), p. 33 (♀).
 Type locality: CAMEROON (H. Trautwein).

Subgenus 6. **Nematocrema** SANTSCHI

- Crematogaster* subg. *Nematocrema* SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.
 Subgenotype: *Crematogaster stadelmanni* MAYR, 1895.
 82. **Crematogaster (Nematocrema) breviventris** (SANTSCHI).
Crematogaster (Nematocrema) breviventris SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 379 (♀).

Type locality: Molundu, CAMEROON (Reichensperger).

83. **Crematogaster (Nematocrema) stadelmanni** (MAYR). See p. 160.
Crematogaster stadelmanni MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 140 (♀, ♀); 1902, Verh. Zool. Bot. Ges. Wien, LII, p. 294 (♀). DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, p. 380. FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 369 (♀).

Crematogaster (Oxygyne) stadelmanni FOREL, 1911, Rev. Zool. Afr., I, p. 277 (♀); 1913, Rev. Suisse Zool., XXI, p. 670 (♀); 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 337 (♀); 1916, Rev. Suisse Zool., XXIV, p. 411 (♀).

Crematogaster (Oxygyne) stadelmanni SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 349 (♀).

Crematogaster (Nematocrema) stadelmanni SANTSCHI, 1918, Bull. Soc. Ent. France, p. 182.

Type locality: LIBERIA.

FRENCH GUINEA: Konakry (F. Silvestri). CAMEROON: (Conradt); Victoria (R. Buchholz). BELGIAN CONGO: Congo da Lemba (Mayné); St. Gabriel (Kohl); Stanleyville (Lang and Chapin). BRITISH EAST AFRICA: Victoria Nyanza (Zimmer). CAPE PROVINCE: Simon's Town (E. Simon).

83₁. Var. **anguliceps** (STITZ).

Crematogaster (Oxygyne) stadelmanni var. *anguliceps* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 384, fig. 4b (♀).

Type locality: Ossidinge, CAMEROON.

83₂. Var. **angustata** (MAYR).

Crematogaster stadelmanni var. *angustata* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 142 (♀).

Type locality: CAMEROON (H. Brauns).

83₂. Var. **dolichocephala** (SANTSCHI). See p. 160.

Crematogaster stadelmanni var. *dolichocephala* SANTSCHI, 1911, Rev. Zool. Afr., I, p. 208, with description (♀, ♀, ♂); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 349, with description (♀). FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 337 (♀).

Crematogaster stadelmanni subsp. *dolichocephala* "Emery in litt." SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 378 (without description). WASMANN, 1911, 1^{er} Congr. Intern. Entom. Bruxelles (1910), II, Mém., p. 231, Pl. XVI, fig. 32 (nest); 1913, Ann. Rept. Smiths. Inst. for 1912, p. 472, Pl. x, fig. 32 (nest).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

NIGERIA: Ibadan (F. Silvestri). FRENCH CONGO: Mindouli; Gomba (A. Weiss). BELGIAN CONGO: St. Gabriel (Kohl); Bengamisa; Manamana; Kwamouth; Ngayu (Lang and Chapin); Kondué (Luja).

83a. Var. **intermedia** (MAYR).

Cremastogaster stadelmanni var. *intermedia* MAYR, 1896, Ent. Tidskr., XVII, p. 242 (♂, ♀). AURIVILLIUS, *ibid.*, p. 254, Pl. v (nest). SJÖSTEDT, 1904, 'I Västafrikas Urskogar,' pp. 502 and 503, fig. (nest).

Type locality: CAMEROON (Sjöstedt).

83b. Var. **ovinodis** (STITZ).

Cremastogaster (Oxygyné) stadelmanni var. *ovinodis* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 383, fig. 4d (♂).

Type locality: Duma, BELGIAN CONGO (Schubotz).

83c. Var. **schereri** (FOREL).

Cremastogaster stadelmanni var. *schereri* FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 273 (♂).

Type locality: Bendov, LIBERIA (Scherer).

Subgenus 7. **Decacrema** FOREL

Cremastogaster subg. *Decacrema* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 18; in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 9.

Subgenotype: *Cremastogaster (Decacrema) decamera* Forel, 1910.

84. **Crematogaster (Decacrema) arthuri-mülleri** (FOREL).

Cremastogaster gallicola subsp. *arthuri-mülleri* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 96 (♂).

Cremastogaster arthuri-mülleri MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 144 (♂).

Cremastogaster (Decacrema) arthur-mülleri ARNOLD, 1920, Ann. South African Mus., XIV, p. 547 (♂).

Type locality: DELAGOA (Arthur Müller).

East Africa.

85. **Crematogaster (Decacrema) edentula** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 253 (♀).

Cremastogaster (Decacrema) edentula SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 349, figs. 18a-b (♂, ♀).

Type locality: Kindia, FRENCH GUINEA (F. Silvestri).

DAHOMY: Kouandé (Desanti).

86. **Crematogaster (Decacrema) liengmei** (FOREL).

Cremastogaster gallicola var. *liengmei* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 96 (♂).

Cremastogaster liengmei WASMANN, 1898, Wien. Ent. Zeitg., XVII, p. 97.

Cremastogaster (Decacrema) liengmei FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 436 (♂). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 21 (♂).

Cremastogaster liengmei REICHENSFELGER, 1915, Med. Göteborgs Mus. Zool. Afd., V, p. 6.

Cremastogaster (Decacrema) liengmei ARNOLD, 1920, Ann. South African Mus., XIV, p. 548 (♂).

Type locality: DELAGOA (Liengme).

NATAL: (L. v. Muralt); Balgowan (I. Trägårdh).

86a. Subsp. **caiculata** (FOREL).

Crematogaster peringueyi subsp. *caiculata* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 341 (♀).

Crematogaster liengmei subsp. *caiculata* ARNOLD, 1920, Ann. South African Mus., XIV, p. 550 (♀).

Type locality: Durban, NATAL (C. B. Cooper).

86b. Subsp. **weitzackeri** (EMERY).

Crematogaster weitzackeri EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 28 (♀), Pl. II, figs. 14 and 15.

Crematogaster (*Decacrema*) *liengmei* var. *weitzackeri* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 21 (♀).

Crematogaster weitzackeri DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, p. 371.

Crematogaster excisa subsp. *andrei* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 236 (♀, according to Arnold, with 10-jointed antennæ).

Crematogaster caelestis var. *kloofensis* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 237 (♀, teste Arnold).

Crematogaster (*Decacrema*) *liengmei* subsp. *weitzackeri* ARNOLD, 1920, Ann. South African Mus., XIV, p. 548, fig. 55 (♀).

Type locality: Pietermaritzburg, NATAL (Weitzacker; I. Trägårdh).

NATAL: Estecourt (R. C. Wroughton); Krantz Kloof, Durban (Marley). CAPE PROVINCE: Cape Town (E. Simon); Nahoon River (Dixey and Longstaff).

86b. Var. **acanthobia** (FOREL).

Crematogaster excisa subsp. *andrei* var. *acanthobia* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 236 (♀).

Crematogaster excisa subsp. *andrei* var. *pensitata* FOREL, 1915, *ibid.*, (5) L, p. 341 (♀).

Crematogaster liengmei subsp. *weitzackeri* var. *acanthobia* ARNOLD, 1920, Ann. South African Mus., XIV, p. 550 (♀).

Type locality: Willowmore, CAPE COLONY (Arnold).

86b. Var. **gordonensis** (FOREL).

Crematogaster excisa subsp. *andrei* var. *gordonensis* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 236 (♀).

Crematogaster liengmei subsp. *weitzackeri* var. *gordonensis* ARNOLD, 1920, Ann. South African Mus., XIV, p. 549 (♀).

Type locality: Gordon Bay, CAPE PROVINCE.

86b. Var. **thais** (FOREL).

Crematogaster excisa subsp. *andrei* var. *thais* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 126 (♀).

Crematogaster liengmei subsp. *weitzackeri* var. *thais* ARNOLD, 1920, Ann. South African Mus., XIV, p. 549 (♀).

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

CAPE PROVINCE: Grahamstown (Sherry).

87. ***Crematogaster* (*Decacrema*) *petiolidens*** (FOREL).

Crematogaster (*Decacrema*) *petiolidens* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 412 (♀).

Type locality: BELGIAN CONGO (Kohl).

88. *Crematogaster* (*Decacrema*) *solenopsides* EMERY.

Crematogaster solenopsides EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 487 (♀).

Type locality: ZANZIBAR.

88a. Subsp. *costeboriensis* (SANTSCHI).

Crematogaster (*Decacrema*) *solenopsoides* subsp. *costeboriensis* SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 86 (♀).

Type locality: Region of San Pedro, IVORY COAST (C. Phore).

88b. Subsp. *flavida* (MAYR).

Crematogaster solenopsides subsp. *flavida* MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.', II, 8, p. 15 (♀).

Crematogaster solenopsides var. *flavida* SJÖSTEDT, 1908, *ibid.*, II, 8, p. 117.

Type locality: Same to Moembe, Usambara, GERMAN EAST AFRICA (Sjöstedt).

88b₁. Var. *convexiclypea* (FOREL).

Crematogaster (*Decacrema*) *solenopsoides* subsp. *flavida* var. *convexiclypea* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 411 (♀, ♀, ♂).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

88b₂. Var. *flaviscapis* (SANTSCHI).

Crematogaster solenopsoides subsp. *flavida* var. *flaviscapis* SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 85 (♀).

Type locality: Landana, PORTUGUESE CONGO (J. Bequaert).

88b₃. Var. *gallarum* (SANTSCHI).

Crematogaster solenopsides subsp. *flavida* var. *gallarum* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 370 (♀).

Crematogaster solenopsoides subsp. *flavida* var. *gallarum* SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 86.

Type locality: Mindouli, FRENCH CONGO (A. Weiss).

FRENCH CONGO: Mbamu (A. Weiss).

Solenopsidini Emery**Monomorium MAYR**

Monomorium MAYR, 1855, Verh. Zool. Bot. Ges. Wien, V, p. 452.

Atta (part) JERDON. *Formica* (part) LINNÆUS, et auct. *Myrmica* (part) SMITH, et auct.

Genotype: *Monomorium minutum* Mayr, 1855.

Subgenus I. Monomorium MAYR, sensu stricto

Subgenotype: same as genotype.

1. *Monomorium altinode* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 18.

Monomorium rhopalocerum var. *altinodis* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 359, fig. 4b (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

1₁. Var. *bondroiti* SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 10, fig. 1f (♀).

Type locality: Upper Lukuga River, BELGIAN CONGO (Gérard).

2. **Monomorium angustinode** FOREL, 1913, Rev. Zool. Afr., II, p. 334 (♀).
BEQUAERT, *ibid.*, p. 428.
Type locality: Welgelegen, BELGIAN CONGO (J. Bequaert).
3. **Monomorium arnoldi** FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 137 (♀).
ARNOLD, 1916, Ann. South African Mus., XIV, p. 232 (♀).
Type locality: Matoppo Hills, SOUTHERN RHODESIA (G. Arnold).
4. **Monomorium braunsi** MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 7 (♀).
ARNOLD, 1916, Ann. South African Mus., XIV, p. 234 (♀).
Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).
4. **Var. shilohense** FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 217 (♀).
ARNOLD, 1916, Ann. South African Mus., XIV, p. 235 (♀).
Type locality: Shiloh, SOUTHERN RHODESIA (G. Arnold).
5. **Monomorium egeus** FOREL, 1910, Ann. Soc. Ent. Belgique, LXV, p. 443 (♀).
Type locality: CAMEROON (v. Muralt).
6. **Monomorium emeryi** MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 132 (♀).
FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 136 (♀).
ARNOLD, 1916, Ann. South African Mus., XIV, p. 212 (♀, ♀, ♂), Pl. v, fig. 62.
Type locality: Mozambique Island, PORTUGUESE EAST AFRICA (H. Brauns).
RHODESIA: Redbank; Nyamandhloru (G. Arnold).
7. **Monomorium explorator** SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 12, fig. 1a-b (♀).
Type locality: Samkita, FRENCH CONGO (F. Faure).
8. **Monomorium fasciatum** SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 10, fig. 1c-e (♀).
Type locality: Kilimanjaro, GERMAN EAST AFRICA (Reichensperger).
9. **Monomorium floricola** (JERDON) EMERY, in Dalla Torre, 1893, 'Cat. Hym.', VII, p. 66.
SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 354 (♀, ♀). See p. 1026.
Atta floricola JERDON, 1851, Madras Journ. Litt. Sc., XVII, p. 107; 1854, Ann. Mag. Nat. Hist., (2) XIII, p. 49 (♀).
Monomorium specularis MAYR, 1866, Sitzb. Ak. Wiss. Wien, LIII, Abt. 1, p. 509 (♀); 1878, Verh. Zool. Bot. Ges. Wien, XXVIII, p. 671.
Type locality: Tellicherry, Southern INDIA (Jerdon).
Tropicopolitan. SOUTHERN NIGERIA: Lagos (F. Silvestri).
10. **Monomorium hanneli** FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 18 (♀).
Type locality: Mte-ya-Kifaru, GERMAN EAST AFRICA (Katona).
11. **Monomorium ilgi** FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 84 (♀).
Type locality: Southern ABYSSINIA (Ilg).
12. **Monomorium leimbachi** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 246 (♀).
ARNOLD, 1916, Ann. South African Mus., XIV, p. 233 (♀).
Type locality: Cape Town, CAPE PROVINCE (G. Arnold).
13. **Monomorium lene** SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 11, fig. 2g-h (♀).
Type locality: Salisbury, SOUTHERN RHODESIA (G. Arnold).
14. **Monomorium mediocre** SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 376, fig. 13 (♀).
Type locality: Kimberley, CAPE PROVINCE (G. Arnold).

15. **Monomorium minutum** MAYR, 1855, Verh. Zool. Bot. Ges. Wien, V, p. 453 (♀); 1861, 'Europ. Formicid.', p. 72 (♀); 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 753 (♀). ROGER, 1863, 'Verzeich. Formicid.', p. 31. MAYR, 1865, 'Reise Novara Zool.', II, 1, Formicidæ, p. 91 (♀, ♂); 1866, Sitzb. Ak. Wiss. Wien, LIII, Abt. I, p. 509 (♀). ERN. ANDRÉ, 1874, Rev. Mag. Zool., (3) II, p. 199; 1883, 'Spec. Hym. Europ. Algérie,' II, pp. 333, 338, and 341 (♀, ♀, ♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 67. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 72 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 217 (♀). See p. 1027.

Myrmica (*Monomarium*) *minuta* F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 129 (♀).

Monomorium carbonarium EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 532 (♀).

Type locality: Lombardy, ITALY (Villa).

UGANDA: Region of Lake Albert (C. Alluaud).

15. Var. **leopoldinum** FOREL, 1905, Ann. Soc. Ent. Belgique, XLIX, p. 179 (♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

15. Var. **pallidipes** FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 252 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 72 (♀).

Type locality: Nefasit, ERITREA (K. Escherich).

BRITISH EAST AFRICA: Kikuyu, Nairobi, 1700 m. (Alluaud and Jeannel).

15a. Subsp. **boerorum** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 244 (♀, ♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 218 (♀, ♀).

Monomorium minutum var. *boerorum* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 442 (♀).

Type locality: ORANGE FREE STATE (R. C. Wroughton).

CAPE PROVINCE: Cape Town (L. Péringuey; Phillip).

15b. Subsp. **hottentotum** EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 26 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 218 (♀).

Type locality: Cape Town, CAPE PROVINCE (E. Simon).

15c. Subsp. **madecassum** FOREL, in Grandidier, 1892, 'Hist. Phys. Nat. Madagascar,' XX, 2, p. 255 (♀, ♀, ♂); 1907, Ann. Mus. Nat. Hungarici, V, p. 18 (♀). See p. 1027.

Monomorium minutum FOREL, in Grandidier, 1891, 'Hist. Phys. Nat. Madagascar,' XX, 2, p. 164 (♀, ♀, ♂).

GERMAN EAST AFRICA: Mto-ya-Kifaru (Katona).

16. **Monomorium mæstum** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 74, fig. 7 (♀).

Type locality: Naivasha, Rift Valley, 1900 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

17. **Monomorium opacum** FOREL, 1913, Rev. Zool. Afr., II, p. 333 (♀). BEQUAERT, *ibid.*, p. 428.

Type locality: Shinsenda, BELGIAN CONGO (J. Bequaert).

18. **Monomorium ophthalmicum** FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 87 (♀).

Type locality: Southern ABYSSINIA (Ilg).

- Monomorium oscaris** FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 86 (♀). EMERY, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 61 (♀).
Type locality: Southern ABYSSINIA (Ilg).
ERITREA: Ghinda (Belli).
191. Var. **excensuræ** FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 342 (♀).
Type locality: Kentani, CAPE PROVINCE (Miss A. Pegler).
192. Var. **nuptiale** FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 216 (♀).
ARNOLD, 1916, Ann. South African Mus., XIV, p. 229 (♀).
Type locality: Bembesi, RHODESIA (G. Arnold).
RHODESIA: Bulawayo (G. Arnold).
19a. Subsp. **musicum** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 442 (♀, ♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 230 (♀, ♀).
Type locality: NATAL (Haviland).
19b. Subsp. **springvalense** FOREL, 1913, Bull. Soc. Ent. Belgique, LVII, p. 136 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 230 (♀).
Type locality: Springvale, Matoppos, SOUTHERN RHODESIA (G. Arnold).
19b1. Var. **paternum** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 248 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 231 (♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 418 (♀).
Type locality: Table Mt., 1800 ft., CAPE PROVINCE (G. Arnold).
BELGIAN CONGO: St. Gabriel (Kohl).
20. **Monomorium osiridis** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 258, fig. 7 (♀).
Type locality: Bura, BRITISH EAST AFRICA (Alluaud and Jeannel).
21. **Monomorium pacis** FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 343 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 234 (♀).
Type locality: Cape Town, CAPE PROVINCE.
22. **Monomorium pharaonis** (LINNÆUS) MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 752; 1863, *ibid.*, XIII, p. 429. ROGER, 1863, 'Verzeich. Formicid.,' p. 32. MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidæ, p. 90 (♂); 1867, Tijdschr. v. Ent., X, p. 95 (♀, ♀, ♂). FRAUENFELD, 1867, Verh. Zool. Bot. Ges. Wien, XVII, p. 441. ERN. ANDRÉ, 1874, Rev. Mag. Zool., (3) II, p. 200. EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 368 (♀, ♀). SAUNDERS, 1880, Trans. Ent. Soc. London, p. 222 (♀, ♀, ♂). EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 532 (♀). ERN. ANDRÉ, 1883, 'Spec. Hym. Europ. Algérie,' II, pp. 333, 338, and 342 (♀, ♀, ♂). MAGRETTI, 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 244 (♀); 1884, Ann. Mus. Civ. Genova, XXI, p. 539 (♀). REUTER, 1884, Öfvers. Finsk. Vet.-Soc. Förh., XXVI, pp. 1-21. PREUDHOMME, 1885, Ann. Soc. Ent. Belgique, XXIX, C. R., p. cxxxvii. KORLEVIC, 1886, Rovart. Lapok, III, pp. 18 and iv. FROMHOLTZ, 1886, Ent. Nachr., XII, p. 122. MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 359. PROVANCHER, 1887, 'Addit. Faune Ent. Canada, Hym.,' p. 249. BELLEVOYE, 1888, Ann. Soc. Ent. France, (6) VIII, Bull., p. clxxvii. RILEY, 1888, 'Insect Life,' II, p. 106, fig. 18. BELLEVOYE, 1890, *ibid.*, II, pp. 230-233. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 69. MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 201; 1895, Ann. Naturh. Hofmus. Wien, X, p. 133. EMERY, 1895, Ann. Soc. Ent. Belgique, XXXIX, p. 337. MAYR, 1896, Ent. Tidskr., XVII, p. 238. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 40. FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p.

81 (♀). SJÖSTEDT, 1904, 'I Västafrikas Urskogar,' p. 29. MAYR, in Sjöstedt, 1907, 'Expd. Kilimandjaro, Meru, etc.,' II, 8, p. 12; 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. 1, p. 387. EMERY, 1908, Deutsch. Ent. Zeitschr., p. 684 (♀, ♀, ♂). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 145 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 351 (♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 354 (♀, ♀). FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 245 (♀, ♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 228 (♀, ♀, ♂). WASMANN, 1918, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VII, p. 147. SANTSCHI, 1920, 'Études Maladies Parasites Cacaoyer, S. Thomé,' X, p. 2. See pp. 162 and 1027.

Formica pharaonis LINNÆUS, 1758, 'Syst. Nat.,' Ed. 10, I, p. 580; 1764, 'Mus. Ludov. Ulric.,' p. 418; 1767, 'Syst. Nat.,' Ed. 12, I, 2, p. 963. P. L. MÜLLER, in Linnæus, 1775, 'Vollst. Natursyst.,' V, p. 913. GMELIN, in Linnæus, 1790, 'Syst. Nat.,' Ed. 13, I, 5, p. 2799. CHRIST, 1791, 'Naturg. d. Insect.,' p. 514. OLIVIER, 1791, 'Encycl. Méthod. Insect.,' VI, p. 503. LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 290.

Formica antiquensis FABRICIUS, 1793, 'Ent. Syst.,' II, p. 357. LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 285. FABRICIUS, 1804, 'Syst. Piez.,' p. 404.

Myrmica domestica SHUCKARD, 1838, Mag. Nat. Hist., (2) II, p. 627 (♀, ♀). F. SMITH, 1851, 'List Brit. Anim. Brit. Mus.,' VI, Aculeat., p. 119 (♀, ♂). DANIELL, 1852, Proc. Linn. Soc. London, II, pp. 172-177. WAKEFIELD, 1853, Zoologist, XI, p. 3810. DANIELL, 1853, *ibid.*, XI, pp. 3769-3773. CURTIS, 1854, Trans. Linn. Soc. London, XXI, p. 217. F. SMITH, 1855, Trans. Ent. Soc. London, (2) III, p. 130 (♀, ♀, ♂). 1859, Zoologist, XVII, p. 6387. SHARP, 1869, Entomologist, IV, p. 232. GILLAM, 1870, *ibid.*, V, p. 83.

Myrmica unifasciata BOSTOCK, 1839, Trans. Ent. Soc. London, II, Proc., pp. li-iii.

Atta minuta JERDON, 1851, Madras Journ. Litt. Sc., XVII, p. 105; 1854, Ann. Mag. Nat. Hist., (2) XIII, p. 47 (♀, ♀).

Myrmica (Ecophthora) domestica NYLANDER, 1856, Ann. Sc. Nat. Zool., (4) V, p. 98 (♀, ♀, ♂), Pl. III, figs. 53, 54.

Myrmica (Diplorhoptrum) molesta F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 122 (*nec Say*); 1858, 'Cat. Brit. Fossor. Hym.,' p. 34 (♀, ♀, ♂); 1858, Trans. Ent. Soc. London, (2) IV, p. 284.

Myrmica (Monomarium) fragilis F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 124 (♀).

Myrmica (Monomarium) contigua F. SMITH, *ibid.*, p. 125 (♀).

Myrmica (Monomarium) molesta F. SMITH, *ibid.*, p. 130 (*nec Say*).

Diplorhoptrum fugax LUCAS, 1858, Ann. Soc. Ent. France, (3) VI, Bull., p. lxxxi (*nec Mayr, nec auct.*).

Myrmica molesta MEINERT, 1860, Danske Vidensk. Selsk. Skrift., naturv. math. Afd., (5) V, p. 335 (♀, ♀, ♂) (*nec Say*). F. SMITH, 1860, Journ. Proc. Linn. Soc. London, Zool., IV, Suppl., p. 73 (*nec Say*); 1862, Trans. Ent. Soc. London, (3) I, p. 33 (*nec Say*); 1863, Ent. Annual for 1863, p. 59 (*nec Say*). RILEY, 1870, Second Ann. Rep. Ins. Missouri, p. 11 (*nec Say*); 1877, Ninth Ann. Rep. Ins. Missouri, p. 43 (*nec Say*). PROVANCHER, 1881, Natural. Canadien, XII, p. 360 (♀, ♀, ♂) (*nec Say*); 1883, 'Faune Ent. Canada, Hym.,' p. 603 (♀, ♀, ♂) (*nec Say*).

Myrmica pharaonis ROGER, 1862, Berlin. Ent. Zeitschr., VI, p. 294 (♀, ♂).

Diplorhoptrum domesticum GASKELL, 1877, Ent. Monthly Mag., XIII, p. 254.
WHITE, 1883, 'Ants and Their Ways,' p. 268, fig. 18.

Type locality: EGYPT.

Cosmopolitan. SENEGAMBIA: Dakar (F. Silvestri). LIBERIA: Junk River. TOGO: Bismarckburg (Conradt). SAN THOMÉ: (de Seabra). CAMEROON: (Sjöstedt); Moliwe Region (v. Maltzahn); Duala (v. Rothkirch). BELGIAN CONGO: Boma (Styczinski); Stanleyville; Thysville (Lang and Chapin); Sankuru (Laja). CAPE PROVINCE: Cape Town (R. Lightfoot). NATAL: Durban (Marley). GERMAN EAST AFRICA: Tanga (Sjöstedt). ZANZIBAR: (Stuhlmann). UGANDA: Gondokoro (F. Werner). ERITREA: Massaua (Beccari). OBOCK: (C. Alluaud).

23. **Monomorium rhopalocorum** EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 25 (♀), Pl. II, fig. 29. SANTSCHI, 1910, *ibid.*, LXXVIII, (1909), p. 359, fig. 4a (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 215 (♀).

Type locality: Cape Town, CAPE PROVINCE (E. Simon).

23. Var. **gabrielesse** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 418 (♀, ♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

23a. Subsp. **speluncarum** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 72, fig. 6 (♀). ALLUAUD and JEANNEL, 1914, Arch. Zool. Gén. Exp., LIII, p. 366.

Type locality: Shimon, BRITISH EAST AFRICA (Alluaud and Jeannel).

24. **Monomorium schultzei** FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 18 (♀, ♀); 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 244 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 231 (♀, ♀).

Type locality: Steinkopf, CAPE PROVINCE (L. Schultze).

CAPE PROVINCE: Lower Albany; Grahamstown (Hewitt). GERMAN SOUTHWEST AFRICA: Prince of Wales Bay, south of Angra Pepuefia (L. Schultze).

25. **Monomorium voeltzkowi** FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 78 (♀).

Type locality: Chake Chake, Pemba Island, BRITISH EAST AFRICA (Voeltzkow).

26. **Monomorium zulu** SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 18 (♀). TRÄGÅRDH, *ibid.*, p. 44. ARNOLD, 1916, Ann. South African Mus., XIV, p. 215 (♀).

Type locality: Junction of the Umfolosi Rivers, ZULULAND (I. Trägårdh).

Subgenus 2. **Syllophopsis** SANTSCHI

Monomorium subg. *Syllophopsis* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 259.

Subgenotype: *Monomorium modestum* Santschi, 1915.

27. **Monomorium (Syllophopsis) modestum** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 250, fig. 8 (♀).

Monomorium modestum SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 17 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 214 (♀).

Type locality: Stamford Hill, NATAL (I. Trägårdh).

27. Var. **smutsi**, new name.

Monomorium (Syllophopsis) modestum var. *boerorum* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 260, fig. 9 (♀) (*nec M. minutum* subsp. *boerorum* Forel, 1910).

Type locality: Pretoria, TRANSVAAL.

Subgenus 3. **Xeromyrmex** EMERY

Monomorium subg. *Xeromyrmex* EMERY, 1915, Bull. Soc. Ent. France, pp. 190 and 191.

Subgenotype: *Formica salomonis* LINNÆUS, 1758.

28. ***Monomorium (Xeromyrmex) afrum*** ERN. ANDRÉ.

Monomorium afrum ERN. ANDRÉ, in Magretti, 1884, Ann. Mus. Civ. Genova, XXI, p. 540 (♀); in Magretti, 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 244 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 65. FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 136 (♀). REICHENSPERGER, 1913, Zool. Jahrb. Abt. Syst., XXXV, p. 198. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 72 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 206 (♀), Pl. v, figs. 58, 58a, 58b, 58c.

Type locality: Atbara River, ANGLO-EGYPTIAN SUDAN (Magretti).

RHODESIA: Bulawayo (G. Arnold). BRITISH EAST AFRICA: Tiwi; Samburu, Wa-Nyika (Alluaud and Jeannel).

28₁. Var. ***asmarense*** FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 250 (♀, ♂). EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 4 (♀).

Type locality: Asmara, ERITREA (K. Escherich).

ERITREA: Ghinda (K. Escherich); Nefasit (F. Silvestri; K. Escherich).

28₂. Var. ***fulvor*** FOREL, 1913, Rev. Zool. Afr., II, p. 332 (♀). BEQUAERT, *ibid.*, p. 427. See p. 163.

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

BELGIAN CONGO: Niapu; Gatamba (Lang and Chapin).

29. ***Monomorium (Xeromyrmex) albopilosum*** EMERY.

Monomorium albopilosum EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 24 (♀). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Af., III, p. 17 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 207 (♀).

Type locality: Bloemfontein, TRANSVAAL (E. Simon).

CAPE PROVINCE: Kimberley (E. Simon). TRANSVAAL: Makapan (E. Simon). ZULULAND: Umfolosi (I. Trägårdh). BASUTOLAND: Leribe (Weitzcker).

29₁. Var. ***clarithorax*** SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 235.

Type locality: NATAL (Haviland).

29₂. Var. ***thales*** FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 136 (♀); 1916, Rev. Suisse Zool., XXIV, p. 417 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 207 (♀, ♀, ♂), Pl. v, fig. 61.

Type locality: Springvale, RHODESIA (G. Arnold).

RHODESIA: Matoppo Hills; Bulawayo (G. Arnold). TRANSVAAL: Shilouwane (Juned); Pretoria (Lounsbury).

29a. Subsp. ***paucipilosum*** SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 235 (♀).

Type locality: NATAL (Haviland).

30. ***Monomorium (Xeromyrmex) bicolor*** EMERY. See p. 162.

Monomorium bicolor EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 368 (♀); 1881, *ibid.*, XVI, pp. 531 and 532 (♀). ERN. ANDRÉ, 1883, 'Spec. Hym. Europ. Algérie,' II, p. 334 (♀). MAGRETTI, 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 244 (♀); 1884, Ann. Mus. Civ. Genova, XXI, p. 540 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 66. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 88; 1901, *ibid.*, X, p. 311. MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 7 (♀, ♀). FOREL, in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 564; 1907, Ann. Mus. Nat. Hungarici,

V, p. 18 (♀); 1907, Mitt. Naturh. Mus. Hamburg, XXIV, p. 15 (♀). EMERY, 1908, Deutsch. Ent. Zeitschr., p. 677 (♀). FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 251 (♀, ♀). SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 163; 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 353 (♀). EMERY, 1915, *ibid.*, X, p. 4 (♀).

Monomorium bicolor var. *cærulescens* SANTSCHI, 1912, Bull. Soc. Hist. Nat. Afr. Nord, IV, p. 148 (♀).

Type locality: Sciotel, ERITREA (Beccari).

SENEGAMBIA: Dakar (F. Silvestri). PORTUGUESE GUINEA: Bissao (Knipping). FRENCH GUINEA: Kindia; Mamou (F. Silvestri). IVORY COAST: (Lohier). BELGIAN CONGO: Leopoldville; Garamba (Lang and Chapin). MOSSAMEDES: between the Cubango and Cuito Rivers (Baum and Van der Kellen). ORANGE FREE STATE: Bothaville (H. Brauns). GERMAN EAST AFRICA: Mto-ya-Kifaru (Katona). SOUTHERN AFRICA: (Ilg). ERITREA: Nefasit (F. Silvestri; K. Escherich); Ghinda (K. Escherich). ANGLO-EGYPTIAN SUDAN: Suakin; Metemma (Magretti). FRENCH SOMALILAND: Obock (Maindron).

30a. Subsp. *dakarense* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 353 (♀).

Type locality: Dakar, SENEGAMBIA (F. Silvestri).

SENEGAMBIA: Longa (Roubaud).

30b. Subsp. *hirsutum* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 251 (♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 353 (♀).

Type locality: Nefasit, ERITREA (K. Escherich).

SOMALILAND.

30c. Subsp. *nitidiventris* EMERY, 1893, Ann. Soc. Ent. France, LXII, p. 256 (♀). MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 4. EMERY, 1908, Deutsch. Ent. Zeitschr., p. 677, fig. 5 (♀). KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 5, figs. 1 and 2 (♀, ♀, ♂). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 353 (♀). FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 244 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 209 (♀, ♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 73 (♀). A. J. CHALMERS, 1919, Journ. Trop. Med. Hyg., London, XXII, p. 117.

Monomorium bicolor var. *nitidiventris* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 441 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 209 (♀, ♀).

Type locality: Aden, SOUTHERN ARABIA (E. Simon).

EGYPT: Lower Egypt; Assuan (I. Trägårdh). ANGLO-EGYPTIAN SUDAN: Khartoum (Karawaiew). BRITISH EAST AFRICA: Naivasha, Rift Valley, 1900 m. (Alluaud and Jeannel). RHODESIA: Victoria Falls (R. C. Wroughton); Bulawayo (G. Arnold). ORANGE FREE STATE: Bothaville (H. Brauns).

31. *Monomorium (Xeromyrmex) hannonis* SANTSCHI.

Monomorium hannonis SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 358 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

32. *Monomorium (Xeromyrmex) medinae* FOREL.

Monomorium medinae FOREL, 1892, Ann. Soc. Ent. Belgique, XXXVI, p. 454 (♀). EMERY, 1908, Deutsch. Ent. Zeitschr., p. 679, fig. 9 (♀).

Type locality: Laguna, CANARY ISLANDS (Medina).

32a. Subsp. **fridae** FOREL, 1905, Ann. Soc. Ent. Belgique, XLIX, p. 183 (♀).
ARNOLD, 1916, Ann. South African Mus., XIV, p. 219 (♀).

Type locality: Willowmore, CAPE PROVINCE (H. Brauns).

33. **Monomorium (Xeromyrmex) salomonis** (LINNÆUS) EMERY, 1915, Bull. Soc. Ent. France, p. 190. See p. 1027.

Formica salomonis LINNÆUS, 1758, 'Syst. Nat.', Ed. 10, I, p. 580.

Monomorium salomonis ERN. ANDRÉ, 1883, 'Spec. Hym. Europ. Algérie,' II, pp. 336, 339 and 342 (♀, ♀, ♂), Pl. XXII, figs. 1-4. MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 4. EMERY, 1908, Deutsch. Ent. Zeitschr., p. 674 (♀, ♀, ♂).

Monomorium salomonis ARNOLD, 1916, Ann. South African Mus., XIV, p. 219 (♀, ♀, ♂).

Type locality: EGYPT.

Cosmopolitan. ANGLO-EGYPTIAN SUDAN: Ondurman; Wadi Halfa (I. Trägårdh).

33i. Var. **parvinode** FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 88 (♀).

Type locality: Southern ABYSSINIA (Ilg).

33j. Var. **pullulum** SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 235 (♀).

Type locality: Senegal, SENEGAMBIA (Claveau).

33a. Subsp. **carbo** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 246 (♀).
EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 4 (♀).

Monomorium salomonis var. **carbo** FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 251 (♀).

Type locality: Ghinda, ERITREA (K. Escherich).

ERITREA: Nefasit (F. Silvestri).

33b. Subsp. **damarense** FOREL, in Schultze, 1910, 'Forschungsreise in Süd-afrika,' IV, p. 17 (♀).

Monomorium salomonis subsp. **damarense** ARNOLD, 1916, Ann. South African Mus., XIV, p. 223 (♀).

Type locality: Gawieb, GERMAN SOUTHWEST AFRICA (L. Schultze).

33c. Subsp. **delagoense** FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 87 (♀); 1910, Ann. Soc. Ent. Belgique, LIV, p. 441 (♀, ♀, ♂). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 17 (♀). TRÄGÅRDH, *ibid.*, p. 44.

Monomorium salomonis subsp. **delagoensis** ARNOLD, 1916, Ann. South African Mus., XIV, p. 226 (♀, ♀, ♂).

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA (Liengme).

NATAL: (Haviland); Estcourt; Pietermaritzburg (I. Trägårdh). ZULULAND: Dukudu; Umfolosi (I. Trägårdh).

33c. Var. **grahamstownense** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 245 (♀).

Monomorium salomonis subsp. **delagoense** var. **grahamstownensis** ARNOLD, 1916, Ann. South African Mus., XIV, p. 227 (♀).

Type locality: Grahamstown, CAPE PROVINCE (G. Arnold).

33d. Subsp. **herero** FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 16 (♀, ♀). SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 284.

Monomorium salomonis subsp. **herero** ARNOLD, 1916, Ann. South African Mus., XIV, p. 222 (♀, ♀).

Type locality: Possession Island, GERMAN SOUTHWEST AFRICA (L. Schultze).

- 33d. Var. **belli** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 245 (♀).
Monomorium salamonis subsp. *herero* var. *belli* ARNOLD, 1916, Ann. South African Mus., XIV, p. 223 (♀).
 Type locality: Willowmore, CAPE PROVINCE (H. Brauns).
- 33d. Var. **willowmorensae** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 245 (♀).
Monomorium salamonis subsp. *herero* var. *willowmorensis* ARNOLD, 1916, Ann. South African Mus., XIV, p. 222 (♀).
 Type locality: Willowmore, CAPE PROVINCE (H. Brauns).
- 33e. Subsp. **junodi** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 441 (♀); 1913, *ibid.*, LVII, p. 137 (♀).
Monomorium salamonis subsp. *junodi* ARNOLD, 1916, Ann. South African Mus., XIV, p. 221 (♀).
 Type locality: Shilouvane, TRANSVAAL (Junod).
 RHODESIA: Bulawayo (G. Arnold).
- 33e. Var. **opacius** FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 136 (♀).
 SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 235.
Monomorium salamonis var. *opacior* ARNOLD, 1916, Ann. South African Mus., XIV, p. 220 (♀, ♀).
 Type locality: Bulawayo, RHODESIA (G. Arnold).
- 33f. Subsp. **ocellatum** SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 377 (♀).
 Type locality: Willowmore, CAPE PROVINCE (H. Brauns).
- 33g. Subsp. **subopacum** (F. SMITH) EMERY, 1908, Deutsch. Ent. Zeitschr., p. 676, fig. 4 (♀, ♀). WASMANN, 1912, Zeitschr. Wiss. Zool., CI, p. 107.
Myrmica (Monomorium) subopaca F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 127 (♀, ♀).
Monomorium mediterraneum MAYR, 1861, 'Europ. Formicid.,' p. 72 (♀, ♀).
 ROGER, 1862, Berlin. Ent. Zeitschr., VI, p. 294.
- Monomorium subopacum* subsp. *mediterraneum* EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 531 (♀).
Monomorium subopacum var. *mediterraneum* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 133.
- Monomorium subopacum* MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 753 (♀); 1863, *ibid.*, XIII, p. 429. ERN. ANDRÉ, 1874, Rev. Mag. Zool., (3) II, p. 199. EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 531 (♀); 1892, *ibid.*, XXXII, p. 117. ERN. ANDRÉ, 1883, 'Spec. Hym. Europ. Algérie,' II, pp. 336 and 339 (♀, ♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 70.
- Monomorium salomonis* var. *subopacum* FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 18 (♀). SANTSCHI, 1908, Ann. Soc. Ent. France, LXXVII, p. 517 (♀, ♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 353 (♀).
- Monomorium salamonis* subsp. *subopacum* ARNOLD, 1916, Ann. South African Mus., XIV, p. 224 (♀, ♀).
 Type locality: MADEIRA (T. V. Wollaston).
 CANARIES, MOROCCO, southern SPAIN, SARDINIA, SICILY, NAXOS, ALGIERS, EGYPT. CAPE PROVINCE: Cape Town (Raffray); Willowmore (H. Brauns). RHODESIA: Bulawayo (G. Arnold). SOMALILAND: Obbia; Gubbet (Bricchetti-Robecchi). GOLD COAST: Kitta (H. Brauns).

33g. Var. **anceps** (EMERY).

Monomorium subopacum var. *anceps* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 24 (♀).

Monomorium salomonis subsp. *subopacum* var. *anceps* ARNOLD, 1916, Ann. South African Mus., XIV, p. 225 (♀).

Type locality: Hamman's Kraal, TRANSVAAL (E. Simon).

33g. Var. **santschiellum**, new name.

Monomorium salomonis subsp. *subopacum* var. *senegalense* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 306 (♀) (*nec Monomorium senegalense* Roger, 1862).

Type locality: St. Louis, SENEGAMBIA (Claveau).

33h. Subsp. **termitarium** FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 17 (♀, ♀); 1910, Ann. Soc. Ent. Belgique, LIV, p. 441.

Monomorium salomonis subsp. *termitarium* ARNOLD, 1916, Ann. South African Mus., XIV, p. 224 (♂, ♀).

Type locality: Kooa, BECHUANALAND (L. Schultze).

MOSSAMEDES: (Baum and Van der Kellen).

33h. Var. **disertum** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 417.

Monomorium salomonis var. *diserta* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 216 (♀).

Monomorium salomonis var. *diserta* ARNOLD, 1916, Ann. South African Mus., XIV, p. 221 (♀).

Type locality: Shiloh, SOUTHERN RHODESIA (G. Arnold).

34. **Monomorium (Xeromyrmex) senegalense** ROGER.

Monomorium senegalense ROGER, 1862, Berlin. Ent. Zeitschr., VI, p. 294, footnote (♀); 1863, 'Verzeich. Formicid.,' p. 31. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 429. EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 530 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 70.

Type locality: Senegal, SENEGAMBIA.

35. **Monomorium (Xeromyrmex) setuliferum** FOREL.

Monomorium setuliferum FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 16 (♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 354 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 227 (♀).

Type locality: Khakhea, BECHUANALAND (L. Schultze).

ANGOLA: Quifangondo (F. Silvestri). GERMAN SOUTHWEST AFRICA: Okahandja (Casper). BECHUANALAND: Severelela (L. Schultze).

35. Var. **notulum** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 441 (♀, ♂); 1913, *ibid.*, LVII, p. 137 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 228 (♀, ♂).

Type locality: NATAL (Haviland).

SOUTHERN RHODESIA: Bulawayo; Springvale (G. Arnold).

36. **Monomorium (Xeromyrmex) subdentatum** FOREL.

Monomorium subdentatum FOREL, 1913, Rev. Zool. Afr., II, p. 332 (♀). BEQUAERT, *ibid.*, p. 427.

Type locality: Elisabethville, BELGIAN CONGO (J. Bequaert).

37. **Monomorium (Xeromyrmex) tchelihofoi** FOREL.

Monomorium tchelihofoi FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 244 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 211 (♀).

Type locality: Willowmore, CAPE PROVINCE (H. Brauns).

38. **Monomorium (Xeromyrmex) venustum** (F. SMITH).

Myrmica (Monomorium) venusta F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 126 (♀).

Monomorium venustum EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 531 (♀).
ERN. ANDRÉ, 1881, Ann. Soc. Ent. France, (6) I, p. 65 (♀, ♀, ♂), Pl. III, figs. 16-18;
1883, 'Spec. Hym. Europ. Algérie,' II, pp. 335, 339, and 343 (♀, ♀, ♂), Pl. XXII,
figs. 6 and 7. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 70. FOREL, 1894, Mitth.
Schweiz. Ent. Ges., IX, p. 88. EMERY, 1908, Deutsch. Ent. Zeitschr., p. 677, fig.
fig. 66 (♀, ♀).

Type locality: SYRIA.

SOUTHERN ABYSSINIA: (Ilg.).

Subgenus 4. **Parholcomyrme** EMERY

Monomorium subg. *Parholcomyrme* EMERY, 1915, Bull. Soc. Ent. France, p. 190.

Monomorium subg. *Parholcomyrme* EMERY, *ibid.*, p. 191.

Subgenotype: *Myrmica gracillima* F. Smith, 1861.

39. **Monomorium (Parholcomyrme?) amblyops** EMERY.

Monomorium amblyops EMERY, 1894, Bull. Soc. Ent. Italiana, XXVI, p. 148.

Type locality: MATTO GROSSO, BRAZIL.

39a. Subsp. *prossae* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 418.

Monomorium amblyops subsp. *bulawayense* FOREL, 1914, Bull. Soc. Vaudoise
Sc. Nat., (5) I, p. 247 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 236
(♀, ♀, ♂). (*nec Monomorium exiguum* var. *bulawayense* Forel, 1913).

Type locality: Hillside, Bulawayo, RHODESIA (G. Arnold).

40. **Monomorium (Parholcomyrme) australe** (EMERY).

Monomorium subopacum subsp. *australe* EMERY, 1886, Bull. Soc. Ent. Italiana,
XVIII, p. 363 (♀, ♀).

Monomorium australe DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 65.

Monomorium subopacum subsp. *australe* var. *laviceps* EMERY, 1886, Bull. Soc.
Ent. Italiana, XVIII, p. 364 (♀).

Monomorium salomonis subsp. *australe* FOREL, 1913, Deutsch. Ent. Zeitschr.,
Beih., p. 216 (♀, ♀).

Monomorium salomonis subsp. *australe* ARNOLD, 1916, Ann. South African Mus.,
XIV, p. 225 (♀, ♀).

Monomorium subopacum var. *australe* DIXEY and LONGSTAFF, 1907, Trans. Ent.
Soc. London, p. 339.

Monomorium (Paraholcomyrme) australe SANTSCHI, 1917, Ann. Soc. Ent. France,
LXXXV, (1916), p. 282 (♀, ♀).

Type locality: Cape of Good Hope, CAPE PROVINCE (L. Péringuey).

CAPE PROVINCE: Willowmore (H. Brauns); Kimberley (Dixey and Longstaff).

40a. Subsp. *havilandi* (FOREL) SANTSCHI, 1917, Ann. Soc. Ent. France,
LXXXV, (1916), p. 282 (♀, ♀).

Monomorium havilandi FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 443 (♀,
♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 216 (♀, ♀).

Type locality: NATAL (Haviland).

CAPE PROVINCE: Cape of Good Hope (L. Péringuey; G. Arnold).

41. **Monomorium (Parholcomyrmex) destructor** (JERDON).

Atta destructor JERDON, 1851, Madras Journ. Litt. Sc., XVII, p. 105; 1854, Ann. Mag. Nat. Hist., (2) XIII, p. 47 (♂).

Myrmica vastator F. SMITH, 1858, Journ. Proc. Linn. Soc. London, Zool., II, p. 71 (♂).

Myrmica ominosa GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 263 (♂); in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 517 (♂).

Myrmica atomaria GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 263 (♀); in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 518 (♀).

Myrmica (Monomarium) vastator F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 123 (♂).

Myrmica (Monomarium) basalis F. SMITH, *ibid.*, p. 125 (♂).

Monomorium ominosum ROGER, 1863, 'Verzeich. Formicid.,' p. 31. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 429.

Monomorium atomarium MAYR, 1863, *ibid.*, XIII, p. 429.

Monomorium basale MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidae, p. 92 (♂). EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 369, fig. (♂, ♀); 1881, *ibid.*, XVI, p. 532 (♂). MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 359 (♂).

Monomorium vastator MAYR, *ibid.*, p. 359. FOREL, 1897, Abhandl. Senckenberg. Naturf. Ges., XXI, p. 188 (♂).

Monomorium destructor EMERY, in Dalla Torre, 1893, 'Cat. Hym.,' VII, p. 66. SANTSCHI, 1908, Ann. Soc. Ent. France, LXXVII, p. 517 (♂). EMERY, 1908, Deutsch. Ent. Zeitschr., p. 671 (♂, ♀).

Type locality: INDIA (Jerdon).

Tropicopolitan. ERITREA: Sciotel (Beccari). ZANZIBAR: (Voeltzkow). PORTUGUESE EAST AFRICA: Quilimane (Peters).

41a. Subsp. **kalahariense** FOREL, in Schultze, 1910, 'Forschungsreise in Süd-afrika,' IV, p. 18 (♂). ARNOLD, 1916, Ann. South African Mus., XIV, p. 235 (♂).

Type locality: KOOA to Sekgoma, BECHUANALAND (L. Schultze).

41a1. Var. **despectum** FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 252 (♂); 1913, Rev. Zool. Afr., II, p. 331 (♂, ♀). BEQUAERT, *ibid.*, p. 427.

Type locality: Ghinda, ERITREA (K. Escherich).

BELGIAN CONGO: Sankisia (J. Bequaert).

42. **Monomorium (Parholcomyrmex) dispar** EMERY.

Monomorium dispar EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 24 (♂). ARNOLD, 1916, Ann. South African Mus., XIV, p. 235 (♂).

Type locality: Makapan, TRANSVAAL (E. Simon).

43. **Monomorium (Parholcomyrmex) gracillimum** (F. SMITH) EMERY, 1915, Bull. Soc. Ent. France, p. 190. See p. 1027.

Myrmica gracillima F. SMITH, 1862, Journ. Proc. Linn. Soc. London, Zool., VI, p. 34 (♂).

Monomorium gracillimum MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 753 (♂); 1863, *ibid.*, XIII, p. 429. EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 369, footnote, fig. (♂, ♀, ♂); 1881, *ibid.*, XVI, pp. 532 and 533 (♂). ERN. ANDRÉ, 1883, 'Spec. Hym. Europ. Algérie,' II, pp. 333, 337, and 340 (♂, ♀, ♂), Pl. xxii, figs. 8 and 9. MAGRETTI, 1884, Ann. Mus. Civ. Genova, XXI, p. 541 (♂); 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 244 (♂). DALLA TORRE, 1893, 'Cat. Hym.,'

VII, p. 67. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 88; 1902, Ann. Soc. Ent. Belgique, XLVI, p. 152 (♀). EMERY, 1908, Deutsch. Ent. Zeitschr., p. 669, fig. 2 (♀, ♀, ♂).

Monomorium (Holcomyrmez) gracillimum SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 354 (♀).

Type locality: SYRIA.

SYRIA, TUNIS, southern ALGERIA, central ASIA, ARABIA, INDIA, JAVA, LAYSAN, etc. FRENCH GUINEA: Mamou (F. Silvestri). ANGLO-EGYPTIAN SUDAN: Kassala; El Hefera, Settit (Magretti). SOUTHERN ABYSSINIA: (Ilg).

43. Var. *karawajewi* (FOREL).

Monomorium destructor subsp. *gracillimum* var. *karawajewi* FOREL, 1913, Rev. Suisse Zool., XXI, p. 437 (♀).

Monomorium minutum var. *pallidipes* KARAWAJEW, 1911, Rev. Russe Ent., XI, p. 7 (♀) (*nec* Forel).

Type locality: Khartum, ANGLO-EGYPTIAN SUDAN (Karawajew).

SYRIA: Rehobot near Jaffa (Aharoni).

43a. Subsp. *robustus* FOREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 352 (♀); 1894, *ibid.*, IX, p. 88. MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.', II, 8, p. 12. See pp. 163 and 1027.

Monomorium robustius DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 69.

Monomorium destructor subsp. *robustus* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 216 (♀).

Type locality: SOMALILAND (C. Keller).

SOUTHERN ABYSSINIA: (Ilg). BELGIAN CONGO: Yakuluku (Lang and Chapin). GERMAN EAST AFRICA: Mt. Meru (Sjöstedt).

Subgenus 5. *Holcomyrmez* MAYR

Holcomyrmez MAYR, 1878, Verh. Zool. Bot. Ges. Wien, XXVIII, p. 671.

Aphenogaster (part) ERN. ANDRÉ. *Atta* (part) ROGER.

Monomorium subg. *Holcomyrmez* EMERY, 1915, Bull. Soc. Ent. France, p. 191.

Subgenotype: *Holcomyrmez scabriceps* Mayr, 1878.

44. *Monomorium (Holcomyrmez) abyssinicum* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 250 (♀). SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 259 (♀).

Holcomyrmez abyssinicus FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 83 (♀).

Type locality: SOUTHERN ABYSSINIA (Ilg).

ERITREA: Ghinda (K. Escherich). DAROMEY: (Desanti).

Subgenus 6. *Lampromyrmez* MAYR

Lampromyrmez MAYR, 1868, Beiträge z. Naturkunde Preussens, I, p. 93.

Monomorium subg. *Mitara* EMERY, 1913, Ann. Soc. Ent. Belgique, LVII, p. 261 (type: *Monomorium lave* Mayr, 1879).

Subgenotype: *Lampromyrmez gracillimus* Mayr, 1868 = *Monomorium mayrianum* Wheeler, 1914, from the Baltic amber.

45. *Monomorium (Lampromyrmez) bequaerti* FOREL.

Monomorium (Martia) bequaerti FOREL, 1913, Rev. Zool. Afr., II, p. 334 (♀). BEQUAERT, *ibid.*, p. 428.

Type locality: Elisabethville, BELGIAN CONGO (J. Bequaert).

46. **Monomorium (Lampromyrmex) exiguum** FOREL.*Monomorium exiguum* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 85 (♀).*Monomorium (Martia) exiguum* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 351 (♀).

Type locality: SOUTHERN ABYSSINIA (Ilg).

BELGIAN CONGO: Leopoldville (J. Maes).

46i. Var. **bulawayense** FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 217

(♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 238 (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

46a. Subsp. **flavescens** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 418 (♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

46b. Subsp. **mictile** (FOREL) EMERY, 1914, Boll. Lab. Zool. Gen. Agrar.

Portici, VIII, p. 4 (♂, ♀).

Monomorium (Martia) atomus subsp. *mictile* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 252 (♂, ♀).*Monomorium (Mitara) atomus* subsp. *mictile* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 248. ARNOLD, 1916, Ann. South African Mus., XIV, p. 239 (♂, ♀).

Type locality: Ghinda, ERITREA (K. Escherich; F. Silvestri).

ERITREA: Nefasit (K. Escherich). NATAL: Durban (G. Arnold; C. B. Cooper).

47. **Monomorium (Lampromyrmex) faurei** SANTSCHI.*Monomorium (Mitara) atomus* subsp. *mictile* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 354 (♂) (nec FOREL).*Monomorium (Mitara) faurei* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 260, fig. 10 (♀).

Type locality: Gaboon, FRENCH CONGO (F. Faure).

FRENCH GUINEA: Mamou (F. Silvestri).

48. **Monomorium (Lampromyrmex) orientale** MAYR.*Monomorium orientale* MAYR, 1878, Verh. Zool. Bot. Ges. Wien, XXVIII, p. 670 (♂). EMERY, 1908, Deutsch. Ent. Zeitschr., p. 685 (♂, ♀).*Monomorium (Mitara) orientale* SANTSCHI, 1915, Bull. Soc. Hist. Nat. Afr. Nord, VII, p. 58, fig. 5 (♂).

Type locality: Calcutta, INDIA (Rothney).

48i. Var. **africanum** "MAYR" FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 78 (♀). This has apparently not been described.

Type locality: unknown.

BRITISH EAST AFRICA: Fundu Island near Pemba (Voeltzkow).

48s. Var. **clavicornis** (ERN. ANDRÉ) MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 4. EMERY, 1908, Deutsch. Ent. Zeitschr., p. 686.*Monomorium clavicornis* ERN. ANDRÉ, 1881, Ann. Soc. Ent. France, LI, p. 68 (♂), Pl. III, fig. 9; 1883, 'Spec. Hym. Europ. Algérie,' II, p. 332 (♂), Pl. XXII, fig. 5. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 66.

Type locality: SYRIA.

ANGLO-EGYPTIAN SUDAN: Kaka, White Nile (I. Trägårdh).

49. **Monomorium (Lampromyrmex) rosæ** SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 13, fig. 2c-f (♂).

Type locality: BELGIAN CONGO (J. Bequaert); no further locality is given, but the collector's field-number (No. 36) refers to Boma, according to verbal information received from Dr. Bequaert.

50. **Monomorium (Lampromyrmex) rotundatum** SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 14, fig. 2a (♀).

Type locality: Durban, NATAL (Marley).

Diplomorium MAYR

Diplomorium MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 16.

Genotype: *Diplomorium longipenne* Mayr, 1901.

1. **Diplomorium longipenne** MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 18 (♀, ♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 240 (♀, ♀).

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

Bondroitia FOREL

Diplomorium subg. *Bondroitia* FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 398.

Bondroitia FOREL, 1915, 'Fauna Ins. Helvetiæ, Hym., Formicidæ,' pp. 10 and 38.

Genotype: *Diplomorium (Bondroitia) lujæ* Forel, 1909.

1. **Bondroitia lujæ** (FOREL).

Diplomorium lujæ FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 72 (♀, ♀, ♂).

Diplomorium (Bondroitia) lujæ FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 398 (♀, ♀, ♂).

Type locality: Sankuru, BELGIAN CONGO (Luja).

Solenopsis WESTWOOD

Solenopsis WESTWOOD, 1840, Ann. Mag. Nat. Hist., VI, p. 86.

Diplorhoptrum MAYR, 1855, Verh. Zool. Bot. Ges. Wien, V, p. 449.

Atta (part) FABRICIUS, JERDON, F. SMITH, etc. *Crematogaster* (part) F. SMITH. *Formica* LATREILLE, etc. *Myrmica* LATREILLE, etc.

Genotype: *Solenopsis mandibularis* Westwood, 1840.

1. **Solenopsis africana** (SANTSCHI) EMERY, 1915, Rend. Accad. Sc. Bologna, N. S., XIX, p. 59, fig. 1 (♀).

Solenopsis fugax subsp. *africana* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 81 (♀, ♂).

Type locality: Blue Post Hotel, Kikuyu, 1550 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

2. **Solenopsis capensis** MAYR, 1866, Verh. Zool. Bot. Ges. Wien, XVI, p. 905 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 75.

Type locality: Cape of Good Hope.

3. **Solenopsis geminata** (FABRICIUS) ROGER, 1863, 'Verzeich. Formicid.,' pp. 32 and 49. MAYR, 1867, Tijdschr. v. Ent., X, p. 109 (♀, ♀, ♂); 1870, Verh. Zool. Bot. Ges. Wien, XX, p. 996 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 359; 1913, Ann. Soc. Ent. Belgique, LVII, p. 306 (2, ♀, ♀).

Atta geminata FABRICIUS, 1804, 'Syst. Piez.,' p. 423 (♀).

Solenopsis geminata var. *innota* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 257, fig. 6 (2, ♀, ♂).

Type locality: South America.

Tropicopolitan. LIBERIA: Monrovia (Delafosse). FRENCH CONGO: Sankita (F. Faure); Brazzaville (Valerio). BELGIAN CONGO: Goma (Valerio).

4. *Solenopsis gnomula* EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 13, fig. 6 (♀).

Type locality: Nefasit, ERITREA (F. Silvestri).

5. *Solenopsis maligna* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 359, fig. 5 (♀). EMERY, 1915, Rend. Accad. Sc. Bologna, N. S., XIX, p. 64, fig. 6 (♀).

Type locality: BRAZZAVILLE, FRENCH CONGO (A. Weiss).

6. *Solenopsis orbuloides* ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 321 (♀). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 77. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 144. SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 354 (♂, ♀). EMERY, 1915, Rend. Accad. Sc. Bologna, N. S., XIX, p. 59.

Type locality: SIERRA LEONE (Mocquerys).

NIGERIA: Olokemeji; Lagos (F. Silvestri).

7. *Solenopsis punctaticeps* MAYR, 1865, 'Reise Novara, Zool.', II, Formicidae, p. 109 (♂); 1870, Verh. Zool. Bot. Ges. Wien, XX, p. 996 (♂); 1887, *ibid.*, XXXVII, p. 616 (♂). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 77. WASMANN, 1899, Deutsch. Ent. Zeitschr., p. 405. MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 14 (♂). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 430; 1911, Rev. Zool. Afr., I, p. 276 (♂); 1913, Ann. Soc. Ent. Belgique, LVII, p. 138 (♂). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 355 (♂); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 19, figs. 1, 2 (♂, ♀). TRÄGÅRDH, *ibid.*, p. 44. EMERY, 1915, Rend. Accad. Sc. Bologna, N. S., XIX, pp. 60 *et seq.*, fig. 2 (♂). ARNOLD, 1916, Ann. South African Mus., XIV, p. 243 (♂, ♀, ♂), Pl. VI, figs. 78, 78a, 78b.

Solenopsis punctatoceps J. HEWITT, 1920, South African Journ. Nat. Hist., II, p. 109.

Type locality: Table Mountain, Cape of Good Hope (Novara Expedition).

RHODESIA: (G. Arnold). NATAL: Pietermaritzburg; Balgowan (I. Trägårdh); Estcourt (R. C. Wroughton); Shiyre (Haviland). ORANGE FREE STATE: Bothaville (H. Brauns). CAPE PROVINCE: Port Elizabeth (H. Brauns); St. Croix Island, Algoa Bay (Hewitt). ANGOLA: Quifangondo (F. Silvestri). BELGIAN CONGO: Congo da Lemba (R. Mayné).

7i. Var. *cleptomana* (SANTSCHI) EMERY, 1915, Rend. Accad. Sc. Bologna, N. S., XIX, p. 65, fig. 7 (♂).

Solenopsis cleptomana SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 80 (♂).

Type locality: Naivasha, Rift Valley, 1400 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

7a. Subsp. *caffra* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 16. EMERY, 1915, Rend. Accad. Sc. Bologna, N. S., XIX, p. 62, fig. 3, I and II (♂).

Solenopsis punctaticeps var. *caffra* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 99 (♂, ♀). EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 27. MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 15 (♂). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 431 (♀); 1913, Rev. Zool. Afr., II, p. 337 (♂). BEQUAERT, *ibid.*, p. 429. FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 242 (♂). ARNOLD, 1916, Ann. South African Mus., XIV, p. 246 (♂, ♀).

Type locality: TRANSVAAL (P. Berthoud).

BELGIAN CONGO: Elisabethville (J. Bequaert). BECHUANALAND: Kooa (L. Schultze). TRANSVAAL: Pretoria (E. Simon). NATAL: (R. C. Wroughton). CAPE PROVINCE: Table Mt. (G. Arnold). PORTUGUESE EAST AFRICA: Delagoa Bay: (Liengme).

7a1. Var. *cyclops* (SANTSCHI).

Solenopsis punctaticeps subsp. *cyclops* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 19, fig. 1, 1 (♀, ♂).

Solenopsis punctaticeps subsp. *caffra* var. (?) *cyclops* EMERY, 1915, Rend. Accad. Sc. Bologna, N. S., XIX, p. 62, fig. 3, III (♀).

Solenopsis punctaticeps var. *cyclops* ARNOLD, 1916, Ann. South African Mus., XIV, p. 246 (♀, ♂).

Type locality: NATAL (I. Trägårdh).

NATAL: Durban (Marley; C. B. Cooper).

7a2. Var. *diversipilosa* (MAYR).

Solenopsis punctaticeps var. *diversipilosa* MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 15 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 247 (♀).

Solenopsis punctaticeps subsp. *caffra* var. (?) *diversipilosa* EMERY, 1915, Rend. Accad. Sc. Bologna, N. S., XIX, pp. 62, 65.

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

CAPE PROVINCE: Grahamstown (Hewitt).

7b. Subsp. *erythraea* EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 12, fig. 5 (♀); 1915, Rend. Accad. Sc. Bologna, N. S., XIX, p. 64, fig. 5 (♀).

Type locality: Asmara, ERITREA (F. Silvestri).

ERITREA: Nefasit (F. Silvestri).

7c. Subsp. *indocilis* (SANTSCHI) EMERY, 1915, Rend. Accad. Sc. Bologna, N. S., XIX, p. 63, fig. 4 (♀).

Solenopsis punctaticeps var. *indocilis* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 355 (♀, ♀).

Type locality: Mamou, FRENCH GUINEA (F. Silvestri).

7d. Subsp. *kibaliensis* Wm. M. WHEELER. See p. 164 (♀, ♂).

Type locality: Vankerekhovenville, BELGIAN CONGO (Lang and Chapin).

8. *Solenopsis sævissima* (F. SMITH) MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 751 (♀).

Myrmica sævissima F. SMITH, 1855, Trans. Ent. Soc. London, (2) III, p. 166 (♀), Pl. XII, fig. 18.

Type locality: Tapajos River, BRAZIL (W. H. Bates).

Neotropical and apparently spreading to other continents.

8i. Var. *itinerans* (FOREL).

Solenopsis pylades var. *itinerans* FOREL, 1911, Rev. Zool. Afr., I, p. 276 (♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 355 (♀).

Type locality: Kigerama, GERMAN EAST AFRICA.

FRENCH GUINEA: Konakry (F. Silvestri).

Anergatides WASMANN.

Anergatides WASMANN, 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, p. 281.

Genotype: *Anergatides kohli* Wasmann, 1915.

1. **Anergatides kohli** WASMANN, 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, p. 283 (♀, ♂), Pls. VII and VIII.

Type locality: Fikilini, near Stanleyville, BELGIAN CONGO (H. Kohl).

Pheidologeton MAYR

Pheidologeton MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 750.

Phidologeton BINGHAM, 1903, 'Fauna Brit. India, Hym.,' II, p. 160.

Myrmica (part) HEER. *Ocodoma* (part) JERDON. *Pheidole* (part) F. SMITH. *Solenopsis* (part) F. SMITH.

Genotype: *Ocodoma diversa* Jerdon, 1851 = *Pheidole ocellifera* F. Smith, 1858.

1. **Pheidologeton diversus** (JERDON) ROGER, 1863, 'Verzeich. Formicid.,' p. 30.

Ocodoma diversa JERDON, 1851, Madras Journ. Litt. Sc., XVII, p. 109 (2 ♀); 1854, Ann. Mag. Nat. Hist., (2) XIII, p. 51 (2 ♀).

Type locality: Wynaad, SOUTHERN INDIA (Jerdon).

Oriental Region.

- 1a. Subsp. **standfussi** FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 386 (2).

Type locality: GUINEA (locality extremely doubtful).

2. **Pheidologeton solitarius** STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 141 (♀).

Type locality: Sokode Basan, Togo (Schröder).

The generic reference of this species is perhaps erroneous.

The following species has been referred to *Pheidologeton* by Mayr, 1866, Verh. Zool. Bot. Ges. Wien, XVI, p. 899; but owing to its 12-jointed antennae it cannot belong in that genus. It is impossible to place it without examining the type specimen.

Atta hostilis F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 165 (♀, ♀).

Type locality: Port Natal, NATAL.

Aneleus EMERY

Pheidologeton subg. *Aneleus* EMERY, 1900, Termes. Füzetek, XXIII, p. 327.

Genotype: *Solenopsis similis* Mayr, 1862.

1. **Aneleus diabolus** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 355, fig. 19 (2, ♀).

Oligomyrmex diabolus SANTSCHI, 1913, Bull. Soc. Ent. France, pp. 459 and 460 (2, ♀).

Type locality: Victoria, CAMEROON (F. Silvestri).

2. **Aneleus perpusillus** (EMERY) ARNOLD, 1916, Ann. South African Mus., XIV, p. 254 (2, ♀).

Pheidologeton perpusillum EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 26 (2, ♀), Pl. II, figs. 8-11. FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 431 (♀).

Type locality: Pretoria, TRANSVAAL (E. Simon).

CAPE PROVINCE: Kimberley (E. Simon). NATAL: in the Mountains (R. C. Wroughton).

- 2a. Subsp. **arnoldi** (FOREL) ARNOLD, 1916, Ann. South African Mus., XIV, p. 255 (2, ♀), Pl. VI, figs. 81, 81a.

Pheidologeton perpusillus subsp. *arnoldi* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 242 (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

2b. Subsp. **condescens** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 77 (♂, ♀).

Type locality: New Meschi, Mt. Kilimanjaro, 800 m., GERMAN EAST AFRICA (Alluaud and Jeannel).

BRITISH EAST AFRICA: Likoni; Cheteni (Alluaud and Jeannel).

2c. Subsp. **spinosus** (FOREL).

Pheidologeton (Aneleus) perpusillus subsp. *spinosus* FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 17 (♂).

Type locality: Kibosho, GERMAN EAST AFRICA (Katona).

3. **Aneleus politus** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 79, fig. 8 (♀).

Type locality: Blue Post Hotel, Kikuyu, 1520 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

4. **Aneleus silvestrii** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 357, fig. 20 (♂, ♀).

Type locality: Aburi, GOLD COAST (F. Silvestri).

Oligomyrmex MAYR

Oligomyrmex MAYR, 1867, Tijdschr. v. Ent., X, p. 110.

Genotype: *Oligomyrmex concinnus* Mayr, 1867.

1. **Oligomyrmex alluaudi** SANTSCHI, 1913, Bull. Soc. Ent. France, pp. 459 and 460 (♂, ♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 81, fig. 9 (♂, ♀).

Type locality: Shimoni, BRITISH EAST AFRICA (Alluaud and Jeannel).

BRITISH EAST AFRICA: Kijabé, Kikuyu Escarpment, 2100 m. (Alluaud and Jeannel).

1a. Var. **catartæ** SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 337 (♂, ♀).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

2. **Oligomyrmex angolensis** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 358, fig. 21 (♂, ♀).

Type locality: Quifangondo, ANGOLA (F. Silvestri).

2a. Subsp. **congolensis** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 417 (♂, ♀). Type locality: BELGIAN CONGO (H. Kohl).

3. **Oligomyrmex arnoldi** FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 123 (♀). SANTSCHI, 1913, Bull. Soc. Ent. France, p. 459 (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

4. **Oligomyrmex debilis** SANTSCHI, 1913, Bull. Soc. Ent. France, p. 459 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 360, fig. 22 (♂, ♀, ♂).

Oligomyrmex (Aëomyrma?) debilis SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 337 (♂, ♀).

Type locality: Kindia, FRENCH GUINEA (F. Silvestri).

FRENCH GUINEA: Konakry; Camayenne (F. Silvestri).

5. *Oligomyrmex erythræus* EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 14, fig. 7 (♂, ♀).

Type locality: Ghinda, ERITREA (F. Silvestri).

6. *Oligomyrmex jeanneli* SANTSCHI, 1913, Bull. Soc. Ent. France, pp. 459 and 460 (♂, ♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 83, fig. 10 (♂, ♀). ALLUAUD and JEANNEL, 1914, Arch. Zool. Gén. Exp., LIII, p. 366. EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 14 (♂).

Type locality: Shimon, BRITISH EAST AFRICA (Alluaud and Jeannel).

ERITREA: Nefasit (F. Silvestri).

Æromyrma FOREL

Æromyrma FOREL, in Grandidier, 1891, 'Hist. Nat. Phys. Madagascar,' XX, 2, p. 198; 1891, Ann. Soc. Ent. Belgique, XXXV, C. R., p. cccvii.

Genotype: *Æromyrma nosindambo* Forel, 1891.

1. *Æromyrma africana* FOREL, in Schultze, 1910, 'Forschungsreise in Süd-afrika,' IV, p. 15 (♂). ARNOLD, 1916, Ann. South African Mus., XIV, p. 256 (♂).

Type locality: Kooa to Sekgoma, BECHUANALAND (L. Schultze).

2. *Æromyrma arnoldiella* (SANTSCHI).

Oligomyrmex (*Æromyrma*) *arnoldiella* SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 340, figs. 3a, j, p, q, v, x (♂, ♀).

Type locality: NATAL (Haviland).

3. *Æromyrma hewitti* (SANTSCHI).

Oligomyrmex (*Æromyrma*) *hewitti* SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 341, figs. 3d, l, o, t, u (♂, ♀).

Type locality: Grahamstown, CAPE PROVINCE (Hewitt).

4. *Æromyrma incerta* (SANTSCHI).

Oligomyrmex (*Æromyrma*) *incertus* SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 340, figs. 3e, h (♂).

Type locality: Hillside, Bulawayo, RHODESIA (G. Arnold).

5. *Æromyrma lucida* (SANTSCHI).

Oligomyrmex (*Æromyrma*) *lucidus* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 284 (♂); 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 339, figs. 3c and i (♂).

Type locality: Bunthorne Mine, Bulawayo, RHODESIA (G. Arnold).

6. *Æromyrma nana* (SANTSCHI).

Oligomyrmex (*Æromyrma*) *nana* SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 338, figs. 3f, m, r, s (♂, ♀).

Type locality: Amatongas Forest, PORTUGUESE EAST AFRICA (G. Arnold).

7. *Æromyrma petulca* WM. M. WHEELER. See p. 166* (♂).

Type locality: Malela, BELGIAN CONGO (Lang and Chapin).

8. *Æromyrma semilævis* (MAYR).

Solenopsis semilævis MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 15 (♂). ARNOLD, 1916, Ann. South African Mus., XIV, p. 247 (♂).

Oligomyrmex (*Æromyrma*) *semilævis* EMERY, 1915, Rend. Accad. Sc. Bologna, N. S., XIX, p. 59.

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

8. Var. *trægaordhi* (SANTSCHI).

Oligomyrmex (Aëromyrma) semilævis var. *trægaordhi* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 285.

Aëromyrma trægaordhi SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 20, fig. 2 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 257 (♀).

Oligomyrmex (Aëromyrma) trægaordhi SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 338, figs. 3c, g (♀).

Type locality: Balgowan, NATAL (I. Trägårdh).

NATAL: Stamford Hill (I. Trägårdh).

9. *Aëromyrma vorax* SANTSCHI, 1913, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 145, fig. LXIX (♀); 1914, *ibid.*, VIII, p. 358 (♀). F. SILVESTRI, 1914, 'Report Exped. to Africa in Search Nat. Enem. Fruit Flies,' p. 128 (♀), Pl. xxiv, fig. 49.

Type locality: Aburi, GOLD COAST (F. Silvestri).

Carebara WESTWOOD

Carebara WESTWOOD, 1840, Ann. Mag. Nat. Hist., VI, p. 86.

Genotype: *Carebara lignata* Westwood, 1840.

1. *Carebara ampla* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 284 (♀, ♂); 1913, Ann. Soc. Ent. Belgique, LVII, p. 307.

Type locality: Majunga, BELGIAN CONGO (Gérard).

2. *Carebara arnoldi* (FOREL) SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 363, fig. 26 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 253 (♀).

Oligomyrmex arnoldi FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 123 (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

3. *Carebara junodi* FOREL, 1904, Ann. Soc. Ent. Belgique, XLVIII, p. 154 (♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, pp. 363 and 364, fig. 24 (♀).

Carebara vidua subsp. *junodi* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 53 (♀, ♂); 1913, Rev. Zool. Afr., II, p. 336 (♀, ♀, ♂). BEQUAERT, *ibid.*, p. 428. ARNOLD, 1916, Ann. South African Mus., XIV, p. 252 (♀, ♀).

Type locality: Shilouvane, TRANSVAAL (H. Junod).

BELGIAN CONGO: Katanga (Lemaire); Sankisia (J. Bequaert); Kalumba; Katumba (Neave); Lake Moero.

4. *Carebara langi* WM. M. WHEELER. See p. 173 (♀).

Type locality: Stanleyville, BELGIAN CONGO (Lang and Chapin).

5. *Carebara osborni* WM. M. WHEELER. See p. 174 (♂, ♀, ♂).

Type locality: Niangara, BELGIAN CONGO (Lang and Chapin).

6. *Carebara sicheli* MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 754 (♀); 1863, *ibid.*, XIII, p. 402. ROGER, 1863, 'Verzeich. Formicid.,' p. 32. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 74. MAYR, 1904, Verh. Zool. Bot. Ges. Wien, LIV, p. 596 (♀). SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 307 (♀).

Type locality: Senegal, SENEGAMBIA.

UPPER DAHOMEY. ANGLO-EGYPTIAN SUDAN: White Nile.

7. *Carebara silvestrii* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, pp. 362 and 364, fig. 23 (♀).

Type locality: Mamou, FRENCH GUINEA (F. Silvestri).

8. *Carebara vidua* F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 179 (♂). ROGER, 1863, 'Verzeich. Formicid.,' p. 32. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 402; 1866, *ibid.*, XVI, p. 904 (♀, ♂). EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 377 (♀, ♂). DISTANT, 1892, 'A Naturalist in the Transvaal,' p. 211. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 74. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 144 (♀). EMERY, 1899, Ann. Mus. Civ. Genova, XXXIX, p. 500 (♀, ♂). FOREL, 1901, Ann. Soc. Ent. Belgique, XLV, p. 392 (♀). BINGHAM, 1903, Ann. Mag. Nat. Hist., (7) XII, p. 61 (♀, ♂). WASMANN, in Jägerskiöld, 1904, 'Exped., XIII, Termitophilen,' p. 13 (♀). ZAVATTARI, 1907, Boll. Mus. Zool. Anat. Comp. Torino, XXII, No. 550, p. 2 (♀). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 12 (♀, ♂). EMERY, 1912, Annuar. Mus. Zool. Univ. Napoli, N. S., III, No. 26, p. 2 (♀). FOREL, 1913, Rev. Zool. Afr., II, p. 337 (♀, ♂). SANTSCHI, 1913, Ann. Soc. Ent. Belgique, XXXVII, p. 306 (♂); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 363, fig. 25 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 84 (♀, ♂). ARNOLD, 1916, Ann. South African Mus., XIV, p. 249 (♀, ♂), Pl. VI, figs. 79, 80. See p. 177.

Carebara colossus GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 263 (♀); in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 512 (♀), Pl. XXXII, fig. 8. ROGER, 1863, 'Verzeich. Formicid.,' p. 32. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 402.

Carebara duz F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 179 (♀). ROGER, 1863, 'Verzeich. Formicid.,' p. 32. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 402.

Carebara vidua var. *duz* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 53 (♀); 1913, Rev. Zool. Afr., II, p. 337 (♀).

Carebara vidua E. WARREN, 1909, Ann. Natal Mus., II, pt. 1, p. 120.

Carebara vidua var. *abdominalis* SANTSCHI, 1912, Bull. Soc. Ent. France, p. 285 (♀).

Carebara vidua subsp. *abdominalis* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 84 (♀).

Type locality: Port Natal, NATAL.

BELGIAN CONGO: Niangara; Faradje (Lang and Chapin); Yakuluku (J. Rodhain); Katanga (Lemaire); valley of the Lubumbashi (Buttgenbach); Kwesi to Kilo (Bayer). ANGLO-EGYPTIAN SUDAN: Gebel Ahmed Aga, White Nile (I. Trägårdh); Niam Niam. ERITREA: Keren (Beccari). ABYSSINIA: Sancarar to Amarr Burgi; Badditu to Dimé; Dimé to Bass Narok (V. Bottego). SOMALILAND: Giuba River (Elena d'Aosta). BRITISH EAST AFRICA: Bura, Wa-Taita; Kibwezi, Wakamba (C. Alluaud). UGANDA: Kampala (C. C. Gowdey). RHODESIA: Hartley (Zeally); Kazungula (Jallá); Southern Rhodesia (G. Arnold); Lake Ngami. PORTUGUESE EAST AFRICA: Tete (Peters). TRANSVAAL: Pretoria (Distant; Wichgraf); Barberton (Rendall). NATAL: (Haviland). GERMAN SOUTHWEST AFRICA: Okahandja (Casper; Dinter).

***Pædalagus* FOREL**

Pædalagus FOREL, in Escherich, 1911, 'Termitenleben auf Ceylon,' p. 217.

Genotype: *Pædalagus escherichi* Forel, 1911.

1. *Pædalagus infimus* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 364, fig. 27 (♀).

Oligomyrmex infimus SANTSCHI, 1913, Bull. Soc. Ent. France, p. 459 (♀).

Type locality: Kindia, FRENCH GUINEA (F. Silvestri).

2. *Pædalagus termitolestes* Wm. M. WHEELER. See p. 177 (♀).

Type locality: Malela, BELGIAN CONGO (Lang, Chapin and J. Bequaert).

Myrmecini Ashmead

Terataner EMERY

Terataner EMERY, 1912, Ann. Soc. Ent. Belgique, LVI, p. 103.

Genotype: *Atopomyrmex foreli* Emery, 1899.

1. *Terataner bottegoi* EMERY, 1912, Ann. Soc. Ent. Belgique, LVI, p. 103. SANTSCHI, 1914, 'Voy. Alluaud, et Jeannel Afr. Orient., Formicidae,' p. 72 (♀).

Atopomyrmex bottegoi EMERY, 1896, Ann. Mus. Civ. Genova, XXXVII, p. 155, fig. (♂).

Atopomyrmex bottegi R. GESTRO, in Vanutelli and Citerni, 1899, 'Seconda Spedizione Böttogo, L'Omo' p. 626, fig. 19 (♀).

Type locality: Lugh, SOMALILAND (V. Bottego).

BRITISH EAST AFRICA: Voi, Taita, 600 m. (Alluaud and Jeannel).

2. *Terataner luteus* EMERY, 1912, Ann. Soc. Ent. Belgique, LVI, p. 103.

Atopomyrmex luteus EMERY, 1899, *ibid.*, XLIII, p. 477 (♂). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 145 (♀).

Type locality: CAMEROON (Conradt).

CAMEROON: Mundame (Conradt).

Atopomyrmex ERN. ANDRÉ

Atopomyrmex ERN. ANDRÉ, 1889, Rev. d'Ent. Caen, VIII, p. 226.

Genotype: *Atopomyrmex mocquerysi* Ern. André, 1889.

1. *Atopomyrmex mocquerysi* ERN. ANDRÉ, 1889, Rev. d'Ent. Caen, VIII, p. 227 (♀). EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 562 (♀), Pl. xv, fig. 7. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 61. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 83. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 3 (♀). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 134. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 41. EMERY, 1899, Ann. Mus. Civ. Genova, XXXIX, p. 500 (♀). MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 4 (♀); 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. 1, p. 387. FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 12 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 145 (♀). FOREL, 1911, Rev. Zool. Afr., I, p. 276 (♀). EMERY, 1912, Ann. Soc. Ent. Belgique, LVI, p. 102. FOREL, 1913, Rev. Zool. Afr., II, p. 335 (♀, ♂). BEQUAERT, *ibid.*, p. 428. SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 351 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 71 (♀). See p. 181.

Atopomyrmex mocquerysi var. *australis* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 16 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 193 (♀). Same as type, *teste* Arnold *in litt.*

Type locality: Dakar, SENEGAMBIA (Mocquerys).

DAHOMY: Kotonou (F. Silvestri). SLAVE COAST. FRENCH CONGO: Ogowe (Mocquerys). BELGIAN CONGO: Lukolela to Basoko; Akenge; Medje; Faradje (Lang and Chapin); Matadi; Sankisia (J. Bequaert); Kondué (Luja). ANGOLA:

ANGLO-EGYPTIAN SUDAN: Kaka, White Nile (I. Trägårdh); Khor Attar (F. Werner). ABYSSINIA: Buditu to Bass Narok (V. Bottego). SOMALILAND. BRITISH EAST AFRICA: Voi, Wa-Taita, 600 m. (C. Alluaud). ZANZIBAR. PORTUGUESE EAST AFRICA: Lobombo Borges, Mozambique (Wilms); Delagoa Bay (Liengme; P. Berthoud); Xalasi (C. W. Howard). NORTH WEST RHODESIA: Mwangwa (H. Dollmann). ZULULAND: (I. Trägårdh).

1. Var. *curvispina* FOREL, 1911, Deutsch. Ent. Zeitschr., p. 311 (♀); 1914, Bull. Soc. Vaudoise Sc. Nat., (5) 1, p. 232; 1916, Rev. Suisse Zool., XXIV, p. 419 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 191 (♀, ♀).

Type locality: Kondué, BELGIAN CONGO (Luja).

BELGIAN CONGO: Lumbulumbu (H. Kohl). SOUTHERN RHODESIA: Sipapoma, Ungusa River; Malindi; Victoria Falls (G. Arnold). ZULULAND: (I. Trägårdh).

1a. Subsp. *cryptoceroides* (EMERY) FOREL, 1913, Rev. Zool. Afr., II, p. 336 (♀). BEQUAERT, *ibid.*, p. 428. FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) 1, p. 343 (♀). See p. 182.

Atopomyrmex cryptoceroides EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 561 (♀), Pl. xv, figs. 5 and 6. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 60. EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 477 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 145, fig. 10 (♀). EMERY, 1912, Ann. Soc. Ent. Belgique, LVI, p. 102. SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 351 (♀). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 374 (♀).

Atopomyrmex deplanatus MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 133 (♀).

Type locality: Assinie, IVORY COAST (C. Alluaud).

FRENCH GUINEA: Kakoulina; Mamou; Kindia (F. Silvestri). SIERRA LEONE: Samlia Falls, River N'Garnie (Moequers). GOLD COAST. SOUTHERN NIGERIA: Ibadan (F. Silvestri). CAMEROON: Mundame (Conradt). FERNANDO PO: (Conradt). SPANISH GUINEA: Alen; Uelleburg (Tessinann). FRENCH CONGO: Fort de Possel to Fort Crampel (Schubotz); Kuilu River. BELGIAN CONGO: Kondué (Luja); St. Gabriel; Bengamisa (Kohl); Malela; Elisabethville (J. Bequaert).

1a. Var. *melanoticus* WM. M. WHEELER. See p. 182 (♀).

Type locality: between Lukolela and Basoko, BELGIAN CONGO (Lang and Chapin).

Atopula EMERY

Atopula EMERY, 1912, Ann. Soc. Ent. Belgique, LVI, p. 104.

Vollenhorvia subg. *Atopula* EMERY, 1912, *ibid.*, LVI, p. 273.

Genotype: *Atopomyrmex nodifer* Emery, 1901.

1. *Atopula nodifera* EMERY, 1912, Ann. Soc. Ent. Belgique, LVI, p. 104 (♀).

Atopomyrmex nodifer EMERY, 1901, Deutsch. Ent. Zeitschr., p. 115, footnote, fig. (♀, ♀). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 421 (♀).

Type locality: CAMEROON (L. Conradt).

Meranoplina Emery

Calyptomyrmex EMERY

Calyptomyrmex EMERY, 1887, Ann. Mus. Civ. Genova, XXV, p. 472.

Genotype: *Calyptomyrmex beccarii* Emery, 1887.

Subgenus 1. **Calyptomyrmex** EMERY, *sensu stricto*

Subgenotype: same as genotype.

1. **Calyptomyrmex arnoldi** (FOREL) ARNOLD, 1917, Ann. South African Mus., XIV, p. 360 (♀), Pl. VIII, figs. 114, 114a.

Dicroaspis arnoldi FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 115 (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

2. **Calyptomyrmex nummulitica** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 352 (♀).

Type locality: Victoria, CAMEROON (F. Silvestri).

3. **Calyptomyrmex stellatus** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 255, fig. 5 (♀).

Type locality: Gaboon, FRENCH CONGO (F. Faure).

Subgenus 2. **Dicroaspis** EMERY*Dicroaspis* EMERY, 1908, Ann. Soc. Ent. Belgique, LII, p. 184.*Calyptomyrmex* subg. *Dicroaspis* EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 15.Subgenotype: *Dicroaspis cryptocera* Emery, 1908.

4. **Calyptomyrmex (Dicroaspis) claviseta** (SANTSCHI).

Dicroaspis claviseta SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 27 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 362 (♀):

Type locality: Pietermaritzburg, NATAL (I. Trägårdh).

5. **Calyptomyrmex (Dicroaspis) cryptocerus** (EMERY).

Dicroaspis cryptocera EMERY, 1908, Ann. Soc. Ent. Belgique, LII, p. 185, fig. I (♀, ♀). WASMANN, 1918, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VII, p. 145.

Type locality: Stanleyville, BELGIAN CONGO (Kohl).

- 5i. Var. **laevidens** SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 88 (♀).

Type locality: Yambuya, BELGIAN CONGO (J. Bequaert).

6. **Calyptomyrmex (Dicroaspis) foreli** EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 15.

Dicroaspis emeryi FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 262 (♀). SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 311 (♀, ♀). (*nec Calyptomyrmex emeryi* Forel, 1901).

Type locality: Ghinda, ERITREA (K. Escherich; F. Silvestri).

ANYSSINIA: HARAR (Reichensperger).

7. **Calyptomyrmex (Dicroaspis) pusillus** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 256 (♀).

Type locality: Gaboon, FRENCH CONGO (F. Faure).

Meranoplus F. SMITH*Meranoplus* F. SMITH, 1854, Trans. Ent. Soc. Lond., (2) II, p. 224.*Cryptocerus* (part) GUÉRIN, F. SMITH, LOWNE. *Myrmica* (part) JERDON.Genotype: *Cryptocerus bicolor* Guérin, 1845.

1. **Meranoplus bondroiti** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 254, fig. 4 (♀).

Type locality: Uzaga, region of the Great Lakes, BELGIAN CONGO (Gérard).

2. *Meranoplus excisus* ARNOLD, 1914, Proc. Rhodesia Sc. Assoc., XIII, p. 29, Pl., fig. 9 (♂); 1917, Ann. South African Mus., XIV, p. 367 (♀).

Type locality: Estcourt, NATAL (R. C. Wroughton).

3. *Meranoplus magretti* ERN. ANDRÉ, in Magretti, 1884, Ann. Mus. Civ. Genova, XXI, p. 543, fig. A (♂); in Magretti, 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 245, fig. 1a (♂). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 137. MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 201 (♂; var.?). FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 79. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 39 (var.?).

Type locality: Suakin, ANGLO-EGYPTIAN SUDAN (Magretti).

SOUTHERN ABYSSINIA: (Hg). GERMAN EAST AFRICA: Bagamoyo (F. Stuhlmann; var.?).

4. *Meranoplus nanus* ERN. ANDRÉ, 1892, Rev. d'Ent. Caen, XI, p. 55 (♂). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 137. SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 351 (♂, ♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 365 (♂).

Type locality: Gaboon, FRENCH CONGO.

FRENCH GUINEA: Kindia (F. Silvestri).

4a. Subsp. *inermis* (EMERY).

Meranoplus inermis EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 41 (♂), Pl. II, fig. 24; 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 15 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 365 (♂).

Type locality: Makapan, TRANSVAAL (E. Simon).

ERITREA: Asmara (F. Silvestri).

4b. Subsp. *nanior* FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 12 (♂). SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 506. ARNOLD, 1917, Ann. South African Mus., XIV, p. 366 (♂).

Type locality: Mto-ya-Kifaru, GERMAN EAST AFRICA (Katona).

RHODESIA: Victoria Falls (G. Arnold).

4b. Var. *kiboshanus* FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 12 (♂).

Type locality: Kibosho, GERMAN EAST AFRICA (Katona).

4c. Subsp. *soriculus* Wm. M. WHEELER. See p. 184 (♂, ♀, ♂).

Type locality: Avakubi, BELGIAN CONGO (Lang and Chapin).

BELGIAN CONGO: Medje (Lang and Chapin).

5. *Meranoplus peringueyi* EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 365 (♂), Pl. XVII, fig. 12. DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 137. FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 116 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 368 (♂), Pl. VIII, figs. 115, 115a.

Type locality: Cape of Good Hope (L. Péringuey).

CAPE PROVINCE: Willowmore (H. Brauns).

6. *Meranoplus simoni* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 41 (♀), Pl. II, fig. 23. ARNOLD, 1917, Ann. South African Mus., XIV, p. 369 (♀).

Type locality: Vrijburg, CAPE PROVINCE (E. Simon).

6a. Subsp. *nitidiventris* MAYR, 1901, Ann. Naturh. Hofmus. Wien, XIV, p. 26 (♂). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 116 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 370 (♂, ♀, ♂).

Type locality: Bothaville, ORANGE FREE STATE (H. Brauns).

RHODESIA: Bulawayo (G. Arnold).

6a. Var. **springvalensis** ARNOLD, 1917, Ann. South African Mus., XIV, p. 372 (♀).

Type locality: Springvale, SOUTHERN RHODESIA (G. Arnold).

6b. Subsp. **auturalis** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 424 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 370 (♀).

Type locality: NATAL (Haviland).

7. **Meranoplus spininodis** ARNOLD, 1917, Ann. South African Mus., XIV, p. 366 (♀).

Type locality: Hillside, Bulawayo, SOUTHERN RHODESIA (G. Arnold).

Leptothoracini Emery

Macromischoides WM. M. WHEELER

Macromischoides WM. M. WHEELER, 1920, Psyche, XXVII, p. 53.

Genotype: *Macromischa aculeata* Mayr, 1866.

1. **Macromischoides aculeatus** (MAYR). See p. 189.

Macromischa aculeata MAYR, 1866, Sitzb. Ak. Wiss. Wien, LIII, Abt. 1, p. 507 (♀). ERN. ANDRÉ, 1889, Rev. d'Ent. Caen, VIII, p. 224 (♀). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 120. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 3 (♀). MAYR, 1896, Ent. Tidskr., XVII, p. 238. FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 301 (♀). MAYR, 1902, Verh. Zool. Bot. Ges. Wien, p. 292 (♀, ♀, ♂). WASMANN, 1906, Tijdschr. v. Ent., XLVIII, p. 213, Pl. VIII, fig. 3 (nest).

Tetramorium aculeatum EMERY, 1896, Bull. Soc. Ent. France, p. 103; 1908, Ann. Soc. Ent. Belgique, LII, p. 187 (♂). FOREL, 1909, ibid., LIII, p. 59 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 385 (♀, ♀, ♂). FOREL, 1913, Rev. Suisse Zool., XXI, p. 668 (♀, ♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 350 (♀, ♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 366 (♀). H. SCHMITZ, 1915, Deutsch. Ent. Zeitschr., p. 494. FOREL, 1916, Rev. Suisse Zool., XXIV, p. 420 (♀, ♀, ♂). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 393 (♀, ♀). Pl. XXI (nest). STRAND, 1917, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VI, p. 41.

Type locality: GOLD COAST.

FRENCH GUINEA: Los Islands (H. Brauns). SIERRA LEONE: (Mocquerys). CAMEROON: (Sjöstedt); Abo (R. Buchholz); Victoria (F. Silvestri); Ekona (Hintz). FRENCH CONGO: Ogowe (Mocquerys); Brazzaville (A. Weiss). BELGIAN CONGO: St. Gabriel (H. Kohl); Kasai, Kondué; Sankuru (Luja); Lukula (Daniel); Mayombe (de Brier); Duma (Schubotz); Kimuenza (Schultz); Bumba (J. Bequaert); Stanleyville; Avakubi; Bafuka; Medje; Leopoldville; Isangi (Lang and Chapin); Lower Congo (Solon); Lingunda.

1. Var. **major** (FOREL).

Tetramorium aculeatum var. **major** FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 344 (♀); 1916, Rev. Suisse Zool., XXIV, p. 420 (♀, ♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

12. Var. **rubroflavus** (FOREL).

Tetramorium aculeatum var. **rubroflavus** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 420 (♀).

13. Var. **wasmanni** (FOREL). See p. 190.

Macromischa wasmanni FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 300 (♀).

Tetramorium aculeatum var. *wasmanni* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 71 (♂, ♀, ♂). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 385 (♀, ♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 420 (♀, ♀).

Tetramorium aculeatum subsp. *wasmanni* WASMANN, 1912, Zeitschr. Wiss. Zool., CI, p. 104. REICHENSPERGER, 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, p. 125.

Type locality: Leopoldville, BELGIAN CONGO.

FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: Sankuru (Laja); St. Gabriel (Kohl); Zambi; Stanleyville (Lang and Chapin).

1a. Subsp. *andricus* (EMERY).

Tetramorium aculeatum subsp. *andricum* EMERY, 1908, Ann. Soc. Ent. Belgique, LII, p. 187 (♂, ♀, ♂). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 385 (♀, ♀, ♂). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 350 (♀).

Type locality: Stanleyville, BELGIAN CONGO (Kohl).

FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: Kisantu (J. Bequaert).

1a. Var. *gladiator* (SANTSCHI).

Tetramorium aculeatum subsp. *gladiator* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 248 (♀).

Type locality: Congo da Lemba, BELGIAN CONGO (R. Mayné).

2. *Macromischoides africanus* (MAYR).

Macromischa africana MAYR, 1866, Sitzb. Ak. Wiss. Wien, LIII, Abt. 1, p. 507 (♀). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 120. FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 301 (♀). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 131 (♀).

Tetramorium africanum EMERY, 1896, Bull. Soc. Ent. France, p. 103. SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 385 (♀, ♀). FOREL, 1911, Rev. Zool. Afr., I, p. 279 (♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 366 (♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 421 (♀).

Type locality: GOLD COAST.

FRENCH GUINEA: Koniakry (F. Silvestri). NIGERIA: Old Calabar (H. Brauns). CAMEROON: Victoria (F. Silvestri); Mbalmajo to Ekeneli (G. Schwab). FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: Congo da Lemba (R. Mayné); St. Gabriel (Kohl).

Leptothorax MAYR

Leptothorax MAYR, 1855, Verh. Zool. Bot. Ges. Wien, V, p. 431.

Formica (part) FABRICIUS, etc. *Macromischa* EMERY. *Myrmica* (part) NYLANDER, etc. *Stenamma* CURTIS, F. SMITH, PATTON.

Genotype: *Myrmica clypeata* Mayr, 1853.

Subgenus Goniothorax EMERY

Leptothorax subg. *Goniothorax* EMERY, 1896, Bull. Soc. Ent. Italiana, XXVIII, p. 26.

Subgenotype: *Leptothorax vicinus* Mayr, 1887.

1. *Leptothorax (Goniothorax) angulatus* MAYR.

Leptothorax angulatus MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 739 (♂); 1863, ibid., XIII, p. 426. ERN. ANDRÉ, 1883, 'Spec. Hym. Europ. Algérie,' II,

- p. 295 (♀). EMERY, 1891, 'Explor. Tunisie, Fourmis,' p. 5. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 123. FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 233 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 107 (♀, ♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 258 (♀), Pl. v, figs. 56, 56a.

Type locality: Sinai Peninsula (R. v. Frauenfeld).

TUNIS. CENTRAL UGANDA: (C. Alluaud). SOUTHERN RHODESIA: Khami River; Matoppos Hills (G. Arnold).

- 1a. Subsp. **concolor** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 107, fig. 15 (♀, ♀). EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 16 (♀, ♀, ♂).

Type locality: Mombasa, BRITISH EAST AFRICA (Alluaud and Jeannel).

ERITREA: Mayahal (F. Silvestri).

- 1b. Subsp. **ilgi** FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 82 (♀). SANTSCHI, 1912, Bull. Soc. Hist. Nat. Afr. Nord, IV, p. 148 (♀, ♀).

Type locality: SOUTHERN ABYSSINIA (Ilg).

FRENCH SOMALILAND: Obock (Maindron).

2. **Leptothorax (Goniothorax) denticulatus** MAYR.

- Leptothorax denticulatus* MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 5 (♀, ♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 260 (♀, ♀).

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

3. **Leptothorax (Goniothorax) evelynæ** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 423 (♀, ♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

4. **Leptothorax (Goniothorax) grisoni** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 423 (♀, ♂).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

5. **Leptothorax (Goniothorax) innocens** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 423 (♀).

Tetramorium (Leptothorax?) innocens FOREL, 1913, Rev. Zool. Afr., II, p. 317 (♀). BEQUAERT, *ibid.*, p. 424.

Type locality: Elisabethville, BELGIAN CONGO (J. Bequaert).

6. **Leptothorax (Goniothorax) latinodis** MAYR. See p. 1029.
Leptothorax latinodis MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 130 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 259 (♀).

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA (H. Brauns).

Ocymyrmecini Emery

Ocymyrmex EMERY

Ocymyrmex EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 364.

Genotype: *Ocymyrmex barbiger* Emery, 1886.

1. **Ocymyrmex barbiger** EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 364 (♀, ♂), Pl. XVII, figs. 9-11. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 72. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 26. ARNOLD, 1916, Ann. South African Mus., XIV, p. 199 (♀, ♂), Pl. v, fig. 53.

Ocymyrmex barbatus EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 117 (♀).

Type locality: Cape of Good Hope (L. Péringuey).

CAPE PROVINCE: Cape Town (E. Simon); Orange River (G. Arnold). BECHUANALAND: Magalapye (G. Arnold).

2. *Ocymyrmex laticeps* FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 306 (♀); in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 561 (♀).

Type locality: between the Cubango and Cuito Rivers, MOSSAMEDES (Baum and Van der Kellen).

3. *Ocymyrmex picardi* FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 306 (♀); in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 561 (♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 13 (♀). ARNOLD, 1916, Ann. South African Mus., XIV, p. 199 (♀).

Type locality: between the Cubango and Cuito Rivers, MOSSAMEDES (Baum and Van der Kellen).

BECHUANALAND: Severelela; Severelela to Kooa (L. Schultze).

4. *Ocymyrmex robecchii* EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, pp. 114 and 116, fig. (♀). FOREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 352 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 72. EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 178.

Type locality: Webi, ABYSSINIA (Bricchetti-Robecchi).

SOMALILAND: (C. Keller); Erdal (Pavesi). ABYSSINIA: Middle Ganale (V. Bottego).

4a. Subsp. *nitidulus* EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 116 (♀); 1896, *ibid.*, XXXVII, p. 157 (♀, ♂).

Ocymyrmex robecchii var. *nitidulus* EMERY, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 83.

Ocymyrmex nitidulus DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 72.

Type locality: Obbia, SOMALILAND (Bricchetti-Robecchi).

ABYSSINIA: Junction of Webi and Ganale Rivers (V. Bottego); southern Abyssinia (Ilg). SOMALILAND: Lugh (V. Bottego).

5. *Ocymyrmex weitzackeri* EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 116, footnote (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 72. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 26 (♀, ♂). MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 12 (♀).

Ocymyrmex weitzackeri ARNOLD, 1916, Ann. South African Mus., XIV, p. 195 (♀, ♂), Pl. v, fig. 54.

Type locality: Leribe, BASUTOLAND (Weitzacker).

CAPE PROVINCE: Vrijburg; Kimberley (E. Simon). TRANSVAAL: Makapan; Hamman's Kraal (E. Simon). ORANGE FREE STATE. GERMAN EAST AFRICA: Ngarena-Nyuki, Meru Plain (Sjöstedt).

51. Var. *arnoldi* (FOREL).

Ocymyrmex arnoldi FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 138 (♀, ♂).

Ocymyrmex weitzackeri subsp. *arnoldi* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 213 (♀).

Ocymyrmex weitzackeri subsp. *abdominalis* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 16 (♀).

Ocymyrmex weitzackeri var. *arnoldi* ARNOLD, 1916, Ann. South African Mus., XIV, p. 197 (♀, ♂), Pl. v, figs. 52, 52a, and 55.

Type locality: Bulawayo, RHODESIA (G. Arnold).

ZULULAND: Entendweni (I. Trägårdh). SOUTHERN RHODESIA: common (G. Arnold).

52. Var. **foreli** (ARNOLD).

Ocymyrmex weitzackeri var. *foreli* ARNOLD, 1916, Ann. South African Mus., XIV, p. 197 (♀).

Type locality: Redbank, SOUTHERN RHODESIA (G. Arnold).

RHODESIA: Victoria Falls (G. Arnold).

5a. Subsp. **fortior** SANTSCHI, 1911, Rev. Zool. Afr., I, p. 209 (♀).

Type locality: Cucala, BENGUELA (J. Cruchet).

5b. Subsp. **hirsutus** FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 13 (♀).

Ocymyrmex weitzackeri subsp. *hirsutus* ARNOLD, 1916, Ann. South African Mus., XIV, p. 198 (♀).

Type locality: Severelela, BECHUANALAND (L. Schultze).

BECHUANALAND: KOOB (L. Schultze).

5b. Var. **flaviventris** (SANTSCHI).

Ocymyrmex hirsutus var. *flaviventris* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 431 (♀); 1915, Ann. Soc. Ent. France, LXXXIV, p. 254 (♀).

Type locality: Windhoek, GERMAN SOUTHWEST AFRICA.

Originally described, by error, from German East Africa.

5c. Subsp. **transversus** SANTSCHI, 1911, Rev. Zool. Afr., I, p. 209 (♀).

Type locality: Cucala, BENGUELA (J. Cruchet).

5d. Subsp. **wroughtoni** FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 13 (♀).

Ocymyrmex weitzackeri subsp. *wroughtoni* ARNOLD, 1916, Ann. South African Mus., XIV, p. 198 (♀).

Type locality: NATAL (R. C. Wroughton).

5d. Var. **micans** (FOREL).

Ocymyrmex weitzackeri var. *micans* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 12 (♀).

Ocymyrmex weitzackeri subsp. *wroughtoni* var. *micans* ARNOLD, 1916, Ann. South African Mus., XIV, p. 198 (♀).

Type locality: Okahandja, GERMAN SOUTHWEST AFRICA (Peters).

Tetramoriini Emery

Tetramorium MAYR

Tetramorium MAYR, 1855, Verh. Zool. Bot. Ges. Wien, V, p. 423.

Tetragmus ROGER, 1857, Berlin. Ent. Zeitschr., I, p. 10.

ATTA ILLIGER. *Formica* LINNÆUS, etc. *Leptothorax* F. SMITH. *Myrmica* LATREILLE, etc.

Genotype: *Formica cæspitum* Linnæus, 1758.

1. **Tetramorium arnoldi** (FOREL) EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 19, footnote.

Triglyphothrix arnoldi FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 220 (♀).

Tetramorium (Triglyphothrix) arnoldi ARNOLD, 1917, Ann. South African Mus., XIV, p. 338 (♀, ♀), Pl. VII, fig. 93.

Type locality: Shiloh, SOUTHERN RHODESIA (G. Arnold).

RHODESIA: Bulawayo (G. Arnold).

2. **Tetramorium bacchus** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 426 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 309 (♀).

Type locality: NATAL (Haviland).

3. **Tetramorium bequaerti** FOREL, 1913, Rev. Zool. Afr., II, p. 318 (♀). BEQUAERT, *ibid.*, p. 424. ARNOLD, 1917, Ann. South African Mus., XIV, p. 281 (♀). SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, p. 122 (♀).

Type locality: Lake Kabwe, BELGIAN CONGO (J. Bequaert).

3a. Subsp. **bulawayense** FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 119 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 281 (♀, ♀). SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, p. 122 (♀).

Type locality: Bulawayo, RHODESIA (Arnold).

3a₁. Var. **bruni** SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, p. 122 (♀).

Tetramorium bequaerti subsp. *bruni* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 285 (♀).

Type locality: Hillside, SOUTHERN RHODESIA (Arnold).

4. **Tetramorium caespitum** (LINNÆUS) MAYR, 1855, Verh. Zool. Bot. Ges. Wien, V, p. 426 (♀, ♀, ♂).

Formica caespitum LINNÆUS, 1758, 'Syst. Nat.,' Ed. 10, I, p. 581 (♀).

Type locality: "Europa."

EUROPE, tending to become circumpolar.

4₁. Var. **nefassitense** FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 260 (♀).

Type locality: Nefasit, ERITREA (K. Escherich).

4a. Subsp. **altivagans** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 103 (♀).

Type locality: Kinangop, Aberdare Mts., 3100 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

4b. Subsp. **nautarum** SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, p. 156 (♀).

Type locality: ANNOBON ISLAND (Reichensperger).

4c. Subsp. **schultzei** FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 19 (♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 367 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 331 (♀).

Type locality: Kgokong to Kang, BECHUANALAND (L. Schultze).

CAPE PROVINCE: Stellenbosch.

5. **Tetramorium camerunense** MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 129 (♀).

Type locality: CAMEROON (H. Brauns).

5₁. Var. **gegaimi** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 421 (♀, ♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

5₂. Var. **waelbroeki** FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 53 (♀). *Tetramorium camerunense* var. *woelbroeckii* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 367, fig. 29 (♀).

Type locality: Kinshasa, BELGIAN CONGO (Waelbroeck).

SOUTHERN NIGERIA: Lagos (F. Silvestri).

6. **Tetramorium capense** MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidæ, p. 89 (♀); 1870, Verh. Zool. Bot. Ges. Wien, XX, p. 972 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 133. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 36 (♀, ♀,

♂). SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 435; 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 366 (♀, ♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 310 (♀, ♀, ♂).

Tetramorium braunsi FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 119 (♀).

Tetramorium capense var. *braunsi* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 220. ARNOLD, 1917, Ann. South African Mus., XIV, p. 311 (♀).

Type locality: Cape of Good Hope (Novara Expedition).

CAPE PROVINCE: Cape Town (E. Simon); Constantia; Willowmore (H. Brauns); East London.

7. *Tetramorium coloreum* MAYR, 1900, Ent. Tidskr., XXI, p. 273 (♀).

Type locality: Mungo River, CAMEROON (R. Buchholz).

71. Var. *postpetiolatum* SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 88 (♀).

Type locality: Penge, BELGIAN CONGO (J. Bequaert).

8. *Tetramorium doriae* EMERY, 1881, Ann. Mus. Civ. Genova, XIV, p. 530 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 133.

Type locality: Assab, ERITREA (Doria and Beccari).

9. *Tetramorium emeryi* MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 23 (♀, ♀, ♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 300 (♀, ♀, ♂).

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

9a. Subsp. *cristulatum* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 218 (♀, ♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 302 (♀, ♀, ♂).

Type locality: Willowmore, CAPE PROVINCE (H. Brauns).

10. *Tetramorium ericæ* ARNOLD, 1917, Ann. South African Mus., XIV, p. 332 (♀).

Type locality: Hillside, Bulawayo, SOUTHERN RHODESIA (G. Arnold).

11. *Tetramorium frenchi* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 229 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 285 (♀).

Type locality: Krantzklouf, Durban, NATAL (Marley).

12. *Tetramorium gladstonei* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 219 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 284 (♀).

Tetramorium sericeiventris subsp. *gladstonei* SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, p. 125 (♀) (by misprint).

Type locality: Shiloh, SOUTHERN RHODESIA (G. Arnold).

12a. Var. *sepositum* SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, pp. 125 and 131 (♀).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

RHODESIA: Bulawayo (G. Arnold).

13. *Tetramorium gracile* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 81 (♀); 1907, Rev. d'Ent. Caen, XXVI, p. 136 (♂?).

Type locality: SOUTHERN ABYSSINIA (Ilg).

ABYSSINIA: Adis Abeba (de Rothschild).

14. *Tetramorium grandinode* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 308 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 312 (♀).

Type locality: Cape of Good Hope.

14a. Var. *hopense* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 223 (♀, ♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 313 (♀, ♀), Pl. VII, figs. 103, 103a, 103b.

Type locality: Hopetown, CAPE PROVINCE.

15. **Tetramorium grassii** EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 37 (♂, ♀). MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 25 (♂, ♀). FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 226 (♂, ♀).

Tetramorium grassii SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 366 (♂, ♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 304 (♂, ♀), Pl. VII, fig. 105.

Type locality: Cape Town, CAPE PROVINCE (E. Simon).

CAPE PROVINCE: Kimberley (E. Simon); Table Mt., 1500 ft. (G. Arnold); Port Elizabeth (H. Brauns); Constantia.

15₁. Var. **laevigatum** MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 25 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 305 (♂).

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

15₂. Var. **simulans** SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afđ., III, p. 24 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 305 (♂).

Type locality: Richmond, NATAL (I. Trägårdh).

16. **Tetramorium guineense** (FABRICIUS) MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 740; 1863, *ibid.*, XIII, p. 456. ROGER, 1863, 'Verzeich. Formicid.,' p. 27. MAYR, 1870, Verh. Zool. Bot. Ges. Wien, XX, p. 972 (♂). BLACKBURN and KIRBY, 1880, Ent. Monthly Mag., XVII, p. 88. ERN. ANDRÉ, 1883, 'Spec. Hym. Europ. Algérie,' II, pp. 288, 289, 290 (♂, ♀, ♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 133. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 130. FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 135 (♂); 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 260 (♂); 1913, Rev. Zool. Afr., II, p. 319 (♀). BEQUAERT, *ibid.*, p. 425. EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 4 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 306 (♂). See pp. 192 and 1030.

Formica guineensis FABRICIUS, 1793, 'Ent. Syst.,' II, p. 357 (♂). LATRÉILLE, 1802, 'Hist. Nat. Fourmis,' p. 285 (♂). FABRICIUS, 1804, 'Syst. Piez.,' p. 404 (♂). F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 32.

Myrmica bicarinata NYLANDER, 1847, Acta Soc. Sc. Fennicae, II, p. 1061 (♂, ♀). F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 130.

Myrmica cariniceps GUÉRIN, 1852, Rev. Mag. Zool., (2) IV, p. 79 (♀).

Myrmica kollari MAYR, 1853, Verh. Zool. Bot. Ges. Wien, III, p. 283 (♂, ♀, ♂).

Tetramorium kollari MAYR, 1855, *ibid.*, V, p. 425 (♂, ♀, ♂). SICHEL, 1856, Ann. Soc. Ent. France, (3) IV, Bull., p. xxiv. WHITE, 1883, 'Ants and Their Ways,' p. 262 (♂).

Tetramorium cariniceps ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 171.

Myrmica reticulata F. SMITH, 1862, Trans. Ent. Soc. London, (3) I, p. 33 (♂); 1871, Ent. Annual for 1871, p. 60.

Myrmica guineensis ROGER, 1862, Berlin. Ent. Zeitschr., VI, p. 293.

Myrmica (Tetramorium) kollari F. SMITH, 1871, Ent. Annual for 1871, p. 60.

Type locality: Guinea (Isert).

Tropicopolitan. FRENCH GUINEA: Los Islands. ERITREA: Nefasit (F. Silvestri). ABYSSINIA: western part (Ilg); confluent of the Akaki River (de Rothschild). BELGIAN CONGO: Elisabethville (J. Bequaert); Ngayu (Lang and Chapin).

16₁. Var. **erectum** EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 37 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 307 (♂).

Type locality: Vrijburg, CAPE PROVINCE (E. Simon).

16. Var. **phasias** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 226 (♂).
ARNOLD, 1917, Ann. South African Mus., XIV, p. 307 (♂).

Type locality: Durban, NATAL (C. B. Cooper).

- 16a. Subsp. **cristatum** (STITZ) FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 117 (♂).

- Tetramorium guineense* var. *cristatum* STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 144 (♂).

Type locality: Bismarckburg, Togo (Conradt).

RHODESIA: Bulawayo (G. Arnold).

- 16b. Subsp. **medje** WM. M. WHEELER. See p. 192 (♂).

Type locality: Medje, BELGIAN CONGO (Lang and Chapin).

- 16c. Subsp. **peutli** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 419 (♂, ♀).

Type locality: BELGIAN CONGO (H. Kohl).

- 16d. Subsp. **striatum** (STITZ) ARNOLD, 1917, Ann. South African Mus., XIV, p. 308 (♂, ♀, ♂).

- Tetramorium guineense* var. *striatum* STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 144 (♂).

Type locality: Bismarckburg, Togo (Conradt).

SOUTHERN RHODESIA: (G. Arnold).

17. **Tetramorium humile** SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 434 (♂, ♀).

Type locality: Morogoro, GERMAN EAST AFRICA.

18. **Tetramorium intextum** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 104, fig. 14 (♂); 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 506, fig. A (♂).

Type locality: Blue Post Hotel, Kikuyu, 1520 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

18. Var. **cataractae** SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 506, fig. B (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 322 (♂).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

19. **Tetramorium jauresi** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 226 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 296 (♂), Pl. VII, fig. 99.

Type locality: Park Rynie, NATAL (Marley).

20. **Tetramorium joffrei** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 228 (♂, ♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 303 (♂, ♀), Pl. VII, fig. 97.

Type locality: Durban, NATAL (G. Arnold).

20. Var. **alga** ARNOLD, 1917, Ann. South African Mus., XIV, p. 304 (♂, ♀).

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

21. **Tetramorium laevithorax** EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 39 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 286 (♂), Pl. VIII, figs. 110, 110a.

Type locality: Pietermaritzburg, NATAL (Weitzacker).

CAPE PROVINCE: Port Elizabeth (H. Brauns).

22. **Tetramorium lobulicorne** SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 504 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 298 (♂, ♀, ♂).

Type locality: Bulawayo, RHODESIA (G. Arnold).

23. **Tetramorium longicorne** FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 13 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 119 (♀); 1913, Deutsch. Ent. Zeitschr., Beih., p. 219 (♀); 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 226 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 283 (♀), Pl. VII, figs. 96, 96a.

Type locality: Mto-ya-Kifaru, GERMAN EAST AFRICA (Katona).

RHODESIA: Bulawayo (G. Arnold).

24. **Tetramorium longoi** FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 344 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 321 (♀).

Type locality: George, CAPE PROVINCE (H. Brauns).

25. **Tetramorium luteipes** SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 24, footnote.

Tetramorium grassi subsp. *luteipes* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 383, fig. 11 (♀, ♀, ♂).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

26. **Tetramorium meressei** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 422 (♀). See p. 192.

Type locality: BELGIAN CONGO (Kohl).

BELGIAN CONGO: Masaki between Masisi and Walikale (J. Bequaert).

27. **Tetramorium microgyna** SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, p. 132 (♀).

Type locality: NATAL (Haviland).

28. **Tetramorium miserabile** SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, p. 153 (♀).

Type locality: BRITISH EAST AFRICA (Reichensperger).

29. **Tetramorium neuvillei** FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 135 (♀).

Type locality: Diré Daus, ABYSSINIA (de Rothschild).

30. **Tetramorium oculatum** FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 116 (♀).

Type locality: Redbank, RHODESIA (G. Arnold).

31. **Tetramorium pauper** FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 14 (♀).

Type locality: Mto-ya-Kifaru, GERMAN EAST AFRICA (Katona).

31a. Subsp. **nigrum** FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 15 (♀).

Type locality: Mto-ya-Kifaru, GERMAN EAST AFRICA (Katona).

- 31b. Subsp. **transformans** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 104 (♀).

Type locality: Shimoni, BRITISH EAST AFRICA (Alluaud and Jeannel).

32. **Tetramorium popovici** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 230 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 311 (♀, ♀).

Type locality: Table Mt., 1500 ft., CAPE PROVINCE (G. Arnold).

33. **Tetramorium pusillum** EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 38 (♀, ♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 382, fig. 10a (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 323 (♀, ♀).

Type locality: Cape Town, CAPE PROVINCE (E. Simon).

- 33i. Var. **anxium** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 365, fig. 28 (♀, ♀).

Type locality: Camayenne, FRENCH GUINEA (F. Silvestri).

332. Var. **bantouanum** SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 382, fig. 106 (♂, ♂).

Type locality: Mbamu, FRENCH CONGO (A. Weiss).

333. Var. **exoletum** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 366 (♀).

Type locality: Lagos, SOUTHERN NIGERIA (F. Silvestri).

334. Var. **hemisi** Wm. M. WHEELER. See p. 193 (♀).

Type locality: Niangata, BELGIAN CONGO (Lang and Chapin).

335. Var. **mossamedense** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 223. ARNOLD, 1917, Ann. South African Mus., XIV, p. 324 (♀).

Tetramorium caespitum var. *mossamedense* FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 306 (♀); in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 560 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 117.

Type locality: between the Cubango and Cuito Rivers, MOSSAMEDES (Baum and Van der Kellen).

RHODESIA: Bulawayo (G. Arnold).

336. Var. **triste** (SANTSCHI).

Tetramorium pusillum subsp. *mosamedense* var. *tristis* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 285 (♀).

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

33a. Subsp. **ghindanum** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 223. EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 4 (♀).

Tetramorium caespitum subsp. *ghindanum* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 260 (♀).

Type locality: Ghinda, ERITREA (K. Escherich; F. Silvestri).

33b. Subsp. **ladismithense** (FOREL) ARNOLD, 1917, Ann. South African Mus., XIV, p. 325 (♀).

Tetramorium caespitum subsp. *ladismithensis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 117 (♀).

Tetramorium pusillum subsp. *ladysmithensis* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 223.

Type locality: Ladismith, CAPE PROVINCE (H. Brauns).

33c. Subsp. **tablense** (FOREL) ARNOLD, 1917, Ann. South African Mus., XIV, p. 326 (♀, ♀, ♂).

Tetramorium pusillum var. *tablensis* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 223 (♀, ♀, ♂).

Type locality: Table Mt., CAPE PROVINCE (G. Arnold).

34. **Tetramorium pygmaeum** EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 371 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 134. EMERY, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 62 (♀, ♂); 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 17 (♀).

Type locality: Keren, ERITREA (Beccari).

ERITREA: Bizen (A. Fiori); Massaua (Belli).

35. **Tetramorium quadridentatum** STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 144 (♀).

Type locality: Mundame, CAMEROON (Conradt).

36. **Tetramorium quadrispinosum** EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 362 (♀), Pl. XVII, fig. 8. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 134.

FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 80. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 36. WASMANN, 1899, Notes Leyden Mus., XXI, p. 51; 1911, Tijdschr. v. Ent., LIV, p. 201; 1912, Zeitschr. Wiss. Zool., CI, p. 107. FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 226 (♀). REICHENSPERGER, 1915, Med. Göteborgs Mus. Zool. Afd., V, pp. 45 and 73. ARNOLD, 1917, Ann. South African Mus., XIV, p. 277 (♀). SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, p. 122 (♀). See p. 1030.

Type locality: Cape of Good Hope (L. Péringuey).

CAPE PROVINCE: Table Mt. (G. Arnold); Cape Town (E. Simon); Willowmore (H. Brauns). NATAL: (R. C. Wroughton). DELAGOA: (Liengme).

36a. Subsp. *elegans* SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, pp. 122 and 125 (♀).

Type locality: Willowmore, CAPE COLONY (G. Arnold).

36b. Subsp. *eudoxium* (FOREL) SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, p. 122 (♀).

Tetramorium blochmanni subsp. *continentis* var. *eudoxia* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 231 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 280 (♀).

Type locality: Willowmore, CAPE PROVINCE (G. Arnold).

37. *Tetramorium semireticulatum* ARNOLD, 1917, Ann. South African Mus., XIV, p. 319 (♀, ♂).

Type locality: Hillside District, Bulawayo, SOUTHERN RHODESIA (G. Arnold).

38. *Tetramorium sericeiventre* EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 370 (♀). MAGRETTI, 1884, *ibid.*, XXI, p. 539 (♀). ERN. ANDRÉ, 1885, 'Spec. Hym. Europ. Algérie,' II, Suppl., p. 851 (♀) (p. 19 of reprint). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 134. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 37; 1896, Ann. Mus. Civ. Genova, XXXVII, p. 158 (♀); 1909, Deutsch. Ent. Zeitschr., p. 697, fig. 1 (♀). FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 261 (♀, ♀). GESTRO, 1911, Bull. Soc. Ent. Italiana, XLI, (1909), p. 258. FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 266 (♀); 1913, Rev. Zool. Afr., II, p. 319 (♀). BEQUAERT, *ibid.*, p. 424. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 101 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 366 (♀). EMERY, 1915, *ibid.*, X, p. 4 (♀, ♀). SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, pp. 123 and 125 (♀). See pp. 191 and 1030.

Tetramorium sericeiventre MAGRETTI, 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 244 (♀).

Type locality: Sciotel, ERITREA (Beccati).

FRENCH GUINEA: Konakty (F. Silvestri). SIERRA LEONE. NIGERIA: Old Calabar (Bates); Ibadan (F. Silvestri). BELGIAN CONGO: Thysville; Bukama (J. Bequaert); Garamba (Lang and Chapin). GERMAN EAST AFRICA: Moschi, Mt. Kilimandjaro, 1120 m. (Alluaud and Jeannel). SOMALILAND: Matagoi to Lugh (V. Bottego). ERITREA: Asmara (K. Escherich); Nefasit (K. Escherich; F. Silvestri); Ghinda (F. Silvestri). ANGLO-EGYPTIAN SUDAN: El Hefera, Settiti (Magretti).

38a. Var. *arenarium* SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, pp. 123 and 126 (♀).

Type locality: Kairouan, TUNIS (Santschi).

ALGERIA. SENEGAMBIA: Fello (Claveau). ABYSSINIA: Harar (Reichensperger).

38s. Var. **bipartitum** SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, pp. 123 and 126 (§).

Type locality: BRITISH EAST AFRICA (Le Mout).

38s. Var. **debile** FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 80 (§). SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, p. 124 (§).

Type locality: SOUTHERN ABYSSINIA (Ilg).

38s. Var. **gamai** SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, pp. 124 and 128 (§).

Type locality: Gwaai, RHODESIA (G. Arnold).

38s. Var. **hori** SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, pp. 123 and 125 (§).

Tetramorium sericeiventre KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 8 (§).

Type locality: Khartum, ANGLO-EGYPTIAN SUDAN (Karawaiew).

38s. Var. **jasonis** SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, pp. 123 and 127 (§, ♀).

Type locality: Jacquleville, IVORY COAST (Lohier).

IVORY COAST: Dimbroko (Le Mout).

38s. Var. **mundum** SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, pp. 123 and 127 (§).

Type locality: Kakoulima, FRENCH GUINEA (F. Silvestri).

38s. Var. **nigriventre** (STITZ) SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, pp. 123 and 126 (§).

Tetramorium blochmanni var. *nigriventre* STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 144 (§); 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 388 (§). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 366 (§). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 393 (§).

Type locality: Misahöhe, TOGO (Smend).

FRENCH GUINEA: Konakry; Kakoulima (F. Silvestri). FRENCH CONGO: Fort de Possel to Fort Crampel; Fort Crampel (Schubotz). BELGIAN CONGO: Kassenje, western shore of Lake Albert (Schubotz). NIGERIA: Ibadan (F. Silvestri).

38s. Var. **vascoi** SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, pp. 123 and 128 (§).

Type locality: Bulawayo, RHODESIA (G. Arnold).

38s. Subsp. **cinnamomeum** "ARNOLD" SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, p. 124 (§).

Type locality: RHODESIA.

I have not found a form of *Tetramorium* described under the name "*cinnamomeum*" by Arnold.

38s. Subsp. **continentis** (FOREL) SANTSCHI, 1918, Bull. Soc. Hist. Nat. Afr. Nord, IX, pp. 124 and 130 (§). See p. 191.

Tetramorium blochmanni EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 36. FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 311; in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 504; in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 19 (§, ♀, ♂).

Tetramorium blochmanni subsp. *continentis* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 426 (§); 1913, *ibid.*, LVII, p. 118 (§, ♀); 1913, Rev. Zool. Afr., II,

p. 319 (♀). BEQUAERT, *ibid.*, p. 424. ARNOLD, 1917, *Ann. South African Mus.*, XIV, p. 278 (♀, ♀, ♂). WASMANN, 1918, *Tijdschr. v. Ent.*, LX, (1917), p. 396.

Type locality: NATAL (R. C. Wroughton; Haviland).

BECHUANALAND: Kang to Kgokong; Severelela (L. Schultze). GERMAN SOUTHWEST AFRICA: Kubub (L. Schultze). TRANSVAAL: Hamman's Kraal; Hebron (E. Simon). NATAL: (Weitzecker). BASUTOLAND: (Weitzecker). MOSSAMEDES: between the Cubango and Cuito Rivers (Baum and Van der Kellen). RHODESIA: Bulawayo (G. Arnold). BELGIAN CONGO: Elisabethville; Sankisia (J. Bequaert); Zambi (Lang, Chapin and Bequaert).

38b. Var. **georgei** SANTSCHI, 1918, *Bull. Soc. Hist. Nat. Afr. Nord*, IX, pp. 124 and 131 (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

38b. Var. **platonis** SANTSCHI, 1918, *Bull. Soc. Hist. Nat. Afr. Nord*, IX, pp. 124 and 130 (♀).

Type locality: BASUTOLAND (R. C. Wroughton).

38c. Subsp. **femoratum** EMERY, 1895, *Ann. Soc. Ent. France*, LXIV, p. 37 (♀). ARNOLD, 1917, *Ann. South African Mus.*, XIV, p. 294 (♀). SANTSCHI, 1918, *Bull. Soc. Hist. Nat. Afr. Nord*, IX, pp. 124 and 128 (♀).

Type locality: Makapan, TRANSVAAL (E. Simon).

38c. Var. **collutum** SANTSCHI, 1918, *Bull. Soc. Hist. Nat. Afr. Nord*, IX, pp. 124 and 129 (♀).

Type locality: Durban, NATAL (Demarchi).

38c. Var. **transversum** SANTSCHI, 1918, *Bull. Soc. Hist. Nat. Afr. Nord*, IX, pp. 124 and 128 (♀).

Type locality: Pretoria, TRANSVAAL (C. U. Brain).

38d. Subsp. **inversum** FOREL, 1916, *Rev. Suisse Zool.*, XXIV, p. 421 (♀). SANTSCHI, 1918, *Bull. Soc. Hist. Nat. Afr. Nord*, IX, pp. 124 and 129 (♀).

Tetramorium sericeiventris var. *inversa* SANTSCHI, 1910, *Ann. Soc. Ent. France*, LXXVIII, (1909), p. 384 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

FRENCH CONGO: M'Pila (A. Weiss). BELGIAN CONGO: St. Gabriel (Kohl).

38d. Var. **defrictum** SANTSCHI, 1918, *Bull. Soc. Hist. Nat. Afr. Nord*, IX, pp. 124 and 129 (♀).

Type locality: Malindi, RHODESIA (G. Arnold).

NATAL: Durban. BASUTOLAND.

38e. Subsp. **petersi** (FOREL).

Tetramorium blochmanni subsp. *petersi* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 19 (♀). ARNOLD, 1917, *Ann. South African Mus.*, XIV, p. 280 (♀).

Type locality: Okahandja, GERMAN SOUTHWEST AFRICA (Peters).

39. **Tetramorium setigerum** MAYR, 1901, *Ann. Naturh. Hofmus. Wien*, XVI, p. 22 (♀). FOREL, 1910, *Ann. Soc. Ent. Belgique*, LIV, p. 424 (♀, ♀, ♂). ARNOLD, 1917, *Ann. South African Mus.*, XIV, p. 287 (♀, ♀, ♂).

Type locality: Bothaville, ORANGE FREE STATE (H. Brauns).

NATAL: (R. C. Wroughton; Standfuss).

39a. Subsp. **quærens** FOREL, 1914, *Bull. Soc. Vaudoise Sc. Nat.*, (5) L, p. 226 (♀). ARNOLD, 1917, *Ann. South African Mus.*, XIV, p. 289 (♀). See p. 193.

Type locality: Bulawayo, RHODESIA (G. Arnold).

NATAL: Durban (Marley). BELGIAN CONGO: Nisapu (Lang and Chapin).

40. *Tetramorium setuliferum* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 36 (♀). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 19 (♀, ♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 118 (♀). ARNOLD, 1913, Proc. Rhodesia Sc. Assoc., XII, p. 14; 1917, Ann. South African Mus., XIV, p. 289 (♀, ♀, ♂), Pl. VII, figs. 101, 101a.

Tetramorium squamiferum FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 80 (♀) (without description); 1907, Mitt. Naturh. Mus. Hamburg, XXIV, p. 15 (♀) (without description).

Type locality: Vrijburg, CAPE PROVINCE (E. Simon).

NORTH WEST RHODESIA: Mwendwa (H. Dollman, in Coll. Wheeler). SOUTHERN RHODESIA: Bulawayo (G. Arnold). BASUTOLAND: (Weitzcker). CAPE PROVINCE: Mafeking. DELAGOA: (Liengine). ORANGE FREE STATE: (Weitzcker); Bothaville (H. Brauns). BECHUANALAND: Khakhea; Severelela; Kooa to Sekgoma (L. Schultze). GERMAN SOUTHWEST AFRICA: Gobabis (Gentz). NATAL: (R. C. Wroughton).

Specimens collected in Natal by Wroughton and received from Forel as "*T. squamiferum* Emery in litt." agree entirely with Emery's description of *T. setuliferum*.

- 40i. Var. *cuculense* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 356 (♀).

Type locality: Cucala near Caconda, BENGUELA (J. Cruchet).

- 40s. Var. *triptolemium* ARNOLD, 1917, Ann. South African Mus., XIV, p. 292 (♀).

Type locality: not mentioned, probably RHODESIA (G. Arnold).

- 40a. Subsp. *cluna* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 218 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 292 (♀).

Type locality: Willowmore, CAPE PROVINCE (H. Brauns).

- 40b. Subsp. *galosasanum* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 356.

Tetramorium setuliferum var. *galosana* SANTSCHI, 1910, *ibid.*, LXXVIII, (1909), p. 381 (♀, ♀, ♂).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

FRENCH CONGO: M'Bounion; Mindouga; Comba-Ibre (A. Weiss).

41. *Tetramorium simillimum* (F. SMITH) MAYR, 1861, 'Europ. Formicid.,' p. 61 (♀); 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 456; 1870, *ibid.*, XX, p. 972 (♀). ERN. ANDRÉ, 1874, Rev. Mag. Zool., (3) II, p. 191; 1881, Ann. Soc. Ent. France, (6) VII, p. 71 (♀, ♀). FOREL, 1881, Mitth. München. Ent. Ver., V, p. 3. ERN. ANDRÉ, 1883, 'Spec. Hym. Europ. Algérie,' II, pp. 287, 289, 290 (♀, ♀, ♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 134. MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 7. FOREL, 1904, Ann. Soc. Ent. Belgique, XLVIII, p. 422; 1907, Ann. Mus. Nat. Hungarici, V, p. 13 (♀). KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 8 (♀). FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 225. EMERY, 1909, Deutsch. Ent. Zeitschr., p. 696 (♀, ♀, ♂). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 421 (♀, ♀, ♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 326 (♀, ♀, ♂). WASMANN, 1918, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VII, p. 136. See pp. 193 and 1030.

Myrmica simillima F. SMITH, 1851, 'List Brit. Anim. Brit. Mus., VI, Aculeat.,' p. 118 (♀). CURTIS, 1854, Trans. Linn. Soc. London, XXI, p. 216 (♀). F. SMITH, 1855, Trans. Ent. Soc. London, (2) III, p. 129 (♀).

Tetragmus caldarius ROGER, 1857, Berlin. Ent. Zeitschr., I, p. 12 (♀, ♀); 1861, *ibid.*, V, p. 172 (♂); 1862, *ibid.*, VI, p. 297.

Myrmica (Leptothorax) simillima F. SMITH, 1858, 'Cat. Brit. Fossor. Hym.,' p. 31 (♀).

Myrmica caldaria MEINERT, 1860, Danske Vidensk. Selsk. Skrift., naturv. math. Afd., (5) V, p. 334 (♀, ♀, ♂).

Tetragmus simillimus ROGER, 1863, 'Verzeich. Formicid.,' p. 27.

Leptothorax simillimus WHITE, 1883, 'Ants and Their Ways,' p. 266.

Tetramorium (Tetragmus) simillimum SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 106 (♀).

Type locality: Dorsetshire, ENGLAND, in a hothouse (Dale).

Tropicopolitan. CAMEROON: Soppo (v. Rothkirch). BELGIAN CONGO: St. Gabriel (H. Kohl); Stanleyville (Lang and Chapin). GERMAN EAST AFRICA: Kibosho (Katona). BRITISH EAST AFRICA: Shimoni (Alluaud and Jeannel). EGYPT: Ismailia (Biró). ANGLO-EGYPTIAN SUDAN: White Nile, south of Ondurman (I. Trägårdh); Khartum (Karawajew). NATAL: Durban (G. Arnold).

41. Var. **poweri** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 225 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 328 (♀).

Type locality: Kimberley, CAPE PROVINCE (Power).

41a. Var. **shilohense** FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 218 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 329 (♀).

Type locality: Shiloh, SOUTHERN RHODESIA (G. Arnold).

RHODESIA: Bembesi (G. Arnold).

41a. Subsp. **bothae** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 425 (♀, ♀, ♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 331 (♀, ♀, ♂).

Type locality: BASUTOLAND (R. C. Wroughton; Haviland).

NATAL: (R. C. Wroughton).

41b. Subsp. **delagoense** FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 80 (♀, ♀, ♂); 1910, Ann. Soc. Ent. Belgique, LIV, p. 425. ARNOLD, 1917, Ann. South African Mus., XIV, p. 330 (♀, ♀, ♂).

Type locality: DELAGOIA (Liengme).

NATAL: (R. C. Wroughton).

41c. Subsp. **isipingense** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 225 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 329 (♀).

Type locality: Isipingo, NATAL (Marley).

41c. Var. **dumezi** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 422 (♀). See p. 193.

Type locality: St. Gabriel, BELGIAN CONGO (H. Kohl).

BELGIAN CONGO: Thysville (J. Bequaert).

42. **Tetramorium simoni** EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 36 (♀), Pl. II, fig. 22. ARNOLD, 1917, Ann. South African Mus., XIV, p. 295 (♀).

Type locality: Makapan, TRANSVAAL (E. Simon).

43. **Tetramorium simulator** ARNOLD, 1917, Ann. South African Mus., XIV, p. 297 (♀), Pl. VII, fig. 102.

Type locality: Malindi, SOUTHERN RHODESIA (G. Arnold).

44. **Tetramorium solidum** EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 362 (♀, ♀), Pl. XVII, fig. 7. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 135. DIXEY

and LONGSTAFF, 1907, Trans. Ent. Soc. London, p. 333. FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 19 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 292 (♀, ♀).

Type locality: Cape of Good Hope (L. Péringuey).

CAPE PROVINCE: Steinkopf (L. Schultze). NATAL: Ladysmith (Dixey and Longstaff).

44j. Var. **signatum** EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 35 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 118 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 293 (♀), Pl. VII, fig. 98.

Type locality: Matjesfontein, CAPE PROVINCE (E. Simon).

CAPE PROVINCE: Kimberley; Willowmore (H. Brauns).

44a. Subsp. **lugubre** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 425 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 293 (♀).

Type locality: MOSSAMEDES (Picard).

44a₁. Var. **grootense** (FOREL) ARNOLD, 1917, Ann. South African Mus., XIV, p. 294 (♀, ♂).

Tetramorium solidum var. *grootensis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 118 (♀, ♂?).

Type locality: Willowmore, CAPE PROVINCE (H. Brauns).

45. **Tetramorium squaminode** SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 356, fig. 2 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 102 (♀, ♀, ♂).

Type locality: Mt. Kilimanjaro, 3200-3800 m., GERMAN EAST AFRICA (Alluaud and Jeannel).

45a. Subsp. **do** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 224 (♀, ♀, ♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 314 (♀, ♀, ♂).

Type locality: Bulawayo, RHODESIA (G. Arnold).

45a₁. Var. **flaviceps** ARNOLD, 1917, Ann. South African Mus., XIV, p. 316 (♀).

Type locality: Matoppe Hills, SOUTHERN RHODESIA (G. Arnold).

45a₂. Var. **mus** ARNOLD, 1917, Ann. South African Mus., XIV, p. 316 (♀).

Type locality: Hillside, Bulawayo, RHODESIA (G. Arnold).

46. **Tetramorium subcæcum** FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 137 (♀).

Type locality: Toullo, ABYSSINIA (de Rothschild).

46₁. Var. **inscium** FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 218 (♀). EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 17 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 318 (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

RHODESIA: Matoppe Hills (G. Arnold). ERITREA: Nefasit (F. Silvestri).

47. **Tetramorium termitobium** EMERY, 1908, Ann. Soc. Ent. Belgique, LII, p. 186 (♀).

Type locality: Sankuru, BELGIAN CONGO (Luja).

48. **Tetramorium tersum** SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 357, fig. 1 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 102 (♀). Type locality: Naivasha, Rift Valley, BRITISH EAST AFRICA (C. Alluaud).

49. **Tetramorium titus** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 427 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 317 (♀).

Type locality: Mountains of NATAL (R. C. Wroughton).

50. **Tetramorium trægaordhi** SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 23 (♂, ♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 319 (♂, ♀).

Type locality: Stamford Hill, NATAL (I. Trägårdh).

Decamorium FOREL

Tetramorium subg. *Decamorium* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 121.

Genotype: *Tetramorium (Decamorium) decem* Forel, 1913.

1. **Decamorium decem** (FOREL).

Tetramorium (Decamorium) decem FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 121 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 349 (♂, ♀, ♂), Pl. VIII, fig. 111.

Type locality: Redbank, RHODESIA (G. Arnold).

SOUTHERN RHODESIA: Nyamandhloru; Hillside, Bulawayo (G. Arnold); Hartley (H. B. Maufe).

1. Var. **ultor** (FOREL).

Tetramorium (Decamorium) decem var. *ultor* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 217 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 351 (♂).

Type locality: Shiloh, SOUTHERN RHODESIA (G. Arnold).

Xiphomyrmex FOREL

Tetramorium subg. *Xiphomyrmex* FOREL, 1887, Mitth. Schweiz. Ent. Ges., VII, p. 385.

Genotype: *Tetramorium (Xiphomyrmex) kelleri* Forel, 1887.

1. **Xiphomyrmex angulinodis** SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 16, fig. 3a (♂). See p. 194.

Tetramorium (Xiphomyrmex) angulinode SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 385, fig. 12 (♂, ♀, ♂). FOREL, 1911, Rev. Zool. Afr., I, p. 279 (♂); 1913, Ann. Soc. Ent. Belgique, LVII, p. 350 (♂).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

BELGIAN CONGO: Leopoldville (Dubois); Congo da Lemba (R. Mayné); Medje; Irumu (Lang and Chapin). FRENCH CONGO: Samkita (F. Faure).

1. Var. **chloe** SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 17, fig. 3c (♂).

Type locality: Sawmills, SOUTHERN RHODESIA (G. Arnold).

1. Var. **daphnis** SANTSCHI, 1920, Ann. Soc. Ent. Belgique, LX, p. 16, fig. 3b (♂, ♀).

Type locality: Hillside near Bulawayo, RHODESIA (G. Arnold).

2. **Xiphomyrmex edouardi** (FOREL) SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 344 (♀).

Tetramorium (Xiphomyrmex) edouardi FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 82 (♂).

Type locality: SOUTHERN ABYSSINIA (Ilg).

ABYSSINIA: (Reichensperger).

3. **Xiphomyrmex escherichi** (FOREL) EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 4 (♂).

Tetramorium (Xiphomyrmex) escherichi FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 259 (♂, ♀).

Type locality: Ghinda, ERITREA (K. Escherich).

ERITREA: Nefasit (F. Silvestri).

4. *Xiphomyrmex fossulatus* (FOREL).

Tetramorium (*Xiphomyrmex*) *fossulatum* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 428 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 348 (♀), Pl. VII, figs. 107, 109.

Type locality: Willbrook, NATAL (R. C. Wroughton).

5. *Xiphomyrmex humbloti* (FOREL). See p. 1031.

5₁. Var. *pembensis* (FOREL).

Tetramorium (*Xiphomyrmex*) *humbloti* var. *pembensis* FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika (1903-05)', II, p. 83 (♀, ♂); 1907, Ann. Mus. Nat. Hungarici, V, p. 13 (♀, ♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 120 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 106 (♀).

Type locality: Fundu Island, W. Pemba, BRITISH EAST AFRICA (Voeltzkow).

GERMAN EAST AFRICA: Moshi; Arusha-chini; Boma Gombe (Katona). BRITISH EAST AFRICA: Mombasa (Alluaud and Jeannel).

5₂. Var. *victoriensis* (FOREL).

Tetramorium (*Xiphomyrmex*) *humbloti* var. *victoriensis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 120 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 346 (♀, ♀, ♂).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

6. *Xiphomyrmex kivuensis* (STITZ).

Tetramorium (*Xiphomyrmex*) *kivuense* STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentral. Afr. Exp. (1907-08)', III, p. 386, fig. 6 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 106 (♀).

Type locality: Kwidjwi Island in Lake Kivu, BELGIAN CONGO (Schubotz).

BRITISH EAST AFRICA: Naivasha, Rift Valley, 1900 m. (Alluaud and Jeannel).

7. *Xiphomyrmex minusculus* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 369, fig. 32 (♀).

Type locality: Victoria, CAMEROON (F. Silvestri).

8. *Xiphomyrmex muralti* (FOREL) SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 88.

Tetramorium (*Xiphomyrmex*) *muralti* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 429 (♀).

Type locality: CAMEROON (L. v. Muralt).

BELGIAN CONGO: Penge (J. Bequaert).

8₁. Var. *trilineatus* SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 88 (♀, ♀).

Xiphomyrmex muralti SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 368 (♀, ♀).

Type locality: Aburi, GOLD COAST (F. Silvestri).

8₂. Subsp. *flavithorax* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 369, fig. 31 (♀).

Type locality: Aburi, GOLD COAST (F. Silvestri).

9. *Xiphomyrmex occidentalis* SANTSCHI, 1916, Bull. Soc. Ent. France, p. 50, fig. 1 (♀).

Type locality: CAMEROON.

9₂. Subsp. *akengensis* Wm. M. WHEELER. See p. 194 (♀, ♀).

Type locality: Akenge, BELGIAN CONGO (Lang and Chapin).

LIBERIA.

10. **Xiphomyrmex orbiceps** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 367, fig. 30 (♂).

Type locality: Victoria, CAMEROON (F. Silvestri).

GOLD COAST: Aburi (F. Silvestri).

11. **Xiphomyrmex weitzckeri** (EMERY).

Tetramorium (Xiphomyrmex) weitzckeri EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 39 (♂). WASMANN, 1911, Tijdschr. v. Ent., LIV, p. 201. REICHENSPERGER, 1915, Med. Göteborgs Mus. Zool. Afd., V, pp. 40 and 73.

Tetramorium (Xiphomyrmex) weitzckeri FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 428 (♂).

Tetramorium (Xiphomyrmex) weitzckeri FOREL, 1913, Rev. Zool. Afr., II, p. 319 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 345 (♂, ♀).

Tetramorium weitzckeri BEQUAERT, 1913, Rev. Zool. Afr., II, p. 425.

Tetramorium (Xiphomyrmex) weitzckeri SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 24 (♂).

Type locality: Verulam, NATAL (Weitzcker).

BELGIAN CONGO: Elisabethville (J. Bequaert). RHODESIA: Victoria Falls (R. C. Wroughton); Redbank; Bulawayo; Hope Fountain (G. Arnold). NATAL: (Haviland); Durban; Stamford Hill (I. Trägårdh).

Tetramyrma FOREL

Dilobocondyla subg. *Tetramyrma* FOREL, 1912, Rev. Suisse Zool., XX, p. 766.

Tetramyrma FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 122; 1917, Bull. Soc. Vaudoise Sc. Nat., (5) LI, p. 245. ARNOLD, 1917, Ann. South African Mus., XIV, p. 358.

Genotype: *Dilobocondyla (Tetramyrma) braunsi* Forel, 1912.

1. **Tetramyrma braunsi** FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 122 (♀ ergat., ♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 358 (♂, ♀), Pl. VII, fig. 86.

Dilobocondyla (Tetramyrma) braunsi FOREL, 1912, Rev. Suisse Zool., XX, p. 767 (♂).

Type locality: Willowmore, CAPE PROVINCE (H. Brauns).

Rhoptromyrmex MAYR

Rhoptromyrmex MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 18.

Genotype: *Rhoptromyrmex globulinodis* Mayr, 1901.

1. **Rhoptromyrmex arnoldi** SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 503 (♂); 1917, *ibid.*, LXXXV, (1916), p. 286.

Type locality: Victoria Falls, RHODESIA (G. Arnold).

2. **Rhoptromyrmex globulinodis** MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 20 (♂, ♀, ♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 352 (♂, ♀, ♂).

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

2a. Subsp. **alberti** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 419 (♂).

Type locality: BELGIAN CONGO (Kohl).

3. **Rhoptromyrmex opacus** FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 59, footnote (♂). EMERY, *ibid.*, p. 133. FOREL, 1916, Rev. Suisse Zool., XXIV, p. 419 (♂). See p. 195.

Type locality: CAMEROON (Conradt).

BELGIAN CONGO: (Kohl); Thysville (J. Bequaert).

- 3i. Var. **estus** FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 59 (♀, ♂).
Type locality: Lower Congo, in stomach of *Manis temmincki* (Solon).
- 3a. Var. **laeviceps** SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 504 (♀).
Type locality: Boma, BELGIAN CONGO.
4. **Rhoptromyrmex solleri** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 430 (♀).
Type locality: Bissao, PORTUGUESE GUINEA (Soller).
5. **Rhoptromyrmex steini** FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 122 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 357 (♀).
Type locality: Ladismith, CAPE PROVINCE (H. Brauns).
6. **Rhoptromyrmex tessmanni** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 421 (♀).
Type locality: Alen, SPANISH GUINEA (Tessmann).
7. **Rhoptromyrmex transversinodis** MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 22 (♀). BERGROTH, 1903, Wien. Ent. Zeitg., XXII, p. 254. ARNOLD, 1917, Ann. South African Mus., XIV, p. 355 (♀, ♂), Pl. VIII, figs. 112, 112a, 113, 113a. *Rhoptromyrmex transversinodis* "MAYR" WASMANN, 1898, Wien. Ent. Zeitg., XVII, p. 99 (without description).
Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).
ORANGE FREE STATE: Bothaville (H. Brauns). TRANSVAAL: Pretoria (C. U. Brain).

Triglyphothrix FOREL

- Triglyphothrix* FOREL, 1890, Ann. Soc. Ent. Belgique, XXXIV, C. R., p. cvi.
Genotype: *Triglyphothrix walshi* Forel, 1890.
1. **Triglyphothrix areolatus** STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 142, fig. 9 (♀).
Type locality: Bibundi, CAMEROON (Tessmann).
 2. **Triglyphothrix auropunctatus** FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 20 (♀, ♂).
Tetramorium (Triglyphothrix) auropunctatus ARNOLD, 1917, Ann. South African Mus., XIV, p. 334 (♀, ♂).
Type locality: NATAL, 2500 m. (Haviland).
 - 2i. Var. **bulawayensis** (ARNOLD).
Tetramorium (Triglyphothrix) auropunctatus var. *bulawayensis* ARNOLD, 1917, Ann. South African Mus., XIV, p. 335 (♀).
Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).
 - 2a. Var. **fusciventris** FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 116 (♀).
Tetramorium (Triglyphothrix) auropunctatus var. *fusciventris* ARNOLD, 1917, Ann. South African Mus., XIV, p. 336 (♀).
Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).
 - 2b. Var. **pallens** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 424 (♀); 1911, Deutsch. Ent. Zeitschr., p. 311 (♀).
Tetramorium (Triglyphothrix) auropunctatus var. *pallens* ARNOLD, 1917, Ann. South African Mus., XIV, p. 336 (♀).
Type locality: NATAL (Haviland).
 - 2c. Var. **rhodesianus** FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 221 (♀).

Tetramorium (Triglyphothrix) zuiopunctatus var. *rhodesiana* ARNOLD, 1917, Ann. South African Mus., XIV, p. 336 (♀).

Type locality: Bembesi, SOUTHERN RHODESIA (G. Arnold).

3. ***Triglyphothrix constanciae*** (ARNOLD).

Tetramorium (Triglyphothrix) constanciae ARNOLD, 1917, Ann. South African Mus., XIV, p. 339 (♂), Pl. VII, figs. 94, 94a.

Type locality: Matoppo Hills, SOUTHERN RHODESIA (G. Arnold).

4. ***Triglyphothrix desertorum*** FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 20 (♀, ♂).

Tetramorium (Triglyphothrix) desertorum ARNOLD, 1917, Ann. South African Mus., XIV, p. 344 (♀, ♂).

Type locality: Kgokong to Kang, BECHUANALAND (L. Schultze).

5. ***Triglyphothrix emini*** FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 79 (♂).

Type locality: SOUTHERN ABYSSINIA (Ilg).

6. ***Triglyphothrix gabonensis*** ERN. ANDRÉ, 1892, Rev. d'Ent. Caen, XI, p. 53 (♂, ♀, ♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 135. STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 144 (♂). FOREL, 1913, Rev. Suisse Zool., XXI, p. 668 (♂). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 393 (♀, ♂). See p. 197.

Type locality: GABOON, FRENCH CONGO.

CAMEROON: Mundame (Conradt). BELGIAN CONGO: Angu; Libenge; Mon-gumba, on the Ubangi River (Schubotz); Akengé; Niapu; Ngayu; Medje (Lang and Chapin).

6i. Var. ***boulognei*** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 423 (♂).

Type locality: BELGIAN CONGO (H. Kohl).

6j. Var. ***brevispinosus*** STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 144 (♂).

Type locality: Bismarekburg, Togo (Conradt).

6a. Subsp. ***soyauxi*** FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 53 (♀).

Type locality: Sibange, FRENCH CONGO (Soyaux).

7. ***Triglyphothrix hepburni*** (ARNOLD).

Tetramorium (Triglyphothrix) hepburni ARNOLD, 1917, Ann. South African Mus., XIV, p. 340 (♂, ♀, ♂).

Type locality: Hillside, Bulawayo, SOUTHERN RHODESIA (G. Arnold).

SOUTHERN RHODESIA: Matoppo Hills (G. Arnold).

8. ***Triglyphothrix imbellis*** EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 18, fig. 9 (♂).

Type locality: Nefasit, ERITREA (F. Silvestri).

9. ***Triglyphothrix inezulæ*** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 221 (♂, ♂).

Tetramorium (Triglyphothrix) inezulæ ARNOLD, 1917, Ann. South African Mus., XIV, p. 343 (♂, ♂).

Type locality: Beach Bush, Durban, NATAL (G. Arnold).

10. ***Triglyphothrix marleyi*** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 221 (♂).

Tetramorium (Triglyphothrix) marleyi ARNOLD, 1917, Ann. South African Mus., XIV, p. 342 (♂).

Type locality: Krantzklouf, NATAL (Marley).

NATAL: Durban (Marley).

11. **Triglyphothrix marthae** FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 265 (♀).

Type locality: ZANZIBAR.

12. **Triglyphothrix microps** MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 25 (♀, ♀).

Tetramorium (Triglyphothrix) microps ARNOLD, 1917, Ann. South African Mus., XIV, p. 337 (♀, ♀).

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

13. **Triglyphothrix mucidus** FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 71 (♀). See p. 197.

Type locality: Sankuru, BELGIAN CONGO (Laja).

BELGIAN CONGO: Medje; Ngayu; Boyulu (Lang and Chapin).

14. **Triglyphothrix pauper** SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 286 (♀).

Type locality: Umgusa River, Cawston Farm, SOUTHERN RHODESIA (G. Arnold).

15. **Triglyphothrix rothschildi** FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 134 (♀).

Type locality: Harar, ABYSSINIA (de Rothschild).

16. **Triglyphothrix silvestrii** EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 17, fig. 8 (♀).

Type locality: Nefasit, ERITREA (F. Silvestri).

17. **Triglyphothrix striatidens** (EMERY) DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 136. FOREL, 1900, Mitth. Schweiz. Ent. Ges., X, p. 284 (♀); 1901, Mitt. Zool. Mus. Berlin, II, p. 10; 1903, Journ. Bombay Nat. Hist. Soc., XIV, p. 704. BINGHAM, 1903, 'Fauna Brit. India, Hym.,' II, p. 173; 1906, Bull. Misc. Inform. Bot. Gard. Kew, Add. Ser. V, p. 28. DONISTHORPE, 1908, *ibid.*, p. 122. WM. M. WHEELER, 1909, Bull. American Mus. Nat. Hist., XXVI, p. 336. DONISTHORPE, 1915, 'British Ants,' p. 341. FOREL, 1911, '1st Congr. Intern. Entom. Bruxelles (1910),' II, Mém., p. 83; 1912, Mém. Soc. Ent. Belgique, XX, p. 1; 1913, Zool. Jahrb. Abt. Syst., XXXVI, p. 82. WM. M. WHEELER, 1916, Journ. Econ. Ent., IX, pp. 566-569, fig. 39 (♀). See p. 1032.

Tetramorium obesum subsp. *striatidens* EMERY, 1889, Ann. Mus. Civ. Genova, XXVII, p. 501 (♀).

- Triglyphothrix obesum* subsp. *striatidens* EMERY, 1891, 'Explor. Tunisie, Fourmis,' p. 4.

Type locality: Bhamo, BURMA (L. Fea).

This Indian ant is becoming tropicopolitan. TUNIS. SIERRA LEONE.

18. **Triglyphothrix trimeni** EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 40 (♀, ♀). FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 13 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 107 (♀).

Tetramorium (Triglyphothrix) trimeni ARNOLD, 1917, Ann. South African Mus., XIV, p. 337 (♀, ♀).

Type locality: Kimberley, CAPE PROVINCE (E. Simon).

GERMAN EAST AFRICA: Kibosho; Mto-ya-Kifaru (Katona); New Moschi; Mt. Kilimanjaro, 800 m. (Alluaud and Jeannel). TRANSVAAL: Hamman's Kraal (E. Simon).

Ochetomyrmicini Emery

Wasmannia FOREL

Wasmannia FOREL, 1893, Trans. Ent. Soc. London, p. 383.

Type: *Tetramorium* (?) *aeropunctatum* ROGER, 1863.

1. *Wasmannia aeropunctata* (ROGER) FOREL, 1893, Trans. Ent. Soc. London, p. 383 (♂, ♀, ♂).

Tetramorium (?) *aeropunctatum* ROGER, 1863, Berlin. Ent. Zeitschr., VII, p. 182 (♂).

Xiphomyrmex atomum SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 370 (♂).

Wasmannia aeropunctata var. *atoma* SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 504 (♂).

Type locality: CUBA (Gundlach).

Neotropical. FRENCH CONGO: Libreville (F. Silvestri); evidently introduced from South America.

Cataulacini Emery

Cataulacus F. SMITH

Cataulacus F. SMITH, 1854, Trans. Ent. Soc. London, (2) II, p. 225.

Cryptocerus (part) LEPELETIER. *Formica* (part) LATREILLE.

Genotype: *Cataulacus taprobanæ* F. Smith, 1854.

1. *Cataulacus baumi* FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 304 (♂, ♀, ♂); in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 560 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 388 (♂, ♀, ♂).

Type locality: between the Cubango and Cuito Rivers, MOSSAMEDES (Baum and Van der Kellen).

1a. Subsp. *batonga* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 114 (♂). ARNOLD, 1920, Ann. South African Mus., XIV, p. 404 (♂).

Cataulacus baumi var. *batonga* ARNOLD, 1917, *ibid.*, XIV, p. 389 (♂), Pl. VIII, figs. 119 and 119a.

Type locality: RHODESIA (G. Arnold).

1a. Var. *bulawayensis* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 218 (♂, ♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 404 (♂).

Cataulacus baumi var. *bulawayensis* ARNOLD, 1917, *ibid.*, XIV, p. 391 (♂, ♀).

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

2. *Cataulacus bequaerti* FOREL, 1913, Rev. Zool. Afr., II, p. 316 (♂). BEQUAERT, *ibid.*, p. 424.

Type locality: Village Kabanza on the River Lovoi near Kikondja, BELGIAN CONGO (J. Bequaert).

3. *Cataulacus coriaceus* STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 138, fig. 7 (♂).

Type locality: Mundame, CAMEROON (Conradt).

4. *Cataulacus egenus* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 359, fig. (♂). See p. 199 (♂, ♀).

Type locality: Madingu, FRENCH CONGO (P. Zimmermann).

BELGIAN CONGO: Medje (Lang and Chapin).

4a. Subsp. **simplex** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicide,' p. 111, fig. 18 (♂).

Type locality: CENTRAL UGANDA (C. Alluaud).

5. **Cataulacus erinaceus** STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 134, fig. 3 (♂). FOREL, 1913, Rev. Zool. Afr., II, p. 315 (♂); 1913, Ann. Soc. Ent. Belgique, LVII, p. 350 (♀); 1913, Rev. Suisse Zool., XXI, p. 668 (♂); 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 345 (♂); 1916, Rev. Suisse Zool., XXIV, p. 427 (♂, ♀). See p. 198.

Cataulacus princeps "EMERY" FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 71 (♂), without description.

Type locality: Alen, SPANISH GUINEA (Tessmann).

CAMEROON: Mundaine (Conradt). BELGIAN CONGO: Kondué, Kasai; Sankuru (Luja); Batiamponde (H. Kohl); Stanleyville; Risimu (Lang and Chapin).

5a. Var. **crassispina** SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 287 (♂).

Type locality: Goda, FRENCH CONGO (P. Chaleuf).

6. **Cataulacus foveolatus** STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 140 (♂).

Type locality: Uelleberg, SPANISH GUINEA (Tessmann).

7. **Cataulacus guineensis** F. SMITH, 1854, Trans. Ent. Soc. London, (2) II, p. 225 (♂), Pl. XX, fig. 5; 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 196; 1862, Trans. Ent. Soc. London, (3) I, p. 414 (♂). MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 403. ROGER, 1863, 'Verzeich. Formicid.,' p. 39. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 138. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 4 (♂). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 129. SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 390 (♂, ♀). FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 263 (♂). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 373 (♂, ♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 427 (♂, ♀). See p. 199.

Type locality: Tropical West Africa.

LIBERIA: Junk River (H. Brauns). DAHOMEY: Kotonou (F. Silvestri). SOUTHERN NIGERIA: Lagos (F. Silvestri); Old Calabar (Bates). FRENCH CONGO: Ogowe (Mocquerys); Brazzaville (A. Weiss). BELGIAN CONGO: Malomé on the Okiavo (Kohl); Leopoldville to Yumbi; Bolobo; Lukolela to Basoko; Isangi; Stanleyville; Medje; Akenge (Lang and Chapin).

7a. Var. **alenensis** (Stitz) FOREL, 1913, Rev. Suisse Zool., XXI, p. 668 (♂).

Cataulacus sulcatus var. **alenensis** STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 137 (♂).

Type locality: Alen, SPANISH GUINEA (Tessmann).

CAMEROON: (Conradt).

7b. Var. **fernandensis** (Stitz).

Cataulacus sulcatus var. **fernandensis** STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 137 (♂).

Type locality: FERNANDO Po (Zenker).

7c. Var. **sulcatus** (Stitz) FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 263 (♂).

Cataulacus sulcatus STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 136, figs. 4-6 (♂, ♀, ♂).

Cataulacus guineensis FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 421.

Type locality: Yaunde, CAMEROON (Zenker).

SOUTHERN NIGERIA: Old Calabar (Bates).

7a. Subsp. **sulcinodis** EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 563 (♂), Pl. xv, fig. 8.

- Cataulacus sulcinodis* DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 139.
Type locality: Assinie, IVORY COAST (C. Alluaud).
8. *Cataulacus hararicus* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 79 (♀). EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 180 (♀).
Type locality: Harar, SOUTHERN ABYSSINIA (Ilg).
SOMALILAND: Lower Ganale (V. Bottego).
9. *Cataulacus huberi* ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 326 (♀).
DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 138. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 128 (♀). ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 4 (♀).
SANTSCHI, 1910, Ann. Soc. Ent. France, LXXXVIII, (1909), p. 390 (♀).
Type locality: SIERRA LEONE (Mocquerys).
FRENCH CONGO: Ubangi (Dybowski); Ogowe (Mocquerys).
- 9i. Var. *longispinus* STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 139, fig. 8 (♀).
Type locality: Mundame, CAMEROON (Conradt).
- 9a. Subsp. *herteri* FOREL, 1913, Rev. Zool. Afr., II, p. 315 (♀). BEQUAERT, *ibid.*, p. 423.
Type locality: Welgelegen, BELGIAN CONGO (J. Bequaert).
10. *Cataulacus intrudens* (F. SMITH) MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 364. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 138. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 78. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 41 (♀, ♀). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 129 (♀); in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 14. Sjöstedt, 1908, *ibid.*, p. 117. ARNOLD, 1917, Ann. South African Mus., XIV, p. 391 (♀, ♀, ♂).
Meranoplus intrudens F. SMITH, 1876, Trans. Ent. Soc. London, p. 609 (♀, ♀, ♂), Pl. XI, figs. 7 and 7a. EMERY, 1892, Zool. Anzeiger, XV, p. 237.
Type locality: Weenen District, NATAL, in thorns of *Acacia* (J. M. Hutchinson).
DELAGOA: (Liangene). TRANSVAAL: Hamman's Kraal (E. Simon). GERMAN EAST AFRICA: Mombo, Usambara (Sjöstedt).
- 10a. Subsp. *intermedius* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 287 (♀).
Cataulacus intrudens var. *intermedius* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 403 and 404 (♀).
Type locality: Bembesi, RHODESIA (G. Arnold).
- 10b. Subsp. *tristiculus* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 237 (♀).
Type locality: Port Elizabeth, CAPE PROVINCE (T. Reeve).
11. *Cataulacus kohli* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 127, fig. 2 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXXVIII, (1909), p. 389, fig. 14 (♀, ♂).
Type locality: Samlia Falls, N'Gamie River, SIERRA LEONE.
FRENCH CONGO: Brazzaville (A. Weiss).
- 11a. Subsp. *brazzavillensis* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXXVIII, (1909), p. 389, fig. 15 (♀, ♂).
Type locality: Brazzaville, FRENCH CONGO (A. Weiss).
12. *Cataulacus lobatus* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 126, fig. 1 (♀).
Type locality: Warship harbor in CAMEROON (H. Brauns).
13. *Cataulacus marleyi* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 219 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 398 (♀, ♀); 1920, *ibid.*, XIV, p. 404 (♀).

Type locality: Krantz Kloof, NATAL (Marley).

NATAL: Durban (Marley).

14. *Cataulacus micans* (MAYR) FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 219. ARNOLD, 1917, Ann. South African Mus., XIV, p. 393 (♀, ♀, ♂).

Cataulacus rugosus subsp. *micans* MAYR, 1901, Ann. Naturh. Hofmus. Wien, X, p. 27 (♀, ♀, ♂).

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

- 14a. Subsp. *durbanensis* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 219 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 394 (♀, ♀, ♂); 1920, *ibid.*, XIV, p. 404 (♀).

Type locality: Durban, NATAL (G. Arnold).

15. *Cataulacus mocquerysi* ERN. ANDRÉ, 1889, Rev. d'Ent. Caen, VIII, p. 229 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 138.

Type locality: SIERRA LEONE (Mocquerys).

- 15i. Var. *nainei* FOREL, 1918, Bull. Soc. Vaudoise Sc. Nat., (5) LI, (1917), p. 724 (♀).

Type locality: BELGIAN CONGO (Kohl).

16. *Cataulacus otii* (FOREL).

Cataulacus wissemanni subsp. *otii* FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 304 (♀).

Type locality: Durban, NATAL (Haviland).

- 16a. Subsp. *fricatidorsus* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 26 (♀). TRÄGÅRDH, *ibid.*, p. 45. ARNOLD, 1917, Ann. South African Mus., XIV, p. 401 (♀).

Type locality: Dukudu, ZULULAND (I. Trägårdh).

17. *Cataulacus parallelus* F. SMITH, 1854, Trans. Ent. Soc. London, (2) II, p. 226 (♀), Pl. XIX, fig. 6; 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 106; 1862, Trans. Ent. Soc. London, (3) I, p. 414 (♀). MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 403. ROGER, 1863, 'Verzeich. Formicid.,' p. 39. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 138. ARNOLD, 1917, Ann. South African Mus., XIV, p. 402 (♀).

Type locality: Cape of Good Hope.

18. *Cataulacus pilosus* SANTSCHI, 1920, Rev. Zool. Afr., VIII, p. 118 (♀, ♀).

Type locality: Avakubi, BELGIAN CONGO (J. Bequaert).

19. *Cataulacus pullus* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 387, fig. 13 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

- 19i. Var. *orientalis* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 108 (♀).

Type locality: Voi, 600 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

20. *Cataulacus pygmaeus* ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 325 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 139. SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 390 (♀).

Type locality: SIERRA LEONE (Mocquerys).

FRENCH CONGO: Brazzaville (A. Weiss).

- 20i. Var. *bakusensis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 350 (♀, ♂).

Type locality: Bakusu, BELGIAN CONGO.

20j. Var. *chariensis* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 358 (♀).

Type locality: Fort Archambault, MOYEN CHARI (Decorse).

20a. Subsp. *brevisetosus* (FOREL) SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, p. 508.

Cataulacus brevisetosus FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 305 (♀); in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 560 (♀).

Type locality: between the Cubango and Cuito Rivers, MOSSAMEDES (Baum and Van der Kellen).

20b. Subsp. *degener* SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 507 (♀).

Type locality: not given.

20c. Subsp. *difficilis* SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 506 (♀).

Type locality: DAHOMEY (Desanti).

20d. Subsp. *jeanneli* SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), pp. 506, footnote, and 508.

Cataulacus jeanneli SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 108, fig. 16 (♀).

Type locality: Gazi near Mombasa, BRITISH EAST AFRICA (Alluaud and Jeannel).

20e. Subsp. *lujae* (FOREL) SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), pp. 506, footnote, and 508. See p. 199.

Cataulacus lujae FOREL, 1911, Deutsch. Ent. Zeitschr., p. 311 (♀); 1913, Rev. Zool. Afr., II, p. 316 (♀).

Cataulacus brevisetosus subsp. *lujae* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 220 (♀, ♂); 1916, Rev. Suisse Zool., XXIV, p. 427 (♀, ♀, ♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 397 (♀, ♀, ♂); 1917, *ibid.*, XIV, p. 404 (♀).

Type locality: KONDÉ, BELGIAN CONGO (Luja).

BELGIAN CONGO: St. Gabriel (Kohl); Garamba (Lang and Chapin). SOUTHERN RHODESIA: Bulawayo (G. Arnold). CAPE PROVINCE: Kimberley (Power).

20e. Var. *gilviventris* (FOREL).

Cataulacus lujae var. *gilviventris* FOREL, 1913, Rev. Zool. Afr., II, p. 316 (♀). BEQUAERT, *ibid.*, p. 423.

Type locality: Village Kabanza on the Lovoi River near Kikondja, BELGIAN CONGO (J. Bequaert).

20e. Var. *plebejus* SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 508 (♀).

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

20e. Var. *weissi* SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 508.

Cataulacus weissi SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 310 (♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 427 (♀, ♀, ♂).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

BELGIAN CONGO: St. Gabriel (Kohl).

21. *Cataulacus rugosus* (FOREL) MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 129 (♀, ♀). H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 43. ARNOLD, 1917, Ann. South African Mus., XIV, p. 392 (♀, ♀).

Cataulacus intrudens var. *rugosus* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 78 (♀).

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA (A. Möller; Liengme). GERMAN EAST AFRICA: Tanga (H. Brauns). ZANZIBAR: (H. Brauns).

21. Var. **subrugosus** SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 26 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 393 (♀).
Type locality: Junction of the Umfolosi Rivers, ZULULAND (I. Trägårdh).
22. **Cataulacus schoutedeni** SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 248 (♀, ♀).
Type locality: Congo da Lemba, BELGIAN CONGO (R. Mayné).
23. **Cataulacus tardus** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 372, fig. 33 (♀).
Type locality: Mamou, FRENCH GUINEA (F. Silvestri).
24. **Cataulacus trægaordhi** SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 24, fig. 3 (♀, ♀, ♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 399 (♀, ♀, ♂).
Type locality: Dukudu, ZULULAND (I. Trägårdh).
ZULULAND: Umfolosi (I. Trägårdh).
- 24a. Var. **plectroniæ** Wm. M. WHEELER. See p. 199 (♀).
Type locality: Stanleyville, BELGIAN CONGO (Lang, Chapin and Bequaert).
- 24s. Var. **ugandensis** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 110 (♀).
Type locality: Unyoro Province, near Hoima, CENTRAL UGANDA (C. Alluaud).
25. **Cataulacus wissmanni** FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 78 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 401 (♀).
Type locality: MOZAMBIQUE (A. Müller).
- 25a. Subsp. **linearis** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 109, fig. 17 (♀).
Type locality: Voi, Wa-Taita, BRITISH EAST AFRICA (Alluaud and Jeannel).
BRITISH EAST AFRICA: Mbuyuni, Pori (Alluaud and Jeannel).

Dacetonini Emery

Microdaceton SANTSCHI

Microdaceton SANTSCHI, 1913, Bull. Soc. Ent. France, p. 478; 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 33.

Genotype: *Microdaceton exornatum* Santschi, 1913.

1. **Microdaceton exornatum** SANTSCHI, 1913, Bull. Soc. Ent. France, p. 478 (♀); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 33, fig. 8 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 384 (♀).

Type locality: Dukudu, ZULULAND (I. Trägårdh).

Strumigenys F. SMITH

Strumigenys F. SMITH, 1860, Journ. of Entom., I, p. 72.

Labidogenys ROGER, 1862, Berlin. Ent. Zeitschr., VI, p. 249.

Pyramica ROGER, ibid., p. 251.

Genotype: *Strumigenys mandibularis* F. Smith, 1860.

Subgenus 1. **Strumigenys** F. SMITH, *sensu stricto*

Subgenotype: same as genotype.

1. **Strumigenys arnoldi** FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 114 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 376 (♀), Pl. VIII, figs. 117, 117a-d. WASMANN, 1918, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VII, p. 143.

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

2. *Strumigenys havilandi* FOREL, 1905, Mitt. Naturh. Mus. Hamburg, XXII, p. 13, footnote (♀). SANTSCHI, 1913, Bull. Soc. Ent. France, p. 257 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 377 (♀). WASMANN, 1918, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VII, p. 143.

Type locality: NATAL, 5300 ft. (Haviland).

- 2a. Subsp. *marleyi* ARNOLD, 1914, Proc. Rhodesia Sc. Assoc., XIII, p. 31 (♀), Pl., fig. 10; 1917, Ann. South African Mus., XIV, p. 378 (♀).

Type locality: Durban, NATAL (Marley).

3. *Strumigenys irrorata* SANTSCHI, 1913, Bull. Soc. Ent. France, p. 257 (♀); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 29, fig. 5 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 375 (♀).

Type locality: Lake Sibayi, ZULULAND (I. Trägårdh).

4. *Strumigenys reticulata* STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 141 (♀, ♀). SANTSCHI, 1913, Bull. Soc. Ent. France, p. 257 (♀).

Type locality: Bibundi, CAMEROON (Tessmann).

5. *Strumigenys rufobrunnea* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 373 (♀, ♀).

Type locality: Konakry, FRENCH GUINEA (F. Silvestri).

NIGERIA: Olokemeji (F. Silvestri).

6. *Strumigenys stygia* SANTSCHI, 1913, Bull. Soc. Ent. France, p. 257 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 113, fig. 20 (♀). ALLUAUD and JEANNEL, 1914, Arch. Zool. Gén. Exp., LIII, p. 366.

Type locality: Cave A at Shimoni, BRITISH EAST AFRICA (Alluaud and Jeannel).

7. *Strumigenys sulfurea* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 261 (♀).

Type locality: Samkita, FRENCH CONGO (F. Faure).

8. *Strumigenys tragaordhi* SANTSCHI, 1913, Bull. Soc. Ent. France, p. 257 (♀).

Strumigenys tragaordhi SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 28, fig. 4 (♀). ARNOLD, 1917, Ann. South African Mus., XIV, p. 374 (♀). EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 19 (♀, ♀).

Type locality: Pietermaritzburg, NATAL (I. Trägårdh).

ERITREA: Ghinda (F. Silvestri). NATAL: Sweetwaters (I. Trägårdh).

Subgenus 2. *Cephaloxys* F. SMITH

Cephaloxys F. SMITH, 1865, Journ. Proc. Linn. Soc. London, Zool., VIII, p. 76.

Trichoscapa EMERY, 1869, Ann. Accad. Natural. Napoli, (2) II, p. 24.

Strumigenys subg. *Cephaloxys* EMERY, 1916, Bull. Soc. Ent. Italiana, XLVII, p. 204.

Subgenotype: *Cephaloxys capitata* F. Smith, 1865.

9. *Strumigenys (Cephaloxys) alluaudi* SANTSCHI.

Strumigenys alluaudi SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 360 (♀, ♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 111 (♀, ♀). ALLUAUD and JEANNEL, 1914, Arch. Zool. Gén. Exp., LIII, pp. 368 and 375.

Strumigenys (Trichoscapa) alluaudi SANTSCHI, 1913, Bull. Soc. Ent. France, p. 258 (♀).

Type locality: Grotto of Tanga, "Kulumuzi," GERMAN EAST AFRICA (C. Alluaud).

9a. Subsp. **nigeriensis** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 376 (♀).

Type locality: Olokemeji, NIGERIA (F. Silvestri).

10. **Strumigenys (Cephaloxys) biconvexa** SANTSCHI.

Strumigenys (Trichoscapa) biconvexa SANTSCHI, 1913, Bull. Soc. Ent. France, p. 258 (♀).

Strumigenys biconvexa SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 111, fig. 19a (♂, ♀).

Type locality: Likoni, BRITISH EAST AFRICA (Alluaud and Jeannel).

11. **Strumigenys (Cephaloxys) concolor** SANTSCHI.

Strumigenys (Trichoscapa) concolor SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 375 (♂, ♀).

Type locality: Aburi, GOLD COAST (F. Silvestri).

12. **Strumigenys (Cephaloxys) emarginata** MAYR.

Strumigenys emarginata MAYR, 1901, Ann. Naturh. Hofmus. Wien, X, p. 26 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 379 (♂).

Strumigenys (Trichoscapa) emarginata SANTSCHI, 1913, Bull. Soc. Ent. France, p. 257 (♂).

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

13. **Strumigenys (Cephaloxys) escherichi** FOREL.

Strumigenys escherichi FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 261 (♂). *Strumigenys (Trichoscapa) escherichi* SANTSCHI, 1913, Bull. Soc. Ent. France, p. 258 (♂).

Type locality: Ghinda, ERITREA (K. Escherich).

13a. Var. **cliens** FOREL, 1913, Rev. Zool. Afr., II, p. 317 (♂).

Strumigenys escherichi subsp. *cognata* var. *cliens* BEQUAERT, 1913, Rev. Zool. Afr., II, p. 424.

Type locality: Elisabethville, BELGIAN CONGO (J. Bequaert).

13a. Subsp. **boerorum** SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 32 (♂). ARNOLD, 1917, Ann. South African Mus., XIV, p. 383 (♂, ♀).

Strumigenys (Trichoscapa) cognata subsp. *boerorum* SANTSCHI, 1913, Bull. Soc. Ent. France, p. 259 (♂).

Type locality: Stamford Hill, NATAL (I. Trägårdh).

ZULULAND: Dukudu (I. Trägårdh).

13b. Subsp. **cognata** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 376 (♂). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 427 (♂, ♀, ♂). WASMANN, 1918, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VII, p. 144.

Strumigenys cognata SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 362 (♂, ♀); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 32, fig. 7 (♂); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 112, fig. 19b (♂).

Strumigenys (Trichoscapa) cognata SANTSCHI, 1913, Bull. Soc. Ent. France, p. 258 (♂).

Type locality: Cucala, BENGUELA (J. Cruchet).

FRENCH GUINEA: Kindia (F. Silvestri). BELGIAN CONGO: (Kohl).

13b. Var. **fusciventris** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 261.

Strumigenys (Trichoscapa) escherichi subsp. *cognata* var. *obscuriventris* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 376 (♂).

Type locality: Olokemeji, NIGERIA (F. Silvestri).

13c. Subsp. *limbata* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 222 (♀).
ARNOLD, 1917, Ann. South African Mus., XIV, p. 381 (♀, ♀), Pl. VIII, fig. 118.

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

NATAL: Durban (Marley).

14. *Strumigenys* (*Cephaloxys*) *glanduscula* SANTSCHI, 1919, Rev. Zool. Afr., VII, p. 88 (♀).

Type locality: Yambuya, BELGIAN CONGO (J. Bequaert).

15. *Strumigenys* (*Cephaloxys*) *lujae* FOREL.

Strumigenys lujae FOREL, 1902, Allgem. Zeitschr. Ent., VII, p. 294, footnote (♀), Pl. I, fig. 1. WASMANN, 1918, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VII, p. 144.

Strumigenys (*Trichoscapa*) *lujae* SANTSCHI, 1913, Bull. Soc. Ent. France, p. 258 (♀).

Type locality: Morumballe, on the Zambesi (E. Luja).

16. *Strumigenys* (*Cephaloxys*) *maynei* FOREL.

Strumigenys (*Trichoscapa*) *maynei* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 427 (♀, ♀, ♂). WASMANN, 1918, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VII, p. 143.

Type locality: Stanleyville, BELGIAN CONGO (Kohl).

16a. Var. *latiuscula* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 428 (♀, ♀).

Type locality: BELGIAN CONGO (R. Mayné).

17. *Strumigenys* (*Cephaloxys*) *rothkirchi* WASMANN.

Strumigenys rothkirchi WASMANN, 1918, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VII, p. 142 (♀), Pl. II, figs. 9 and 10.

Type locality: Soppo, CAMEROON (v. Rothkirch).

18. *Strumigenys* (*Cephaloxys*) *serrula* SANTSCHI.

Strumigenys lujae var. *serrula* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 390 (♀).

Strumigenys serrula SANTSCHI, 1910, ibid., LXXIX, p. 361 (♀).

Strumigenys (*Trichoscapa*) *serrula* SANTSCHI, 1913, Bull. Soc. Ent. France, p. 258 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

19. *Strumigenys* (*Cephaloxys*) *simoni* EMERY.

Strumigenys simoni EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 42 (♀), Pl. II, fig. 21. ARNOLD, 1917, Ann. South African Mus., XIV, p. 380 (♀).

Strumigenys (*Trichoscapa*) *simoni* SANTSCHI, 1913, Bull. Soc. Ent. France, p. 258 (♀).

Type locality: Makapan, TRANSVAAL (E. Simon).

20. *Strumigenys* (*Cephaloxys*) *transversa* SANTSCHI.

Strumigenys (*Trichoscapa*) *transversa* SANTSCHI, 1913, Bull. Soc. Ent. France, p. 258 (♀); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 31, fig. 6 (♀).

Strumigenys transversa ARNOLD, 1917, Ann. South African Mus., XIV, p. 380 (♀).

Type locality: Pietermaritzburg, NATAL (I. Trägårdh).

Epitritus EMERY

Epitritus EMERY, 1869, Bull. Soc. Ent. Italiana, I, p. 136.

Genotype: *Epitritus argiolus* Emery, 1869.

1. *Epitritus mandibularis* SZABÓ, 1909, Archivum Zoologicum, Budapest, I, p. 1, fig. 2 (♀). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 34 (♀).
ARNOLD, 1917, Ann. South African Mus., XIV, p. 385 (♀, ♀).

Type locality: Mto-ya-Kifaru, GERMAN EAST AFRICA (Katona).

NATAL: Stamford Hill (I. Trägårdh).

2. *Epitritus marginatus* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 114, fig. 21 (♂, ♀). ALLUAUD and JEANNEL, 1914, Arch. Zool. Gén. Exp., LIII, p. 366.

Type locality: Shimoni, BRITISH EAST AFRICA (Alluaud and Jeannel).

DOLICHODERINÆ Forel¹

(Formicinae Mayr, part)

Tapinomini Emery

Iridomyrmex MAYR

Iridomyrmex MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 702. EMERY, 1912, 'Gen. Insect., Dolichoderinæ,' p. 21.

Liometopum (part) MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 704.

Formica (part) F. SMITH, LOWNE, ROGER. *Tapinoma* (part) F. SMITH, ROGER.

Acantholepis (part) LOWNE, 1865, Entomologist, II, p. 332.

Hypoclinea (part) MAYR, 1870, Verh. Zool. Bot. Ges. Wien, XX, p. 953.

Tapinoma subg. *Doleromyrma* FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 28.

Genotype: *Formica purpurea* F. Smith, 1858 = *Formica detecta* F. Smith, 1858.

1. *Iridomyrmex humilis* (MAYR) EMERY, 1888, Zeitschr. Wiss. Zool., XLVI, p. 386, Pl. XXVIII, figs. 17-19. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 169. H. v. IHERING, 1894, Berlin. Ent. Zeitschr., XXXIX, p. 378. TITUS, 1905, Bull. 52 U. S. Dept. Agric. Bur. Ent., pp. 79-84, fig. 7. NEWELL, 1907, Circ. 13 Louisiana State Crop Pest Com., p. 9. MARTINS, 1907, Broteria, VI, p. 101. FOREL, 1908, Verh. Zool. Bot. Ges. Wien, LVIII, p. 395. FOSTER, 1908, Journ. Econ. Ent., I, pp. 289-293. NEWELL, 1908, 2nd Bien. Rep. Louisiana State Crop Pest Com. (1906-07), pp. 7-9, fig. 1; 1908, Journ. Econ. Ent., I, pp. 21-34 and 152. LOUNSBURY, 1909, Rep. Gov. Ent., Cape Town, (1908), pp. 66-68. WOODWORTH, 1908, Circ. 38 California Agric. Exper. Stat., p. 11, figs. NEWELL and ROSENFELD, 1909, Circ. 27 Louisiana State Crop Pest Com., pp. 112-116, fig. 16. NEWELL, 1909, Journ. Econ. Ent., II, pp. 174-192, fig. 4, and pp. 324-332 (♂, ♀, ♂), Pl. VII; 1909, 3rd Bien. Rep. Louisiana State Crop Pest Com. (1908-09), pp. iv-v. WOODWORTH, 1910, Bull. 207 California Agric. Exper. Stat., pp. 53-83, figs. 1-7. LOUNSBURY, 1910, Rep. Gov. Ent., Cape Town, (1909), pp. 90-91. WHEELER, 1910, 'Ants, Their Structure, etc.,' pp. 153-155, 233, and 542. SAJÓ, 1910, Prometheus, XXI, p. 634, fig. NICKELS, 1911, Journ. Econ. Ent., IV, pp. 353-358. NEWELL, 1911, Bull. 142 Texas Agric. Exper. Stat., p. 37. EMERY, 1912, 'Gen. Insect., Dolichoderinæ,' p. 26, Pl. I, figs. 14, 14b. GALLARDO, 1912, Bol. Soc. Physis, I, p. 133; 1913, *ibid.*, I, p. 264. FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 223 (♂). NEWELL and BARBER, 1913, Bull. 122 U. S. Dept. Agric. Bur. Ent., pp. 1-98, fig. 6 and Pl. IV. ARNOLD, 1915, Ann. South African Mus., XIV, p. 145 (♂, ♀, ♂), Pl. IV, figs. 41 and 42. BARBER, 1916, Bull. 377 U. S. Dept. Agric., pp. 1-23. WASMANN, 1917, Ent. Mitt.

¹Raffray (1887, Rev. d'Ent., Caen, p. 21) and Wasmann (1894, 'Verzeichn. Myrmecoph. Termitoph. Arthrop.,' p. 100) mention a *Bothriomyrmex pumicatus* "Rey," from the Cape Colony, evidently a manuscript name. Moreover, no species of the genus *Bothriomyrmex* is at present known from the Ethiopian Region.

Deutsch. Ent. Mus. Berlin, VI, pp. 184-186. C. W. MALLY, 1917, South African Journ. Sci., XIV, pp. 245-247. BARBER, 1920, Farmers' Bull. 1101 U. S. Dept. Agric., pp. 1-11, fig. 1 (♀, ♀, ♂).

Hypoclinea (Iridomyrmex) humilis MAYR, 1868, Ann. Soc. Nat. Modena, III, p. 164 (♀); 1870, Verh. Zool. Bot. Ges. Wien, XX, p. 959 (♀).

Iridomyrmex anceps? BINGHAM, 1906, Trans. Ent. Soc. London, Proc., p. xxv (♀). Type locality: Buenos Aires, ARGENTINA (P. de Strobel).

CAPE PROVINCE: Cape Town (Bingham; G. Arnold; H. Brauns). BASUTOLAND: Maseru (R. M. Stoley). Also in MADEIRA (Lounsbury).

Engramma FOREL

Engramma FOREL, 1905, Ann. Soc. Ent. Belgique, XLIX, p. 180. EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 38.

Tapinoma (part) EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 487.

Genotype: *Engramma lujæ* Forel, 1905.

1. ***Engramma allectum*** STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 394, fig. 9 (♀).

Type locality: Koloka near Angu, BELGIAN CONGO (Schubotz).

2. ***Engramma denticulatum*** Wm. M. WHEELER. See p. 205 (♀).

Type locality: Lukolela to Basoko, BELGIAN CONGO (Lang and Chapin).

BELGIAN CONGO: Masaki between Masisi and Walikale (J. Bequaert).

3. ***Engramma gowdeyi*** Wm. M. WHEELER. See p. 207 (♀).

Type locality: Kampala, UGANDA (C. C. Gowdey).

4. ***Engramma griseopubens*** Wm. M. WHEELER. See p. 206 (♀).

Type locality: Lukolela to Basoko, BELGIAN CONGO (Lang and Chapin).

5. ***Engramma ilgi*** FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 264 (♀, ♂). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 38 (♀, ♂). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 431 (♀).

Type locality: WESTERN ABYSSINIA (Ilg).

BELGIAN CONGO: Makanga (Kohl).

5. **Var. *stygium*** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicide,' p. 117 (♀, ♀, ♂).

Engramma stygium SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 363 (♀, ♂). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 38.

Type locality: Nairobi, BRITISH EAST AFRICA (C. Alluaud).

BRITISH EAST AFRICA: Blue Post Hotel, 1520 m.; Lumbwa, Mau Escarpment, 1891 m.; El Burgon to Ndjoro, 2100 m. (Alluaud and Jeannel). GERMAN EAST AFRICA: Mt. Kilimanjaro, 2740 m. (Alluaud and Jeannel).

6. ***Engramma kohli*** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 429 (♀, ♀). See p. 203 (♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

BELGIAN CONGO: Niapu (Lang and Chapin); Lubutu to Kirundu; Tshopo River near Stanleyville (J. Bequaert).

7. ***Engramma laurenti*** EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 38.

Tapinoma laurenti EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 487 (♀, ♀).

Type locality: BELGIAN CONGO (Laurent).

CAMEROON: (Conradt).

71. Var. **congolense** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 431 (♀, ♀ ?).
Type locality: St. Gabriel, BELGIAN CONGO (Kohl).
BELGIAN CONGO: Bengamisa (Kohl).
8. **Engramma lujæ** FOREL, 1905, Ann. Soc. Ent. Belgique, XLIX, p. 181, fig. (♂, ♂); 1911, Rev. Zool. Afr., I, p. 279 (♂, ♂). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 38. See p. 203 (♀).
Type locality: Kondué, BELGIAN CONGO (Luja).
BELGIAN CONGO: Niapu (Lang and Chapin).
- 8a. Subsp. **wasmanni** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 432 (♀, ♀, ♂).
Type locality: BELGIAN CONGO (Kohl).
9. **Engramma wolff** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 432, fig. 1 (♀).
See p. 204 (♀, ♀, ♂).
Type locality: St. Gabriel, BELGIAN CONGO (Kohl).
BELGIAN CONGO: Akenge; Ngayu; Medje (Lang and Chapin); Walikale to Lubutu (J. Bequaert).
10. **Engramma zimmeri** FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 370 (♀). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 38.
Type locality: Amani, GERMAN EAST AFRICA (Zimmer).
- 10a. Subsp. **okiavoense** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 431 (♀).
Type locality: Okiavo River, near St. Gabriel, BELGIAN CONGO (Kohl).

Tapinoma FÖRSTER

- Tapinoma* FÖRSTER, 1850, 'Hym. Stud.,' I, p. 43. FOREL, 1878, Bull. Soc. Vaudoise Sc. Nat., XV, p. 385. EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 38.
Microgaster DUFUR, 1857, Ann. Soc. Ent. France, (2) V, p. 60. ROGER, 1862; EMERY, 1887.
Formica (part) FABRICIUS, LATREILLE, etc. *Lasius* (part) FABRICIUS, 1804.
Myrmica (part) LEPELETIER, F. SMITH.
Technomyrmex (part) EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 43.
Genotype: *Tapinoma collina* Förster, 1850 = *Formica erratica* Latreille, 1798.

Subgenus 1. *Tapinoma* FÖRSTER, *sensu stricto*

- Subgenotype: same as genotype.
1. **Tapinoma acuminatum** FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 85 (♀, ♀ ?).
Type locality: Fundu Island, W. Pemba, BRITISH EAST AFRICA (Voeltzkow).
2. **Tapinoma arnoldi** FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 139 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 154 (♀, ♀, ♂), Pl. iv, fig. 44. FOREL, 1916, Rev. Suisse Zool., XXIV, p. 429 (♀ ?).
Type locality: Plumtree, SOUTHERN RHODESIA (G. Arnold).
SOUTHERN RHODESIA: Bulawayo (G. Arnold). NATAL: (Haviland).
- 2a. Var. **tectum** (SANTSCHI) EMERY *in litt.*
Tapinoma tenue "FOREL" var. *tectum* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 288 (♀).
Type locality: Durban, NATAL (Marley).
- According to information given by Prof. Forel and kindly communicated by Prof. Emery, "*Tapinoma tenue* Forel" was never described and this name was used by mistake for *T. arnoldi*.

3. **Tapinoma danitschi** FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 345 (♀).

Type locality: Durban, NATAL (C. B. Cooper).

NATAL: Umbilo (L. Bevis).

3. Var. **bevisi** FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 346 (♀).

Type locality: Umbilo, NATAL (L. Bevis).

NATAL: Durban (C. B. Cooper).

4. **Tapinoma gracile** FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 140 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 156 (♀).

Type locality: Matetsi near the Victoria Falls, RHODESIA (G. Arnold).

4. Var. **lugubre** SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 288 (♂, ♂).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

5. **Tapinoma luridum** EMERY, 1908, Ann. Soc. Ent. Belgique, LII, p. 188 (♀); 1912, 'Gen. Insect., Dolichoderinae,' p. 41.

Type locality: Sankuru, BELGIAN CONGO (Luja).

- 5a. Subsp. **connexum** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 118, fig. 22 (♀, ♂).

Type locality: Lumbwa, Mau Escarpment, 1897 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

- 5b. Subsp. **longiceps** Wm. M. WHEELER. See p. 209 (♀).

Type locality: Zambi, BELGIAN CONGO (Lang, Chapin and Bequaert).

6. **Tapinoma luteum** (EMERY) MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru., etc.,' II, 8, p. 18 (♀). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 42. SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 35 (♀). TRÄGÅRDH, *ibid.*, p. 45. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 118 (♀, ♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 152 (♀).

Technomyrmex luteus EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 43 (♀).

Type locality: Makapan, TRANSVAAL (E. Simon).

ZULULAND: Junction of the Umfolosi Rivers (I. Trägårdh). GERMAN EAST AFRICA: Meru Plain (Sjöstedt). BRITISH EAST AFRICA: River Ramisi; Shimoni (Alluaud and Jeannel).

- 6a. Subsp. **emeryi** (FOREL) EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p.

42. ARNOLD, 1915, Ann. South African Mus., XIV, p. 152 (♀, ♀, ♂).

Technomyrmex luteus subsp. **emeryi** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 447 (♀, ♂).

Type locality: NATAL (Haviland).

SOUTHERN RHODESIA: Bulawayo (G. Arnold).

7. **Tapinoma melanocephalum** (FABRICIUS) MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 651. FOREL, 1881, Mitth. München. Ent. Ver., V, p. 3 (♀). MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 359. EMERY, 1887, Ann. Mus. Civ. Genova, XXIV, p. 249 (♀, ♂). BIELUPS, 1887, Trans. Ent. Soc. London, Proc., p. xxvii. EMERY, 1888, Zeitschr. Wiss. Zool., XLVI, p. 392, Pl. XXIX, figs. 37-38. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 165. BINGHAM, 1903, 'Fauna Brit. India, Hym.,' II, p. 304 (♀). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 41. SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 377 (♀, ♀). See p. 1034.

Formica melanocephala FABRICIUS, 1793, Ent. Syst., II, p. 353. COQUEBERT, 1799, 'Illustr. Iconogr. Ins.,' I, p. 25 (♀), Pl. VI, fig. 8. LATREILLE, 1802, 'Hist. Nat. Fourmis.,' p. 269 (♀). F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 46.

- Lasius melanocephalus* FABRICIUS, 1804, 'Syst. Piez.,' p. 417.
Myrmica melanocephala LEPELETIER, 1836, 'Hist. Nat. Ins. Hym.,' I, p. 185.
Formica nana JERDON, 1851, Madras Journ. Litt. Sc., XVII, p. 125 (♀); 1854, Ann. Mag. Nat. Hist., (2) XIII, p. 108 (♀).
Myrmica pellucida F. SMITH, 1858, Journ. Proc. Linn. Soc. London, Zool., II, p. 71 (♀).
Myrmica (Monomarium) pellucida F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 124 (♀).
Formica familiaris F. SMITH, 1860, Journ. Proc. Linn. Soc. London, Zool., IV, Suppl., p. 96 (♀).
Micromyrma melanocephala ROGER, 1862, Berlin. Ent. Zeitschr., VI, p. 258 (♀, ♀).

Type locality: Cayenne, FRENCH GUIANA.

Tropicopolitan. FRENCH GUINEA: Konakry (F. Silvestri).

8. *Tapinoma minimum* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 147 (♀, ♀). H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 39. EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 41. SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 35 (♀). TRÄGÅRDH, *ibid.*, p. 45.

Type locality: Tanga, GERMAN EAST AFRICA (H. Brauns).

ZULULAND: Dukudu (I. Trägårdh).

9. *Tapinoma voeltzkowi* FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 84 (♀, ♂ ?).

Type locality: Fundu Island, W. Pemba, BRITISH EAST AFRICA (Voeltzkow).

GERMAN EAST AFRICA: Malindi (Voeltzkow).

- 9a. Var. *rhodesiae* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 139 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 155 (♀), Pl. IV, fig. 43.

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

Subgenus 2. *Ecphorella* FOREL

- Tapinoma* subg. *Ecphorella* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 65. EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 42.

Subgenotype: *Tapinoma (Ecphorella) wellmani* Forel, 1909.

10. *Tapinoma (Ecphorella) wellmani* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 66 (♀). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 42.

Type locality: BENGUELA (C. Wellman).

Technomyrmex MAYR

- Technomyrmex* MAYR, 1872, Ann. Mus. Civ. Genova, II, p. 147. FOREL, 1878, Bull. Soc. Vaudoise Sc. Nat., XV, p. 380. EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 42.

Tapinoma (part) F. SMITH, MAYR.

Genotype: *Technomyrmex strenuus* Mayr, 1872.

1. *Technomyrmex albipes* (F. SMITH) EMERY, 1888, Zeitschr. Wiss. Zool., XLVI, pp. 392, 394, Pl. XXIX, fig. 49. DALLA TORRE; 1893, 'Cat. Hym.,' VII, p. 166. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 74; 1895, Journ. Bombay Nat. Hist. Soc., IX, p. 466 (♀). BINGHAM, 1903, 'Fauna Brit. India, Hym.,' II, p. 301, fig. 91 (♀). FOREL, 1908, Bull. Soc. Vaudoise Sc. Nat., (5) XLIV, p. 21 (♂ ergatom.) EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 43. See p. 1035.

Formica (Tapinoma) albipes F. SMITH, 1862, Journ. Proc. Linn. Soc. London, Zool., VI, p. 38 (♀).

Tapinoma nigrum MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 703 (♀); 1865, 'Reise Novara, Zool., II, Formicidae,' p. 62 (♀); 1867, Tijdschr. v. Ent., X, p. 78 (♀).

Tapinoma albitarse MOTSCHOUJSKY, 1863, Bull. Soc. Nat. Moscou, XXXVI, 3, p. 14 (♀, ♀).

Tapinoma albipes MAYR, 1876, Journ. Mus. Godeffroy, XII, p. 83.

Type locality: Tondano, CELEBES (A. R. Wallace).

DELAGOA: (Liengme).

1. Var. *brunneipes* FOREL, 1895, Journ. Bombay Nat. Hist. Soc., IX, p. 466 (♀). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 43.

Technomyrmex albipes var. *brunneipes* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 391.

Type locality: INDIA; not otherwise designated.

FRENCH CONGO: Mbamu (A. Weiss).

1a. Subsp. *foreli* EMERY, 1893, Ann. Soc. Ent. France, LXII, p. 249 (♀); 1895, *ibid.*, LXIV, p. 43. WASMANN, 1899, Notes Leyden Mus., XXI, p. 51. FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 446 (♀, ♀, ♂). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 43. FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 140 (♀); 1913, Deutsch. Ent. Zeitschr., Beih., p. 223 (♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 377 (♀); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 35 (♀). ARNOLD, 1915, Ann. South African Mus., XIV, p. 150 (♀, ♀, ♂), Pl. iv, fig. 40. REICHENSPERGER, 1915, Med. Göteborgs Mus. Zool. Afd., V, pp. 45 and 73. SKAIFE, 1921, Trans. Roy. Soc. South Africa, IX, 3, p. 221. See p. 1035.

Type locality: MADAGASCAR.

ANGOLA: St. Paul de Loanda (F. Silvestri). ORANGE FREE STATE: Bloemfontein (E. Simon). NATAL: (R. C. Wroughton; L. v. Muralt; Haviland); Durban (G. Arnold); Mountain Rise near Pietermaritzburg (Skaife). ZULULAND: Dukudu (I. Trägårdh). CAPE PROVINCE: Willowmore (H. Brauns); Vrijburg (E. Simon).

1a. Var. *affinis* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 119 (♀).

Type locality: Naivasha, Rift Valley, 1900 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

BRITISH EAST AFRICA: Molo, Mau Escarpment, 2080 m. (Alluaud and Jeannel).

2. *Technomyrmex andrei* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 488, fig. (♀, ♀); 1912, 'Gen. Insect., Dolichoderinae,' p. 43.

Technomyrmex mayri ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 3 (♀) (*nec* Forel).

Type locality: Ogowe, FRENCH CONGO (Mocquerys).

CAMEROON: (Conradt). BELGIAN CONGO: Kinshasa (Waelbroeck).

2. Var. *camerunensis* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 489 (♀); 1912, 'Gen. Insect., Dolichoderinae,' p. 43.

Type locality: CAMEROON.

2. Var. *schereri* FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 283 (♀). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 43.

Type locality: Cape Mount, LIBERIA (Scherer).

3. **Technomyrmex arnoldinus** FOREL, 1913, *Deutsch. Ent. Zeitschr.*, Beih., p. 222 (§). ARNOLD, 1915, *Ann. South African Mus.*, XIV, p. 148 (§), Pl. iv, figs. 39, 39a.

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

NATAL: Durban (C. B. Cooper).

4. **Technomyrmex hypoclinoides** SANTSCHI, 1919, *Rev. Zool. Afr.*, VII, p. 89 (§).

Type locality: Avakubi, BELGIAN CONGO (J. Bequaert).

5. **Technomyrmex mœrens** SANTSCHI, 1913, *Ann. Soc. Ent. Belgique*, LVII, p. 312, fig. 4 (§); 1914, *Boll. Lab. Zool. Gen. Agrar. Portici*, VIII, p. 377 (§).

Type locality: Mbamu, FRENCH CONGO (A. Weiss).

FRENCH GUINEA: Kakoulima (F. Silvestri).

6. **Technomyrmex nigriventris** SANTSCHI, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 22, footnote (§); 1910, *Ann. Soc. Ent. France*, LXXVIII, (1909), p. 391 (§). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 44. ARNOLD, 1915, *Ann. South African Mus.*, XIV, p. 149 (§). See p. 210 (§).

Technomyrmex nigriventris "SANTSCHI" FOREL, 1909, *Ann. Soc. Ent. Belgique*, LIII, p. 54 (♀, ♂), without description.

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

BELGIAN CONGO: Kinshasa (Waelbroeck); Thysville (J. Bequaert).

6a. Subsp. **albinasis** FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 22 (§). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 44. ARNOLD, 1915, *Ann. South African Mus.*, XIV, p. 150 (§).

Type locality: Table Mt. near Cape Town, CAPE PROVINCE (L. Schultze).

7. **Technomyrmex pilipes** EMERY, 1899, *Ann. Soc. Ent. Belgique*, XLIII, p. 490 (♀). FOREL, 1901, *Mitt. Naturh. Mus. Hamburg*, XVIII, p. 65 (♀). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 44 (♀).

Type locality: CAMEROON (Conradt).

FRENCH CONGO: (Dinklage).

8. **Technomyrmex schoutedeni** FOREL, 1910, *Ann. Soc. Ent. Belgique*, LIV, p. 447 (§, ♀, ♂). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 44.

Type locality: Bena Dibebe, BELGIAN CONGO (Luja).

9. **Technomyrmex semiruber** EMERY, 1899, *Ann. Soc. Ent. Belgique*, XLIII, p. 489 (§).

Type locality: CAMEROON (Conradt).

Semonius FOREL

Semonius FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 21. EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 44.

Genotype: *Semonius schultzei* Forel, 1910.

1. **Semonius schultzei** FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 21 (§). EMERY, 1912, 'Gen. Insect., Dolichoderinae,' p. 44. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicide,' p. 120, figs. 23 and 24 (§, ♂). ARNOLD, 1915, *Ann. South African Mus.*, XIV, p. 157 (§, ♀, ♂), Pl. iv, fig. 45. FOREL, 1916, *Rev. Suisse Zool.*, XXIV, p. 429 (§).

Semonius schultzei FOREL, 1913, *Ann. Soc. Ent. Belgique*, LVII, p. 139 (§).

Type locality: Khakhea, BECHUANALAND (L. Schultze).

BELGIAN CONGO: St. Gabriel (Kohl). SOUTHERN RHODESIA: Redbank (G. Arnold). NATAL: Durban (Marley). GERMAN EAST AFRICA: Kilema, 1440 m., Mt. Kilimanjaro (Alluaud and Jeannel).

FORMICINÆ Lepeletier

(Formicinæ Mayr, part; Camponotinae Forel)

Santschiellini Forel

Santschiella FOREL

Santschiella FOREL, 1916, Rev. Suisse Zool., XXIV, p. 434.

Genotype: *Santschiella kohli* Forel, 1916.

1. *Santschiella kohli* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 435, figs. 2-4 (♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

Plagiolepidini Forel

Acropyga ROGER

Acropyga ROGER, 1862, Berlin. Ent. Zeitschr., VI, p. 242.

Plagiolepis (part) MAYR.

Genotype: *Acropyga acutiventris* Roger, 1862.

1. *Acropyga silvestrii* EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 21, fig. 11 (♀).

Type locality: Ghinda, ERITREA (F. Silvestri).

Plagiolepis MAYR

Plagiolepis MAYR, 1861, 'Europ. Formicid.', p. 42.

Formica (part) LATREILLE. *Prelepis* (part) MAYR. *Tapinoma* (part) SCHENCK, MAYR.

Genotype: *Formica pygmaea* Latreille, 1798.

Subgenus 1. *Plagiolepis* MAYR, *sensu stricto*

Subgenotype: same as genotype.

1. *Plagiolepis alluaudi* EMERY, 1894, Ann. Soc. Ent. France, LXIII, p. 71 (♀). FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 30 (♀, ♀). DONISTHORPE, 1908, Bull. Misc. Inform. Bot. Gard. Kew, III, p. 122; 1908, Trans. Leicester Lit. Phil. Soc., XII, p. 229; 1915, 'British Ants,' p. 343. See p. 1035.

Plagiolepis flavidula ERN. ANDRÉ, 1881, 'Spec. Hym. Europ. Algérie,' II, p. 208

- (♀). E. SAUNDERS, 1896, 'Hym. Acul. British Islands,' p. 26 (♀) (*nec* Roger).

Type locality: La Misère, Mahé, SEYCHELLES (C. Alluaud).

Introduced in greenhouses in Europe and apparently spreading.

GERMAN EAST AFRICA: Arusha-chini (Katona).

2. *Plagiolepis brunni* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 148 (♀). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 449 (♀); 1913, *ibid.*, LVII, p. 140 (♀). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 35 (♀, ♂); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 377 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicide,' p. 121 (♀).

Plagiolepis brunni FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 248 (♀, ♀, ♂).

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA (H. Brauns).

FRENCH GUINEA: Maniou (F. Silvestri). BRITISH EAST AFRICA: Ramisi River; Shimoni (Alluaud and Jeannel). RHODESIA: Bulawayo; Springvale (G. Arnold). ZULULAND: Umfolosi (I. Trägårdh). NATAL: Richmond (I. Trägårdh); Durban (Haviland; G. Arnold).

2. Var. *nilotica* MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 7 (♀). EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 19 (♀).

Plagiolepis brunni subsp. *nilotica* FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 30 (♀).

Type locality: Kaka, White Nile, ANGLO-EGYPTIAN SUDAN (I. Trägårdh).

ERITREA: Nefasit (F. Silvestri). GERMAN EAST AFRICA: Kibosho (Katons).

2a. Var. *pubescens* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 140 (♀). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 35 (♀). TRÄGÅRDH, *ibid.*, p. 45.

Type locality: Plumtree, SOUTHERN RHODESIA (G. Arnold).

ZULULAND: Umfolosi (I. Trägårdh).

3. *Plagiolepis capensis* MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidae, p. 55 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 172. FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 24 (♀).

Type locality: Cape of Good Hope (Novara Expedition).

CAPE PROVINCE: Kamaggas (L. Schultze).

4. *Plagiolepis exigua* FOREL, 1894, Journ. Bombay Nat. Hist. Soc., VIII, pp. 415 and 417 (♀, ♀). BINGHAM, 1903, 'Fauna Brit. India, Hym.,' II, p. 323. FOREL, 1913, Rev. Zool. Afr., II, p. 338 (♀). BEQUAERT, *ibid.*, p. 429. FOREL, 1916, Rev. Suisse Zool., XXIV, p. 437 (♀). See p. 1035.

Type locality: Poona, INDIA (Wroughton).

BELGIAN CONGO: (Kohl); Lake Kabwe (J. Bequaert).

4a. Subsp. *abyssinica* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 73 (♀); 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 266 (♀).

Type locality: SOUTHERN ABYSSINIA (Ilg).

ERITREA: (K. Escherich).

5. *Plagiolepis funicularis* SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 344 (♀).

Type locality: NATAL, 3500 ft. (Haviland).

6. *Plagiolepis fuscula* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 45 (♀). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 449 (♀). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 36 (♀).

Type locality: Cape Town, CAPE PROVINCE (E. Simon).

NATAL: (Haviland); Stamford Hill (I. Trägårdh).

7. *Plagiolepis mediorufa* (FOREL). See p. 213.

Plagiolepis pygmaea var. *mediorufa* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 437 (♀).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

BELGIAN CONGO: Stanleyville (Lang and Chapin).

8. *Plagiolepis pygmaea* (LATREILLE) MAYR, 1861, 'Europ. Formicid.,' p. 43 (♀, ♀, ♂). SCHENCK, 1861, Jahrb. Ver. Naturk. Nassau, XVI, p. 200 (♀, ♀, ♂); 1864, Progr. Gymnas. Weilburg, p. 13 (♀, ♀, ♂). FOREL, 1874, Nouv. Mém. Soc. Helvét. Sci. Nat. Zurich, XXVI, pp. 45 and 338 (♀, ♀, ♂). ERN. ANDRÉ, 1874,

Rev. Mag. Zool., (3) II, p. 179 (♀, ♂). MAYR, in Fedtschenko, 1877, 'Voy. Turkestan, Formicid.,' p. 5. ERN. ANDRÉ, 1882, 'Spec. Hym. Europ. Algérie,' II, pp. 208, 209 (♀, ♂), Pls. VI and VII. LUBBOCK, 1882, 'Ants, Bees and Wasps,' Ed. 5, p. 11. FOREL, 1888, Berlin. Ent. Zeitschr., XXII, p. 265. BRYANT, 1890, 'Fourmis France Centr.,' p. 56. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 173. FOREL, 1902, Ann. Soc. Ent. Belgique, XLVI, p. 155 (♀). SANTSCHI, 1908, Ann. Soc. Ent. France, LXXVII, p. 517. FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 224 (♀, ♂).

Formica pygmaea LATREILLE, 1798, 'Ess. Hist. Fourmis France,' p. 45 (♀, ♂); 1802, 'Hist. Nat. Fourmis,' p. 183 (♀). LEPELETIER, 1836, 'Hist. Nat. Ins. Hym.,' I, p. 209 (♀, ♂).

Tapinoma pygmaea SCHENCK, 1852, Jahrb. Ver. Naturk. Nassau, VIII, pp. 68, 130 (♀, ♂); 1853, Stettin. Ent. Zeitg., XIV, p. 185 (♀, ♂). F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 56.

Tapinoma pygmaeum MAYR, 1855, Verh. Zool. Bot. Ges. Wien, V, p. 375 (♀, ♂); 1856, Progr. Realsch. Pest, p. 15 (♀). GREDLER, 1858, VIII. Progr. Gymnas. Bozen, p. 15.

Formica (Tapinoma) pygmaea NYLANDER, 1856, Ann. Sc. Nat. Zool., (4) V, p. 72 (♀, ♂), Pl. III, figs. 6 and 22.

Type locality: FRANCE.

SOUTHERN EUROPE, NORTH AFRICA, ASIA MINOR, CANARIES. RHODESIA: Bulawayo; Bembesi (G. Arnold).

S. Var. *intermedia* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 45 (♀, ♂). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 448 (♀, ♂); 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 248 (♀).

Type locality: Bloemfontein, ORANGE FREE STATE (E. Simon).

CAPE PROVINCE: Cape Town; Vrijburg (E. Simon). BASUTOLAND: (R. C. Wroughton). NATAL: (R. C. Wroughton); Isipingo (C. B. Cooper).

S. Var. *punctum* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 448 (♀, ♂). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 35 (♀, ♂). FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 248 (♀); 1916, Rev. Suisse Zool., XXIV, p. 437 (♀).

Type locality: Mountains of NATAL (R. C. Wroughton).

NATAL: Richmond (I. Trägårdh); Durban. BASUTOLAND: (R. C. Wroughton). BELGIAN CONGO: St. Gabriel (Kohl).

Subgenus 2. *Anacantholepis* SANTSCHI

Plagiolepis subg. *Anacantholepis* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 36.

Subgenotype: *Plagiolepis (Anacantholepis) decora* Santschi, 1914.

9. *Plagiolepis (Anacantholepis) bothæ* FOREL.

Plagiolepis bothæ FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 86 (♀, ♂). Type locality: Fundu Island, W. Pemba, BRITISH EAST AFRICA (Voeltzkow).

10. *Plagiolepis (Anacantholepis) decora* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 36, fig. 9 (♀).

Type locality: Stamford Hill, NATAL (I. Trägårdh).

11. *Plagiolepis (Anacantholepis) deweti* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 249.

Plagiolepis deweti FOREL, 1904, Ann. Soc. Ent. Belgique, XLVIII, p. 423 (♀, ♀).

Type locality: Mountains of NATAL (R. C. Wroughton).

CAPE PROVINCE: Cape of Good Hope (Burchell); King William's Town (R. Godfrey).

12. *Plagiolepis (Anacantholepis) jouberti* FOREL. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 122 (♀).

Plagiolepis jouberti FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 23 (♀).

Type locality: Kainaggas, CAPE PROVINCE (L. Schultze).

BRITISH EAST AFRICA: Shimon (Alluaud and Jeannel).

13. *Plagiolepis (Anacantholepis) pictipes* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 122, fig. 25 (♀).

Type locality: Mbuyuni, Pori, 1110 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

14. *Plagiolepis (Anacantholepis) vanderkelleni* FOREL.

Plagiolepis vanderkelleni FOREL, 1894, Mitth. Schweiz. Ent. Ges., X, p. 310 (♀); in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 564 (♀).

Type locality: between the Cubango and Cuito Rivers, MOSSAMEDES (Baum and Van der Kellen).

14. Var. *tricolor* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 24 (♀).

Type locality: Kooa to Sekgoma, BECHUANALAND (L. Schultze).

14a. Subsp. *polita* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 37, fig. 10 (♀).

Type locality: Stamford Hill, NATAL (I. Trägårdh).

Subgenus 3. *Anoplolepis* SANTSCHI

Plagiolepis subg. *Anoplolepis* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 123.

Subgenotype: *Formica longipes* Jerdon, 1851.

15. *Plagiolepis (Anoplolepis) braunsi* FOREL.

Plagiolepis braunsi FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 141 (♀).

Plagiolepis (Anoplolepis) braunsi SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 123.

Type locality: Willowmore, CAPE PROVINCE (H. Brauns).

16. *Plagiolepis (Anoplolepis) carinata* EMERY. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 123.

Plagiolepis carinata EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 491 (♀, ♀). FOREL, 1907, Mitt. Naturh. Mus. Hamburg, XXIV, p. 15 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 391 (♀, ♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 354 (♀).

Type locality: CAMEROON (Conradt).

FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: Mayombe (Deleval).

17. *Plagiolepis (Anoplolepis) custodiens* (F. SMITH) SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 123 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 377 (♀); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 38 (♀, ♀, ♂). See p. 213.

Formica custodiens F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 33 (♂). ROGER, 1863, 'Verzeich. Formicid.,' p. 15. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 414.

Camponotus hendecarthrus ROGER, 1863, Berlin. Ent. Zeitschr., VII, p. 128 (♂, ♀); 1863, 'Verzeich. Formicid.,' p. 2. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 458.

Formica berthoudi FOREL, 1876, Bull. Soc. Vaudoise Sc. Nat., (2) XIV, p. 33 (♀).

Plagiolepis custodiens MAYR, 1866, Sitzb. Ak. Wiss. Wien, LIII, Abt. 1, p. 493.

FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 215 (♀). EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 117 (♀). FOREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 350 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 172. MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 197. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 73. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 43. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 148 (♂, ♀, ♂). WASMANN, 1896, Notes Leyden Mus., XVIII, p. 75. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 39. EMERY, 1899, Ann. Soc. Ent. Belgique, XLV, p. 491 (♂, ♀). WASMANN, 1900, Illustr. Zeitschr. Ent., V, p. 103; 1904, Notes Leyden Mus., XXV, pp. 15, 73, and 75. ZAVATTARI, 1907, Boll. Mus. Zool. Anat. Comp. Torino, XXII, No. 550, p. 2 (♀). KOLBE, 1907, Ann. Soc. Ent. Belgique, LI, p. 364. FOREL, 1909, *ibid.*, LIII, p. 54 (♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 23 (♂, ♀, ♂). WASMANN, 1911, Tijdschr. v. Ent., LIV, p. 202; 1912, Zeitschr. Wiss. Zool., CI, p. 106. TRÄGÄRDH, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 46. REICHENSPERGER, 1915, *ibid.*, V, p. 18. WASMANN, 1917, Ent. Mitt. Deutsch. Ent. Mus. Berlin, VI, p. 185; 1920, Tijdschr. v. Ent., LXII, (1919), pp. 119 and 120. LAMBORN, 1920, Trans. Ent. Soc. London, (1919), Proc., p. lvi.

Type locality: Port Natal, NATAL.

BELGIAN CONGO: Banana (Busschodts; Lang and Chapin). ANGOLA: St. Paul de Loanda (F. Silvestri); San Antonio (Lang and Chapin). RHODESIA: Kazungula (Jallá). GERMAN SOUTHWEST AFRICA: (Lübbert); Okahandja (Dinter). BECHUANALAND: Lehututu to Kang; Khakhea to Kang; Kooa to Sekgoma (L. Schultze). CAPE PROVINCE: Kimberley; Vrijburg (E. Simon); Willowmore (H. Brauns). NATAL: Pietermaritzburg (I. Trägårdh); Malvern (O'Neil); Durban. BASUTOLAND: Lessouto (P. Berthoud). ORANGE FREE STATE: Bothaville (H. Brauns). TRANSVAAL: Lydenburg (Wilms); Makapan; Pretoria; Hamman's Kraal (E. Simon); Valdezia (P. Berthoud). ZULULAND: Umfolosi (I. Trägårdh). PORTUGUESE EAST AFRICA: Delagoa Bay (Liengme). GERMAN EAST AFRICA: Darassalaam (H. Brauns); Bagamoyo (Stuhlmann; Le Roy). ZANZIBAR. BRITISH EAST AFRICA: Kikuyu Escarpment, 2250 m. (Alluaud and Jeannel). SOMALILAND: (C. Keller).

17. Var. *detrita* EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 118 (♀). FOREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 350 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 172.

Type locality: Obbia, SOMALILAND (Bricchetti-Robecchi).

SOMALILAND: (C. Keller).

17s. Var. *hirsuta* EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 118 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 172.

Type locality: Ellahelaj, SOMALILAND (Bricchetti-Robecchi).

17s. Var. *pilipes* EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 118 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 172. EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 180; 1897, *ibid.*, XXXVIII, p. 601.

Type locality: Milmil, SOMALILAND (Bricchetti-Robecchi).

ABYSSINIA: Ogaden (Ruspoli); Boran Galla, Middle Ganale; Webi (Bricchetti-Robecchi). SOMALILAND: Wells of Laffarugh to wells of Aberio (V. Bottego).

17a. Subsp. *fallax* (MAYR) EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 43. FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 249 (♀).

Plagiolepis fallax MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidae, p. 54 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 172. WASMANN, 1894, 'Verzeichn. Myrmecoph. Termitoph. Arthrop.,' p. 113; 1896, Notes Leyden Mus., XVIII, p. 75. E. SIMON, 1899, Bull. Soc. Ent. France, p. 179. WASMANN, 1900, Zool. Jahrb. Abt. Syst., XIV, p. 68. PÉRINGUEY, 1900, Ann. Soc. Ent. France, LXIX, p. 67. FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 54 (♀ ?); 1910, *ibid.*, LIV, p. 449 (♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 24 (♀, ♀).

Plagiolepis (Anoplolepis) custodiens var. *fallax* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 377 (♀).

Type locality: Cape of Good Hope (Novara Expedition; Raffray).

CAPE PROVINCE: Constantia (F. Silvestri); Stormsvlei; Swellendam (F. Purell); Cape Town (Wilms; E. Simon); Willowmore (H. Brauns). NATAL: (Haviland). BELGIAN CONGO: Umangi (Wilwerth; this record probably does not refer to the subsp. *fallax*).

18. *Plagiolepis (Anoplolepis) decolor* EMERY.

Plagiolepis decolor EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 44 (♀). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 23 (♀, ♀, ♂).

Type locality: Kimberley, CAPE PROVINCE (E. Simon).

TRANSVAAL: Hamman's Kraal (E. Simon). GERMAN SOUTHWEST AFRICA: Cape Cross (L. Schultze). BECHUANALAND: Khakhea to Kooa (L. Schultze).

19. *Plagiolepis (Anoplolepis) longipes* (JERDON) SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 123. See p. 1036.

Formica longipes JERDON, 1851, Madras Journ. Litt. Sc., XVII, p. 122 (♀); 1854, Ann. Mag. Nat. Hist., (2) XIII, p. 105 (♀).

Formica gracilipes F. SMITH, 1857, Journ. Proc. Linn. Soc. London, Zool., II, p. 55 (♀); 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 22 (♀).

Formica trifasciata F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 27 (♀).

Prenolepis gracilipes MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 698.

Plagiolepis gracilipes MAYR, 1867, Tijdschr. v. Ent., X, p. 73 (♀, ♀); 1870, Verh. Zool. Bot. Ges. Wien, XX, p. 947. ROTHNEY, 1889, Trans. Ent. Soc. London, p. 373. MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 197.

Plagiolepis longipes EMERY, 1887, Ann. Mus. Civ. Genova, XXIV, p. 247; 1888, Zeitschr. Wiss. Zool., XLVI, p. 385, Pl. XXVII, figs. 8-9. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 173. FOREL, 1894, Journ. Bombay Nat. Hist. Soc., VIII, p. 414 (♀). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 149. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 39. BINGHAM, 1903, 'Fauna Brit. India, Hym.,' II, p. 320, fig. 97 (♀).

Type locality: INDIA (Jerdon).

Tropicopolitan. ZANZIBAR: (H. Brauns; Stuhlmann).

20. *Plagiolepis (Anoplolepis) nuptialis* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 289 (♂).

Type locality: Willowmore, CAPE PROVINCE (G. Arnold).

CAPE PROVINCE: Keurbooms River near Plettenberg Bay (H. Brauns).

21. *Plagiolepis (Anoplolepis) opaciventris* EMERY.*Plagiolepis opaciventris* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 492 (♀).*Plagiolepis steingröveri* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 44 (♀; nec ♀, nec Forel).

Type locality: Gaboon, FRENCH CONGO.

22. *Plagiolepis (Anoplolepis) rufescens* SANTSCHI, 1917, Ann. Soc. Ent.

France, LXXXV, (1916), p. 289 (♀).

Type locality: Cape Town, CAPE PROVINCE (Reichensperger).

23. *Plagiolepis (Anoplolepis) steingröveri* FOREL. See p. 1036.*Plagiolepis steingröveri* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 72 (♀); 1910, Ann. Soc. Ent. Belgique, LIV, p. 449 (♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 23 (♀); 1911, Sitzb. Bayer. Akad. Wiss., p. 287 (♀).*Plagiolepis steingröveri* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 44 (♀; nec ♀). WASMANN, 1904, Notes Leyden Mus., XXV, pp. 73 and 75; 1911, Tijdschr. v. Ent., LIV, p. 202; 1912, Zeitschr. Wiss. Zool., CI, p. 106; 1920, Tijdschr. v. Ent., LXII, (1919), p. 120.*Plagiolepis (Anoplolepis) steingröveri* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 123.

Type locality: Angra Pequena, GERMAN SOUTHWEST AFRICA (Steingröver).

MOSSAMEDES: (Baum and Van der Kellen). GERMAN SOUTHWEST AFRICA: (Lübbert); Lüderitzbucht (L. Schultze). BECHUANALAND: KOOA; Severelela to Khakhea (L. Schultze). CAPE PROVINCE: Cape Town (E. Simon; Wilms); Willowmore (H. Brauns). ORANGE FREE STATE: Bothaville (H. Brauns); Bloemfontein (E. Simon). TRANSVAAL: Makapan; Hebron (E. Simon).

24. *Plagiolepis (Anoplolepis) tenella* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 123. See p. 214.*Plagiolepis tenella* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 364 (♀); 1911, Rev. Zool. Afr., I, p. 210 (♀). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 61 (♀; without description).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

BELGIAN CONGO: Lower Congo, in stomach of *Manis temminckii* (Solon); Niapu; Akenge; Medje; Bafwasende; Garamba (Lang and Chapin).25. *Plagiolepis (Anoplolepis) trimeni* FOREL.*Plagiolepis trimeni* FOREL, 1895, Ann. Soc. Ent. Belgique, XXXIX, p. 430 (replete ♀).*Plagiolepis trimeni* EMERY, 1899, ibid., XLIII, p. 492 (♀). GILCHRIST, 1910, Agric. Journ. Cape of Good Hope, XXXVI, p. 328. FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 223 (♀).

Type locality: NATAL (Mutschinson).

RHODESIA: Bulawayo; Bembesi (G. Arnold).

26. *Plagiolepis (Anoplolepis) tumidula* EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 19, fig. 10 (♀, ♂).

Type locality: Ghinda, ETRURIA (F. Silvestri).

Acantholepis MAYR*Acantholepis* MAYR, 1861, 'Europ. Formicid.,' p. 42.*Formica* (part) F. SMITH. *Hypoclinea* (part) MAYR, ROGER.Genotype: *Hypoclinea frauenfeldi* Mayr, 1855.

1. *Acantholepis arenaria* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 555 and 560, fig. 56 (♀).

Type locality: Sawmills, Umgusa River, SOUTHERN RHODESIA (G. Arnold).

2. *Acantholepis arnoldi* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 142 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 563 (♀).

Type locality: Bulawayo, RHODESIA (G. Arnold).

RHODESIA: Redbank (G. Arnold).

3. *Acantholepis capensis* MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, pp. 699 (♀) and 770 (♂); 1863, *ibid.*, XIII, p. 394. ROGER, 1863, 'Verzeich. Formicid.', p. 11. MAYR, 1865, 'Reise Novara, Zool.', II, Formicidae, p. 56 (♀, ♂), Pl. II, fig. 16. EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 366 (♀, ♂, ♂). MAGRETTI, 1884, *ibid.*, XXI, p. 537 (♀); 1884, Bull. Soc. Ent. Italiana, XV, (1883) p. 244 (♀). PÉRINGUEY, 1886, Trans. Ent. Soc. London, Proc., p. xxxvi. FOREL, 1892, Ann. Soc. Ent. Belgique, XXXVI, p. 42 (♀). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 171. WASMANN, 1894, 'Verzeichn. Myrmecoph. Termitoph. Arthrop.', pp. 108, 118, and 215. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 74. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 46. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 149. EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 180 (♀). WASMANN, 1896, Notes Leyden Mus., XVIII, p. 76; 1898, Wien. Ent. Zeitg., XVII, p. 97; 1899, Notes Leyden Mus., XXI, pp. 39 and 51. FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 311; in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 564; 1904, Mém. Ac. Sc. St. Pétersbourg, IX, (1903), p. 387 (♀). WASMANN, 1904, Notes Leyden Mus., XXV, pp. 37 and 73. DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, pp. 310 and 379. FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 54 (♀, ♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 25 (♀, ♀, ♂); 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 265 (♀, ♀, ♂). WASMANN, 1911, Tijdschr. v. Ent., LIV, p. 202; 1912, Zeitschr. Wiss. Zool., CI, pp. 104 and 106. REICHENSPERGER, 1913, Zool. Jahrb. Abt. Syst., XXXV, p. 203. FOREL, 1913, Rev. Zool. Afr., II, p. 338 (♀). BEQUAERT, *ibid.*, p. 429. FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 249 (♀, ♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 377 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 126 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 557 and 568 (♀, ♂). See p. 1036.

Type locality: Table Mountain at the Cape of Good Hope, CAPE PROVINCE (Novara Expedition).

FRENCH GUINEA: Los Islands (H. Brauns). SIERRA LEONE: (Mocqueries). BELGIAN CONGO: Elisabethville; Sankisia (J. Bequaert). MOSSAMEDES: between the Cubango and Cuito Rivers (Baum and Van der Kellen). GERMAN SOUTHWEST AFRICA: (Peters); Kubub (L. Schultze); Gobabis (Boschmann); Cape Cross to Swakopmund (Dinter). BECHUANALAND: Kooa; Khakhea to Kang (L. Schultze). CAPE PROVINCE: Port Elizabeth; Willowmore (H. Brauns); Sir Lowry Pass (F. Purcell); Cape Town; Matjesfontein (E. Simon). TRANSVAAL: Makapan (E. Simon); Pretoria (E. Simon; F. Silvestri). NATAL: Estcourt (R. C. Wroughton). DELAGOA: (Liengme). ERITREA: Asmara; Nefasit; Ghinda (K. Escherich); Keren; Sciotel (Beccari). ABYSSINIA: (Ilg); Boran Galla; Auata (V. Bottego); Adis Abeba (Kachovsky). ANGLO-EGYPTIAN SUDAN: Sebderat near Kassala (Magretti).

3. Var. *anceps* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 438 (♀). See p. 215.

Type locality: BELGIAN CONGO (Kohl).

BELGIAN CONGO: Stanleyville; Medje (Lang and Chapin).

3₂. Var. **guineensis** MAYR, 1902, Verh. Zool. Bot. Ges. Wien, LII, p. 296 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 125 (♀). See p. 215.

Type locality: Akra, GOLD COAST (R. Buchholz).

BELGIAN CONGO: Thysville (Lang and Chapin).

3₃. Var. **simplicoides** (FOREL) SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 125 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 572 (♀).

Acantholepis capensis subsp. *simplicoides* FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 86, footnote (♀).

Type locality: BASUTOLAND (R. C. Wroughton).

3₄. Var. **validiuscula** EMERY, 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 602 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 143 (♀); 1913, Rev. Zool. Afr., II, p. 338 (♀); 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 249 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 126 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 557 and 570 (♀, ♂). See p. 216.

Type locality: Webi, ABYSSINIA (Ruspoli).

BELGIAN CONGO: Sankisia (J. Bequaert); Thysville (J. Bequaert; Lang and Chapin). RHODESIA: very common (G. Arnold). CAPE PROVINCE: very common (G. Arnold); Kimberley (Power); King William's Town (R. Godfrey).

3₅. Subsp. **canescens** (EMERY) REICHENSPERGER, 1913, Zool. Jahrb. Abt. Syst., XXXV, pp. 187, 189, 190, 197, 198, and 215. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' pp. 124 and 126 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 378 (♀). EMERY, 1915, *ibid.*, X, p. 4 (♀). REICHENSPERGER, 1915, Ent. Mitt. Deutsch. Ent. Mus. Berlin, IV, p. 122. WASMANN, 1918, Tijdschr. v. Ent., LX, (1917), p. 397. See p. 216.

Acantholepis capensis var. *canescens* EMERY, 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 601 (♀); 1899, *ibid.*, XXXIX, p. 500. MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 8; 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. 1, p. 389. SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 391 (♀). FOREL, 1911, Rev. Zool. Afr., I, p. 280 (♀). GESTRO, 1911, Bull. Soc. Ent. Italiana, XLI, (1909), pp. 257 and 260. STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 390 (♀). KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 9 (♀).

Type locality: Coromina, ABYSSINIA (Ruspoli).

SENEGAMBIA: Dakar (F. Silvestri). FRENCH GUINEA: Kindia (F. Silvestri). GOLD COAST: Aburi (F. Silvestri). SOUTHERN NIGERIA: Lagos (F. Silvestri). FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: Congo da Lemba (R. Mayné); Thysville (J. Bequaert); Avakubi (Lang and Chapin). GERMAN EAST AFRICA: Lake Mohasi (Schubotz). BRITISH EAST AFRICA: Ramisi; Blue Post Hotel, Kikuyu, 1520 m. (Alluaud and Jeannel). UGANDA: Unyoro Province, near Hoima; region of Lake Albert (C. Alluaud); Gondokoro (F. Werner). ANGLO-EGYPTIAN SUDAN: Kaka, White Nile (I. Trägårdh); Khartum (Karawaiew); near Renk; between Khartum and Fashoda. ABYSSINIA: Buditu to Dimé (V. Bottego); Diré Dava; Abuker; Bisa Timo; Gebel Hakim; Harar (Kristensen); Lake Abaja (Ruspoli). ERITREA: Ghinda; Nefasit (F. Silvestri); Sciotel, Bogos (Beccari). Saganeiti; Adi Ugri; Adi Caié (Andreini).

3a₁. Var. *cacozela* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 126 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 378 (♀). See p. 216.

Acantholepis capensis var. *cacozela* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 395 (♀).

Type locality: FRENCH CONGO.

SOUTHERN NIGERIA: Olokemeji (F. Silvestri). BELGIAN CONGO: Faradje (Lang and Chapin). FRENCH CONGO: Fort Crampel (Schubotz).

3b. Subsp. *depilis* EMERY, 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 602 (♀, ♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 126 (♀).

Acantholepis carbonaria EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 119 (♀, part and ♀).

Type locality: Arigalga, SOMALILAND (Ruspoli).

ABYSSINIA: Leboi (Ruspoli).

3c. Subsp. *hirsuta* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' pp. 124 and 126, fig. 26 (♀).

Type locality: Molo, Mau Escarpment, 2420 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

3c₁. Var. *elevata* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 249 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 573 (♀).

Type locality: Hillside, Bulawayo, RHODESIA (G. Arnold).

3d. Subsp. *incisa* FOREL, 1913, Rev. Zool. Afr., II, p. 338 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 125, fig. 27 (♀).

Type locality: Elisabethville, BELGIAN CONGO (J. Bequaert).

BRITISH EAST AFRICA: Kijabe, Kikuyu Escarpment, 2100 m. (Alluaud and Jeannel).

3e. Subsp. *junodi* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 438 (♀). ARNOLD, 1920, Ann. South African Mus., XXIV, p. 572 (♀).

Type locality: Shilouvane, TRANSVAAL (Junod).

3f. Subsp. *laevis* (SANTSCHI).

Acantholepis simplex subsp. *laevis* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 378 (♀).

Acantholepis laevis SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 312, fig. 5 (♀).

Type locality: St. Louis, SENEGAMBIA (Claveau).

FRENCH GUINEA: Konakry (F. Silvestri).

3f₁. Var. *alexis* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 557 and 573 (♀).

Type Locality: Hillside, Bulawayo, RHODESIA (G. Arnold).

3g. Subsp. *simplex* FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 86, footnote. ARNOLD, 1920, Ann. South African Mus., XIV, pp. 557 and 572 (♀, ♂).

Acantholepis simplex FOREL, 1892, Ann. Soc. Ent. Belgique, XXXVI, p. 42 (♀); 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 350 (♀, ♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 172. WASMANN, 1894, 'Verzeichn. Myrmecoph. Termitoph. Arthrop.,' p. 107; 1912, Zeitschr. Wiss. Zool., CI, p. 103. REICHENSFELGER, 1913, Zool. Jahrb. Abt. Syst., XXXV, p. 197. FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 143 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 123 (♀).

Type locality: SOMALILAND (C. Keller).

RHODESIA: Bulawayo (G. Arnold). BRITISH EAST AFRICA: Ramisi (Alluaud and Jeannel). UGANDA: Kadjura Swamp, near Hoima, region of Lake Albert (C. Alluaud). CAPE PROVINCE: Grahamstown (Baines and Cherry).

3g. Var. *minuta* (FOREL).

Acantholepis simplex var. *minuta* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 438 (♀).

Acantholepis capensis var. *minuta* ARNOLD, 1920, Ann. South African Mus., XIV, p. 572 (♀).

Type locality: Shilouvane, TRANSVAAL (Junod).

4. *Acantholepis capitata* FOREL, 1913, Rev. Zool. Afr., II, p. 338 (♀).

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

5. *Acantholepis carbonaria* EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 119 (♀, part; nec ♀). FOREL, 1892, Ann. Soc. Ent. Belgique, XXXVI, p. 42 (♀). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 171. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 74. STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08)', III, p. 390 (♀). See p. 216.

Type locality: Obbia, SOMALILAND (Bricchetti-Robecchi).

SOUTHERN ABYSSINIA: (Ilg). BELGIAN CONGO: Kwidjwi Island, Lake Kivu; Mawambi to Avakubi (Schubotz); Banana (Lang and Chapin).

5i. Var. *erythræa* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 265 (♀, ♀). Wasmann, 1912, Zeitschr. Wiss. Zool., CI, p. 104. EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 4 (♀).

Type locality: Nefasit, ERITREA (K. Escherich; F. Silvestri).

ERITREA: Ghinda (K. Escherich). ABYSSINIA: Harar (Ilg).

5s. Var. *obtusa* EMERY, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 63 (♀).

Type locality: Saati, ERITREA (Belli).

ERITREA: Sabarguma (Belli).

5a. Subsp. *baumi* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 449 (♀).

Type locality: MOSSAMEDES (Baum and Van der Kellen).

6. *Acantholepis crinita* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 149 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 565 (♀).

Type locality: Port Natal, NATAL (H. Brauns).

7. *Acantholepis curta* EMERY, 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 603, fig. (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 354 (♀).

Acantholepis capensis EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 527 (♀). (nec Mayr).

Type locality: Assab, ERITREA (Doria).

ERITREA: Ras Doumeira (Doria). BELGIAN CONGO: Yambata (R. Mayné).

8. *Acantholepis deplanata* STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08)', III, p. 390, fig. 8 (♀).

Type locality: Buddu Forest, northwest of Bukoba, GERMAN EAST AFRICA (Schubotz).

9. *Acantholepis depressa* SANTSCHE, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicide,' p. 126, fig. 28 (♀).

Type locality: Ramisi River, BRITISH EAST AFRICA (Alluaud and Jeannel).

10. *Acantholepis egregia* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 143 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 557 and 574 (♀).

Type locality: Redbank, SOUTHERN RHODESIA (G. Arnold).

- 10a. Subsp. *santschii* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 575 (♀).
Type locality: Cawston Farm, Ungusa River, SOUTHERN RHODESIA (G. Arnold).
11. *Acantholepis foreli* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 564 (♀), Pl. VIII, fig. 121.
Type locality: Sipapoina, SOUTHERN RHODESIA (G. Arnold).
- 11i. Var. *convexa* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 565 (♀).
Type locality: Hillside, Bulawayo, SOUTHERN RHODESIA (G. Arnold).
- 11s. Var. *impressa* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 565 (♀).
Type locality: Hillside, Bulawayo, SOUTHERN RHODESIA (G. Arnold).
12. *Acantholepis gerardi* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 262 (♀).
Type locality: Kalanga, Upper Lukuga, BELGIAN CONGO (Gérard).
BELGIAN CONGO: Kataki (Gérard).
13. *Acantholepis gracilicornis* FOREL, 1892, Ann. Soc. Ent. Belgique, XXXVI, p. 42 (♀). EMERY, 1895, Ann. Soc. Ent. Belgique, XXXIX, p. 337; 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 63. FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 265 (♀). KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 9 (♀).
Type locality: Aden, SOUTHERN ARABIA (F. Ris).
ERITREA: Asmara (K. Escherich); Sabarguma (Belli). ANGLO-EGYPTIAN SUDAN: Port Sudan; Khartum; Assuan (Karawaiew).
- 13a. Subsp. *abdominalis* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 73 (♀).
Type locality: SOUTHERN ABYSSINIA (Ilg).
14. *Acantholepis longinoda* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 555 and 559 (♀, ♀), Pl. VIII, figs. 120 and 120a.
Type locality: Hillside, Bulawayo, SOUTHERN RHODESIA (G. Arnold).
15. *Acantholepis nigriventris* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 493 (♀).
Type locality: CAMEROON (Conradt).
16. *Acantholepis rubrovaria* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 142. ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 562 (♀, ♂).
Acantholepis simplex var. *rubrovaria* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 449 (♀).
Type locality: BASUTOLAND (R. C. Wroughton).
- 16a. Subsp. *pilosa* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 141 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 562 (♀).
Type locality: Redbank, SOUTHERN RHODESIA (G. Arnold).
- 16a. Var. *avunculus* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 563 (♀).
Type locality: Sawmills, Ungusa River, SOUTHERN RHODESIA (G. Arnold).
17. *Acantholepis silvicola* ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 576, fig. 57 (♀, ♀, ♂).
Type locality: Amatongas Forest, PORTUGUESE EAST AFRICA (G. Arnold).
18. *Acantholepis spinosior* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 144 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 557 (♀, ♀).
Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

18i. Var. **ballaensis** ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 558 (♂).

Type locality: Balla-Balla, SOUTHERN RHODESIA (G. Arnold).

18s. Var. **natalensis** ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 559 (♂).

Type locality: Pietermaritzburg, NATAL (C. Akerman).

19. **Acantholepis submetallica** ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 566 (♂, ♀, ♂).

Type locality: Bulawayo, SOUTHERN RHODESIA (G. Arnold).

19i. Var. **aspera** ARNOLD, 1920, Ann. South African Mus., XIV, pp. 556 and 568 (♂).

Type locality: Amatongas Forest, PORTUGUESE EAST AFRICA (G. Arnold).

20. **Acantholepis vestita** (F. SMITH) DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, pp. 333, 336, and 365.

Formica vestita F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 32 (♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 214.

Type locality: Port Natal, NATAL.

NATAL: Colenso; Newcastle (Dixey and Longstaff). CAPE PROVINCE: Stormberg Junction; Shanks Station (Dixey and Longstaff).

Myrmelachistini Forel

Aphomomyrmex EMERY

Aphomomyrmex EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 493.

Genotype: *Aphomomyrmex afer* Emery, 1899.

1. **Aphomomyrmex afer** EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 494, fig. (♂, ♀). FOREL, 1910, *ibid.*, LIV, p. 449 (♂); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 25 (♀). ARNOLD, 1920, Ann. South African Mus., XIV, p. 553 (♂).

Type locality: CAMEROON (Conradt; v. Muralt).

West Africa: (Fülleborn).

2. **Aphomomyrmex muralti** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 449 (♂). ARNOLD, 1920, Ann. South African Mus., XIV, p. 553 (♂).

Type locality: NATAL (L. v. Muralt).

Prenolepidini Forel

Prenolepis MAYR

Prenolepis MAYR, 1861, 'Europ. Formicid.,' p. 52.

Paratrechina MOTSCHOUISKY, 1863, Bull. Soc. Natural. Moscou, XXXVI, 3, p. 13.

Formica (part) LATREILLE *et auctor*. *Lasius* MAYR. *Tapinoma* F. SMITH, MAYR, etc.

Genotype: *Formica imparis* Say, 1836.

Subgenus 1. *Prenolepis* MAYR, *sensu stricto*

Subgenotype: same as genotype.

1. **Prenolepis kohli** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 438 (♂, ♀?).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

Subgenus 2. *Nylanderia* EMERY

Prenolepis subg. *Nylanderia* EMERY, 1906, Ann. Soc. Ent. Belgique, L, p. 134.

Subgenotype: *Formica vividula* Nylander, 1846.

2. *Prenolepis (Nylanderia) albipes* EMERY.

Prenolepis albipes EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 497 (♀).

Type locality: CAMEROON (Conrads).

3. *Prenolepis (Nylanderia) bourbonica* FOREL. See p. 1037.

Prenolepis nodifera subsp. *bourbonica* FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 210 (♀, ♂).

Prenolepis bourbonica FOREL, in Grandidier, 1891, 'Hist. Phys. Nat. Madagascar,' XX, 2, p. 82 (♀, ♀, ♂), Pl. III, fig. 2; in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 87 (♀).

Type locality: RÉUNION.

BRITISH EAST AFRICA: Chake-Chake, Pemba Island (Voeltzkow).

4. *Prenolepis (Nylanderia) grisoni* FOREL, 1914, Rev. Suisse Zool., XXIV, p. 440, fig. 5 (♀).

Type locality: Bengamisa, BELGIAN CONGO (Kohl).

5. *Prenolepis (Nylanderia) incallida* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 263, fig. 12 (♂).

Type locality: SAN THOMÉ Island.

6. *Prenolepis (Nylanderia) jägerskiöldi* MAYR.

Prenolepis jägerskiöldi MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 8 (♂). EMERY, 1906, Ann. Soc. Ent. Belgique, L, p. 130. KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 9, fig. 3 (♀, ♀, ♂).

Prenolepis (Nylanderia) jägerskiöldi EMERY, 1910, Deutsch. Ent. Zeitschr., pp. 127 and 130, figs. 4 and 5 (♀, ♀, ♂).

Prenolepis vividula ERN. ANDRÉ, 1881, Ann. Soc. Ent. France, (6) I, p. 61 (♀, ♀, ♂); 1882, 'Spec. Hym. Europ. Algérie,' II, pp. 204 and 206 (♀, ♀, ♂), Pl. x, figs. 8-10 and 12 (*ex typ.*). FOREL, in Grandidier, 1891, 'Hist. Phys. Nat. Madagascar,' XX, 2, p. 88 (♂), Pl. II, fig. 1 (*ex typ.*) (*nec vividula* Nylander).

Prenolepis jägerskiöldi var. *borcardi* SANTSCHI, 1908, Ann. Soc. Ent. France, LXXVII, p. 533, fig. 12 (♀, ♀, ♂).

Prenolepis (Nylanderia) jägerskiöldi SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 128 (♂).

Type locality: CAIRO, EGYPT (Jägerskiöld; Karawaiew; Borcard).

EGYPT, CYPRUS, SYRIA: GERMAN EAST AFRICA: Tanga (Alluaud and Jeannel).

7. *Prenolepis (Nylanderia) lepida* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 262, fig. 11 (♂).

Type locality: VICTORIA, CAMEROON (Reichensperger).

8. *Prenolepis (Nylanderia) longicornis* (LATREILLE) SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 127 (♂). LAMBORN, 1920, Trans. Ent. Soc. London, (1919), Proc., p. lvi. See pp. 217 and 1038.

Formica longicornis LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 113 (♂). F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 31. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 417.

Formica vagans JERDON, 1851, Madras Journ. Litt. Sc., XVII, p. 124 (♀, ♀); 1854, Ann. Mag. Nat. Hist., (2) XIII, p. 107 (♀, ♀).

Formica (Tapinoma) gracilescens NYLANDER, 1856, *Ann. Sc. Nat. Zool.*, (4) V, p. 73 (♀), Pl. III, fig. 20.

Formica gracilescens NYLANDER, 1856, *Ann. Soc. Ent. France*, (3) IV, Bull., p. xxviii (♀).

Tapinoma gracilescens F. SMITH, 1858, 'Cat. Hym. Brit. Mus.', VI, p. 56. WHITE, 1883, 'Ants and Their Ways,' p. 257 (♀). FOWLER, 1885, *Ent. Monthly Mag.*, XXII, p. 276.

Prenolepis gracilescens MAYR, 1862, *Verh. Zool. Bot. Ges. Wien*, XII, p. 698.

Paratrechina currens MOTSCHOUJSKY, 1863, *Bull. Soc. Nat. Moscou*, XXXVI, 3, p. 14.

Prenolepis longicornis ROGER, 1863, 'Verzeich. Formicid.,' p. 10. MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidæ, p. 50 (♀); 1867, *Tijdschr. v. Ent.*, X, p. 72 (♀, ♀). FRAUENFELD, 1867, *Verh. Zool. Bot. Ges. Wien*, XVII, p. 442. MAYR, 1870, *ibid.*, XX, p. 947 (♀); 1876, *Journ. Mus. Godeffroy*, XII, p. 77. ERN. ANDRÉ, 1881, *Ann. Soc. Ent. France*, (6) I, p. 60 (♀, ♂). FOREL, 1881, *Mitth. München. Ent. Ver.*, V, p. 2. ERN. ANDRÉ, 1882, 'Spec. Hym. Europ. Algérie,' II, pp. 203, 205, and 206 (♀, ♀, ♂), Pl. X, fig. 11. EMERY, 1888, *Zeitschr. Wiss. Zool.*, XLVI, p. 379, Pl. XXVII, fig. 5. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 179. FOREL, 1909, *Ann. Soc. Ent. Belgique*, LIII, p. 54 (♀); 1910, *Zool. Jahrb. Abt. Syst.*, XXIX, p. 266 (♀). KARAWAIEW, 1911, *Rev. Russe Ent.*, XI, p. 9 (♀). LONGSTAFF, 1911, *Ent. Monthly Mag.*, XLVII, p. 124. FOREL, 1913, *Rev. Zool. Afr.*, II, p. 339 (♀); 1913, *Ann. Soc. Ent. Belgique*, LVII, p. 354 (♀, ♀); 1914, *Bull. Soc. Vaudoise Sc. Nat.*, (5) L, p. 249 (♀). SANTSCHI, 1914, *Boll. Lab. Zool. Gen. Agrar. Portici*, VIII, p. 378 (♀). STRIZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 395 (♀). SANTSCHI, 1920, 'Études Maladies Parasites Cacaoyer S. Thomé,' X, p. 3 (♀).

Prenolepis longicornis var. *hagemanni* FOREL, 1901, *Mitt. Naturh. Mus. Hamburg*, XVIII, p. 65 (♀); 1907, *ibid.*, XXIV, p. 15; 1916, *Rev. Suisse Zool.*, XXIV, p. 441 (♀).

Type locality: Senegal.

Tropicopolitan. WEST AFRICA: (Jourdan). FRENCH GUINEA: Konakry; Kakoulima (F. Silvestri). SOUTHERN NIGERIA: Lagos (F. Silvestri). SAN THOMÉ: (de Seabra). FRENCH CONGO: Fort de Possel to Fort Crampel (Schubotz). BELGIAN CONGO: Boma (C. Hagemann; Styczinski); Congo da Lemba (R. Mayné); Leopoldville; Kwamouth (J. Maes); St. Gabriel (Kohl); Zambi; Stanleyville (Lang and Chapin); Libenge (Schubotz). ANGLO-EGYPTIAN SUDAN: Khartum (Karawaiew). GERMAN EAST AFRICA: Tanga (Alluaud and Jeannel); Lindi (Lamborn); Nguela, Usambara. NATAL: Durban (G. Arnold; C. B. Cooper).

9. *Prenolepis (Nylanderia) trægærdhi* FOREL.

Prenolepis trægærdhi FOREL, in Wasmann, 1904, 'Termitophilen aus dem Sudan, Res. Swed. Zool. Exped. Jägerskiöld,' XIII, p. 14 (♀, ♂).

Type locality: Kaka, White Nile, ANGLO-EGYPTIAN SUDAN (I. Trägårdh).

9. Var. *natalensis* FOREL, 1915, *Bull. Soc. Vaudoise Sc. Nat.*, (5) L, p. 348 (♀).

Type locality: Durban, NATAL (G. Arnold).

10. *Prenolepis (Nylanderia) vividula* (NYLANDER) EMERY, 1906, *Ann. Soc. Ent. Belgique*, L, p. 134, figs. 1-4 (♀, ♀, ♂). See p. 218.

Formica vividula NYLANDER, 1847, *Acta Soc. Sc. Fennicæ*, II, 2, p. 900 (♀, ♀, ♂), Pl. XVIII, figs. 2 and 10-14.

Prenolepis vividula MAYR, 1861, 'Europ. Formicid.,' p. 52; 1870, Verh. Zool. Bot. Ges. Wien, XX, pp. 948 and 949 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 180.

Type locality: in the hothouses of the Botanical Garden at Helsingfors, FINLAND (Nylander).

Tropicopolitan. BELGIAN CONGO: Niapu (Lang and Chapin).

11. *Prenolepis (Nylanderia) waelbroeckii* EMERY.

Prenolepis waelbroeckii EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 496, fig. (♀, ♂).

Prenolepis waelbroeckii FOREL, 1909, ibid., LIII, p. 54 (♂); 1911, Rev. Zool. Afr., I, p. 280 (♀?).

Type locality: Kinshasa, BELGIAN CONGO (Waelbroeck).

BELGIAN CONGO: Congo da Lemba (R. Mayné).

12. *Prenolepis (Nylanderia) weissii* SANTSCHI, 1911, Rev. Zool. Afr., I, p. 210 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

13. *Prenolepis (Nylanderia) zelotypa* SANTSCHI.

Prenolepis (Nylanderia) jagerskioldi var. *zelotypa* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 264 (♀).

Prenolepis zelotypa SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 238.

Type locality: BRITISH EAST AFRICA.

RHODESIA: Victoria Falls (G. Arnold).

Formicini Forel¹

Pseudolasius EMERY

Pseudolasius EMERY, 1887, Ann. Mus. Civ. Genova, XXIV, p. 244.

Formica (part) F. SMITH. *Lasius* (part) MAYR.

Genotype: *Formica familiaris* F. Smith, 1860.

1. *Pseudolasius bucculentus* Wm. M. WHEELER. See p. 222 (♀, ♂).

Type locality: Medje, BELGIAN CONGO (Lang and Chapin).

2. *Pseudolasius bufonum* Wm. M. WHEELER. See p. 220 (♀, ♀, ♂).

Type locality: Medje, BELGIAN CONGO (Lang and Chapin).

3. *Pseudolasius gowdeyi* Wm. M. WHEELER. See p. 223 (♀).

Acropyga gowdeyi GOWDEY, 1917, Bull. Ent. Research, VIII, pt. 2, p. 187 (without description).

Type locality: Entebbe, UGANDA (C. C. Gowdey).

4. *Pseudolasius weissii* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 391, fig. 16 (♀, ♀). EMERY, 1911, Ann. Soc. Ent. Belgique, LV, pp. 214 and 219 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

4. Var. *sordidus* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 378 (♀). See p. 219.

Type locality: Aburi, GOLD COAST (F. Silvestri).

BELGIAN CONGO: Akenge (Lang and Chapin).

¹Bingham, 1903, Ann. Mag. Nat. Hist., (7) XII, p. 62, records *Formica rufibarbis* Fabricius (♀, ♀) as taken by Distant at Pretoria, Transvaal. It would be interesting to know whether this Palearctic ant has really been introduced into South Africa.

Cataglyphis FÖRSTER

Cataglyphis FÖRSTER, 1850, Verh. Naturh. Ver. Preuss. Rheinl., VII, p. 493.

Monocombus MAYR, 1855, Verh. Zool. Bot. Ges. Wien, V, p. 381.

Formica (part) FABRICIUS, etc. *Tapinoma* F. SMITH. *Myrmecocystus* EMERY, FOREL, MACCOCK, ERN. ANDRÉ, etc.

Genotype: *Formica megalocola* Förster, ♀, 1850 = *Cataglyphis fairmairei* Förster, ♂, 1850.

1. ***Cataglyphis cursor*** (BOYER DE FONSCOLOMBE) MAYR, 1861, 'Europ. Formicid.,' p. 45 (♀, ♀, ♂).

Formica cursor BOYER DE FONSCOLOMBE, 1846, Ann. Soc. Ent. France, (2) IV, p. 41 (♀, ♀, ♂); *ibid.*, Bull., p. lxxix (♀).

Myrmecocystus cursor EMERY, 1898, Öfvers. Finsk. Vet.-Soc. Förh., XI, p. 147, footnote (♀).

Type locality: Aix, FRANCE (Boyer de Fonscolombe).

SOUTHERN EUROPE, ANATOLIA, and CENTRAL ASIA. Emery (*loc. cit.*) records it from the GOLD COAST.

2. ***Cataglyphis viaticus*** (FABRICIUS) MAYR, 1861, 'Europ. Formicid.,' p. 45 (♀, ♀, ♂). RADOSZKOWSKY, 1876, Hor. Soc. Ent. Rossicae, XII, p. 140. EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 366 (♀, ♀).

Formica viatica FABRICIUS, 1787, 'Mantissa Insect.,' I, p. 308 (♀).

Myrmecocystus viaticus EMERY, 1880, Ann. Mus. Civ. Genova, XV, p. 389 (♀). MAGRETTI, 1884, *ibid.*, XXI, p. 537 (♀). FOREL, 1894, Journ. Bombay Nat. Hist. Soc., VIII, p. 402.

Type locality: SPAIN (Vahl).

EASTERN EUROPE, NORTH AFRICA, WESTERN and CENTRAL ASIA, NORTHERN INDIA. EGYPT. ABYSSINIA: Harar (Ilg). ANGLO-EGYPTIAN SUDAN: Sebderat; Kassala (Magretti). ERITREA: Keren; Sciotel; Anseba (Beccari); Sogodas (Magretti).

2a. Subsp. ***abyssinicus*** (FOREL).

Myrmecocystus viaticus subsp. *abyssinicus* FOREL, 1904, Ann. Mus. Zool. Ac. Sc. St. Pétersbourg, VIII, (1903), p. 382 (♀, ♀).

Type locality: Ingfal, ABYSSINIA (Kachovskij).

ABYSSINIA: Schoa; western Abyssinia (Ilg).

2b. Subsp. ***adenensis*** (FOREL).

Myrmecocystus viaticus subsp. *adenensis* FOREL, 1904, Ann. Mus. Zool. Ac. Sc. St. Pétersbourg, VIII, (1903), p. 382, footnote (♀).

Type locality: Aden, southern ARABIA (F. Ris).

2c. Subsp. ***bicolor*** (FABRICIUS).

Formica bicolor FABRICIUS, 1793, 'Ent. Syst.,' II, p. 351 (♂). LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 173 (♂). FABRICIUS, 1804, 'Syst. Piez.,' p. 398 (♂).

Myrmecocystus viaticus var. *desertorum* FOREL, 1894, Journ. Bombay Nat. Hist. Soc., VIII, p. 402 (♀); 1894, Mitth. Schweiz. Ent. Ges., IX, p. 72. EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 180.

Myrmecocystus viaticus subsp. *desertorum* K. ESCHERICH, 1902, Allgem. Zeitschr. Ent., VII, p. 354. MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 10.

Myrmecocystus bicolor MAYR, 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. I, p. 389.

Myrmecocystus viaticus subsp. *bicolor* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 450.

Cataglyphis bicolor EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 22 (♀, ♂).

Type locality: BARBARY (Desfontaines).

ALGERIAN SAHARA. EGYPT. ANGLO-EGYPTIAN SUDAN: Edeloud in the Kordofan Desert (F. Werner); Gebelein (I. Trägårdh). ERITREA: Mayabal; Nefasit (F. Silvestri). ABYSSINIA: southern Abyssinia (Hg); Ogaden (V. Bottego). SOMALILAND: wells of Laffarugh to wells of Aberio (V. Bottego). GOLD COAST (doubtful record).

2c. Var. *congolensis* (Stitz).

Myrmecocystus bicolor var. *congolensis* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 396, fig. 10 (♀).

Type locality: Fort Archambault, FRENCH CONGO (Schubotz).

2d. Subsp. *setipes* (FOREL).

Myrmecocystus viaticus subsp. *setipes* FOREL, 1894, Journ. Bombay Nat. Hist. Soc., VIII, p. 401 (♀).

Type locality: Nusseerabad, INDIA (Gladston).

2d. Var. *seticornis* (EMERY).

Myrmecocystus viaticus subsp. *setipes* var. *seticornis* EMERY, 1898, Öfvers. Finsk. Vet. Soc. Förh., XL, p. 147, footnote (♀).

Type locality: GOLD COAST.

Ecophyllini Forel

Ecophylla F. SMITH

Ecophylla F. SMITH, 1861, Journ. Proc. Linn. Soc. London, Zool., V, p. 101.

Formica (part) FABRICIUS, et auct. *Myrmica* (part) HEER.

Genotype: *Formica smaragdina* Fabricius, 1775.

1. *Ecophylla longinoda* (LATREILLE) GERSTÄCKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 354; 1873, in 'v. d. Decken's Reisen in Ost Afrika, Gliederthiere,' p. 343 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 176. H. KOHL, 1906, Natur u. Offenbarung, LII, pp. 166-167; 1909, *ibid.*, LV, p. 169. WASMANN, 1911, 1^{er} Congr. Intern. Entom. Bruxelles (1910), II, Mém., p. 231; 1913, Ann. Rept. Smiths. Inst. for 1912, p. 472. FOREL, 1913, Rev. Zool. Afr., II, p. 339 (♀). See p. 227.

Formica longinoda LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 184 (♀), Pl. XI, fig. 72.

Echophylla nivescens RADOSZKOWSKY, 1881, Journ. Sci. Ac. Lisboa, VIII, No. 31, p. 197.

Ecophylla brevinodis ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 313 (♀). STITZ, 1913, Mitt. Zool. Mus. Berlin, V, p. 148 (♀). SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 345 (♀, ♀).

Ecophylla smaragdina subsp. *longinoda* EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 564 (♀, ♀). FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 72. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 2 (♀, ♀). MAYR, 1896, Ent. Tidskr., XVII, p. 249. FOREL, 1897, Abhandl. Senckenberg. Naturf. Ges., XXI, p. 188 (♀). H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 39. EMERY, 1899, Ann. Mus. Civ. Genova, XXXIX, p. 500 (♀, ♀). MAYR, in Sjöstedt, 1907, 'Exped. Kili-

mandjaro, Meru, etc., II, 8, p. 18. FOREL, 1907, Mitt. Naturh. Mus. Hamburg, XXIV, p. 15 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXXVIII, (1909), p. 391. FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, pp. 54 (♀, ♀), 61 (♀), and 73 (♀, ♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 25 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 148 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 354 (♀, ♂). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 378 (♀). LAMBORN, 1914, Trans. Ent. Soc. London, (1913), p. 444. STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 396 (♀).

Ecophylla longinoda var. *brevinoda* DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 176.

Ecophylla smaragdina var. *longinoda* L. REH, 1905, Zeitschr. f. Pflanzenkrankh., XV, p. 135. AULMANN and LA BAUME, 1912, 'Fauna d. Deutsch. Kolon.,' V, 3, p. 65, fig. 37. AULMANN, 1913, *ibid.*, V, 5, p. 95, fig. 80. FOREL, 1913, Rev. Suisse Zool., XXI, p. 670 (♀, ♂).

Ecophylla smaragdina subsp. *longinoda* var. *brevinoda* SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 345 (♀, ♀).

Ecophylla smaragdina subsp. *brevinoda* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 396 (♀).

Type locality: Senegal.

SÉNÉGAMBIA: Dakar (F. Silvestri). GAMBIA: Bathurst (C. Alluaud). FRENCH GUINEA: Kindia (F. Silvestri). SIERRA LEONE: (Mocquerys). LIBERIA. IVORY COAST: Grand Bassam (Lohier; Bonet); Assinie (C. Alluaud). GOLD COAST. SLAVE COAST. SOUTHERN NIGERIA: Oni Camp, east of Lagos (Lamborn). CAMEROON: (Sjöstedt; Conradt); Bibundi (Tessmann); Molundu (Reichensperger); Victoria (Winkler). SPANISH GUINEA: Alen (Tessmann). FRENCH CONGO: Ogowé (Mocquerys); Brazzaville; Madingu (A. Weiss); Fort Crampel (Schubotz); Boda. BELGIAN CONGO: Duma (Schubotz); Kondué; Sankuru (Lujá); Mondombe (E. Mayné); Leopoldville to Stanleyville (Weyns); Mayombe (de Briey); Banana (Buschodts); Kinshasa (Waelbroeck); Malela; Faradje (Lang and Chapin); Katala (J. Bequaert); Boma. ANGOLA: (Welwitsch); San Antonio (Lang and Chapin). GERMAN EAST AFRICA: Tanga (Sjöstedt). BRITISH EAST AFRICA: Mombasa (v. d. Decken). ZANZIBAR: (Stuhlmann; Voeltzkow). PORTUGUESE EAST AFRICA: Delagoa (Liengme). ABYSSINIA: Buditu to Dimé; Dimé to Bass Narok (V. Bottego).

1. Var. *annectens* Wm. M. WHEELER. See p. 230 (♀, ♀, ♂).

Type locality: Avakubi, BELGIAN CONGO (Lang and Chapin).

BELGIAN CONGO: Stanleyville; Niangara (Lang and Chapin); Malela (J. Bequaert).

1. Var. *fusca* (EMERY). See p. 231.

Ecophylla fusca EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 496 (♀, ♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 148 (♀); 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 396 (♀).

Ecophylla smaragdina subsp. *fusca* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 354.

Type locality: CAMEROON (Conradt).

LIBERIA: Monrovia (J. Morris). NIGERIA. SPANISH GUINEA: Alen (Tessmann). BELGIAN CONGO: Stanleyville; Garamba (Lang and Chapin); Duma (Schubotz).

1. Var. *rubriceps* (FOREL). See p. 231.

Ecophylla smaragdina subsp. *fusca* var. *rubriceps* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 354 (♀).

Type locality: BELGIAN CONGO.

BELGIAN CONGO: Stanleyville (Lang and Chapin).

14. Var. **textor** (SANTSCHI).

Ecophylla smaragdina subsp. *longinoda* var. *textor* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 128, fig. 29a (♀, ♀).

Type locality: Mangapwani, ZANZIBAR ISLAND (Alluaud and Jeannel).

ZANZIBAR ISLAND: Bububu (Alluaud and Jeannel). BRITISH EAST AFRICA: Likoni; Cheteni; Tiwi; Gazi near Mombasa (Alluaud and Jeannel).

2. *Ecophylla smaragdina* (FABRICIUS) F. SMITH, 1861, Journ. Proc. Linn. Soc. London, Zool., V, p. 102 (♀). MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 695; 1863, *ibid.*, XIII, p. 438; 1867, Tijdschr. v. Ent., X, p. 70 (♀, ♀). FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 199 (♂). EMERY, 1887, Ann. Mus. Civ. Genova, XXIV, p. 242 (♀, ♀); 1888, Zeitschr. Wiss. Zool., XLVI, p. 379, Pl. xxvii, fig. 7. ROTHNEY, 1889, Trans. Ent. Soc. London, p. 358. EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 564 (♀, ♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 177. MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 197. H. STADELMANN, 1893, Mitth. Deutsch. Schutzgeb., VI, p. 217. EMERY, 1894, Mem. Accad. Sc. Bologna, (5) IV, p. 112 (♀, ♀). BINGHAM, 1903, 'Fauna Brit. India, Hym.,' II, p. 311, fig. 93 (♀, ♀, ♂); 1903, Ann. Mag. Nat. Hist., (7) XII, p. 62 (♂). FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 88 (♀); 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 371; 1913, Rev. Zool. Afr., II, p. 339 (♀). BEQUAERT, *ibid.*, p. 430. CARPENTER, 1920, 'A Naturalist on Lake Victoria,' pp. 100 and 283.

Formica smaragdina FABRICIUS, 1775, 'Syst. Ent.,' p. 828 (♀); 1781, 'Spec. Insect.,' I, p. 488 (♀); 1787, 'Mantissa Insect.,' I, p. 307. GMELIN, in Linnæus, 1790, 'Syst. Nat.,' Ed. 13, I, 5, p. 2797. CHRIST, 1791, 'Naturg. d. Insect.,' p. 505, Pl. LX, fig. 1. OLIVIER, 1791, 'Encycl. Méthod. Insect.,' VI, p. 491. FABRICIUS, 1793, 'Ent. Syst.,' II, p. 350 (♀). LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 176 (♀), Pl. III, fig. 18. FABRICIUS, 1804, 'Syst. Piez.,' p. 397. LEPELETIER, 1836, 'Hist. Nat. Ins. Hym.,' I, p. 218. JERDON, 1851, Madras Journ. Litt. Sc., XVII, p. 121 (♀, ♀, ♂); 1854, Ann. Mag. Nat. Hist., (2) XIII, p. 104 (♀, ♀, ♂). F. SMITH, 1857, Journ. Proc. Linn. Soc. London, Zool., II, p. 53; 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 17, Pl. III, fig. 3. TENNENT, 1861, 'Sketch Nat. Hist. Ceylon,' p. 422.

Formica macra GUÉRIN, in Duperrey, 1830, 'Voy. Coquille, Zool.,' II, 2, p. 202 (♀); Atlas, Ins., Pl. VIII, fig. 1.

Formica zonata GUÉRIN, *ibid.*, p. 205 (♀).

Ecophylla smaragdina var. *longinoda* EMERY, 1887, Ann. Mus. Civ. Genova, XXIV, p. 243 (*nec* Latreille).

Type locality: INDIA.

Indomalayan and Papuan Regions.

The occurrence of this species in the Ethiopian Region is doubtful. It was recorded from the following African localities, at the specimens in question may have been *longinoda*:

BELGIAN CONGO: Bukama (J. Bequaert). BRITISH EAST AFRICA: Mombasa (H. Prell); Pemba Island (Burt). GERMAN EAST AFRICA: Mafia Island (Voeltzkow); Amani; Uelenge (Zimmer). ZANZIBAR: (Stuhlmann). PORTUGUESE EAST AFRICA: Inhambane; Magarra River (Fornasini). UGANDA: Ngamba Island, Victoria Nyanza (Carpenter).

Camponotini Forel**Camponotus MAYR¹**

Camponotus MAYR, 1861, 'Europ. Formicid.,' p. 35.

Formica (part) LINNÆUS, etc.

Genotype: *Formica herculeana* Linnæus, 1758.

Subgenus 1. Myrmoturba FOREL

Camponotus subg. *Myrmoturba* FOREL, 1912, Mém. Soc. Ent. Belgique, XX, p. 91; 1914, Rev. Suisse Zool., XXII, p. 259. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 235.

Subgenotype: *Formica maculata* Fabricius, 1781.

1. Camponotus (Myrmoturba) acvapimensis MAYR.

Camponotus acvapimensis MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 664 (♂); 1863, ibid., XIII, p. 397. ROGER, 1863, 'Verzeich. Formicid.,' p. 2. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 220. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 150; in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 20 (♀). See p. 236.

Camponotus acvapimensis EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 365 (♂); 1897, ibid., XXXVIII, p. 604; 1896, Mem. Accad. Sc. Bologna, (5) V, p. 770.

Camponotus aquvapimensis EMERY, 1899, Ann. Mus. Civ. Genova, XXXIX, p. 501.

Camponotus akvapimensis L. REH, 1905, Zeitschr. f. Pflanzenkrankh., XV, p. 134.

Camponotus akvapimensis EMERY, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 63. FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 90 (♂, ♀); 1907, Ann. Mus. Nat. Hungarici, V, p. 31 (♂); 1907, Mitt. Naturh. Mus. Hamburg, XXIV, p. 16 (♀); 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 268 (♂); 1910, Ann. Soc. Ent. Belgique, LIV, p. 451 (♂); 1911, Rev. Zool. Afr., I, p. 281 (♀); 1911, Bull. Soc. Vaudoise Sc. Nat., (5) LXVII, p. 371 (♂). AULMANN and LA BAUME, 1912, 'Fauna d. Deutsch. Kolon.,' V, 3, p. 65. AULMANN, 1913, ibid., V, 5, p. 95. BEQUAERT, 1913, Rev. Zool. Afr., II, p. 430. FARQUHARSON, 1914, Trans. Ent. Soc. London, Proc., p. xxiv.

Camponotus (Myrmoturba) akvapimensis FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, pp. 145 and 355 (♂, ♀); 1913, Rev. Zool. Afr., II, p. 340 (♂, ♀, ♂); 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 253 (♂, ♂).

Camponotus (Myrmoturba) acvapimensis FOREL, 1914, Revue Suisse Zool., XXII, p. 266. SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 380 (♂). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 442 (♂, ♀).

Camponotus mombassæ FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 180 (♂). EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 119 (♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 243. EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 181. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 37.

Camponotus kersteni subsp. *mombassæ* EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 770.

¹*Camponotus angolensis* "Smith," recorded from Angola (Welwitsch) by Radoszkowsky, 1881, Journ. Sci. Ac. Lisboa, VIII, No. 31, p. 197, has apparently never been described. Dixey and Longstaff, 1907, Trans. Ent. Soc. London, p. 336, mention *Camponotus marginatus* (Latreille) as taken by them at Johannesburg, Transvaal; but this record is probably due to misidentification, since this is a holarctic species.

Camponotus (Myrmoturba) ackwapimensis EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 255.

Type locality: Akwapim Mountains, GOLD COAST.

FRENCH GUINEA: Kindia; Camayenne; Mamou (F. Silvestri); Los Islands. SIERRA LEONE. LIBERIA: Grand Bassa; Junk River (H. Brauns). SOUTHERN NIGERIA: Moor Plantation near Ibadan (Farquharson); Old Calabar (H. Brauns). CAMEROON: (H. Brauns); Victoria (Fickendey). BELGIAN CONGO: Banza Masola; Ganda Sundi; Mondombe; Congo da Lemba (R. Mayné); Yumbi; Zambi (J. Bequaert); Faradje; Garamba; Bolengi near Coquilhatville; Stanleyville; Vankerekhovenville; Akenge; Niangara (Lang and Chapin); Thysville (Lang, Chapin, and J. Bequaert); Eala. RHODESIA: Bulawayo (G. Arnold). GERMAN EAST AFRICA: Kigerama; Mombasa (Hildebrandt; H. Prell); Tanga (Zimmer); Mto-ya-Kifaru (Katona); Kilimanjaro (Sjöstedt); Patta; Manda. UGANDA: Chacansengula (Bayer). BRITISH EAST AFRICA: Fundu Island, W. Pemba (Voeltzkow). SOMALILAND: Lower Ganada (V. Bottego); Giari Bulé (Ruspoli); between Obbia and Berbera (Bricchetti-Rebecchi). ABYSSINIA: Arussi Galla, Ganale Gudda; Dimé to Bass Narok (V. Bottego); Bela; Coremma (Ruspoli); western Abyssinia (Ilg). ERITREA: Keren (Beccari); Asmara; Keren (Mancini and Ruggeri).

1. *Var. poultoni* (FOREL).

Camponotus ackwapimensis var. *poultoni* FOREL, 1913, Rev. Zool. Afr., II, p. 353 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 355 (♀). LAMBORN, 1914, Trans. Ent. Soc. London, (1913), p. 444. FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 348 (♀, ♂).

Camponotus (Myrmoturba) ackwapimensis var. *poultoni* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 380 (♀).

Camponotus (Myrmoturba) ackwapimensis var. *poultoni* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 442 (♀).

Type locality: Lagos, SOUTHERN NIGERIA (Lamborn).

SIERRA LEONE. CAMEROON: Victoria (F. Silvestri). BELGIAN CONGO: St. Gabriel (Kohl); Congo da Lemba (R. Mayné).

2. *Camponotus (Myrmoturba) arnoldinus* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 251 (♀).

Type locality: Durban, NATAL (G. Arnold).

3. *Camponotus (Myrmoturba) belligerum* SANTSCHI.

Camponotus belligerum SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 384, fig. 15 (♀).

Type locality: Ababis, GERMAN SOUTHWEST AFRICA (G. Arnold).

4. *Camponotus (Myrmoturba) bianconii* EMERY. FOREL, 1914, Rev. Suisse Zool., XXII, p. 267. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 255.

Camponotus bianconii EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 52 (♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 771.

Type locality: Cape Town, CAPE PROVINCE (E. Simon).

5. *Camponotus (Myrmoturba) cleobulus* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 238 (♀).

Type locality: Drakensberg, NATAL, 11,000 ft. (Haviland).

6. *Camponotus (Myrmoturba) maculatus* (FABRICIUS). FOREL, 1914, Rev. Suisse Zool., XXII, p. 267. DONISTHORPE, 1915, Ent. Record, XXVII, p. 221 (♂). EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 255. See pp. 234 and 1040.

Formica maculata FABRICIUS, 1781, 'Spec. Ins.,' I, p. 491 (♂); 1787, 'Mantissa Insect.,' I, p. 308 (♂). GMELIN, in LINNÆUS, 1790, 'Syst. Nat.,' Ed. 13, I, 5, p. 2800. OLIVIER, 1791, 'Encycl. Méthod. Insect.,' VI, p. 495. FABRICIUS, 1793, 'Ent. Syst.,' II, p. 356 (♂). LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 283. FABRICIUS, 1804, 'Syst. Piez.,' p. 403 (♂). LEPELETIER, 1836, 'Hist. Nat. Ins. Hym.,' I, p. 215 (♂). F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 28 (♂).

Camponotus maculatus MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 654 (♂, ♀, ♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 241. EMERY, 1915, Bull. Soc. Ent. France, p. 79.

Camponotus maculatus maculatus EMERY, 1905, Rend. Accad. Sc. Bologna, (1904-05), p. 37; 1908, Deutsch. Ent. Zeitschr., pp. 190 and 191 (♂, ♀, ♂).

Type locality: "in Africa equinoctiali" (probably SIERRA LEONE).

BELGIAN CONGO: Medje; Yakuluku; Garamba; Vankerkhovenville; Faradje (Lang and Chapin).

The distribution of the typical form cannot be traced at present. Probably one of the subspecies or varieties listed below is a synonym of it; perhaps such is the case for *C. maculatus* subsp. *melanocnemis* var. *lohieri* Santschi.

The following references of "*Camponotus maculatus*" probably include some of the forms and not the type.

Formica maculata GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 262; in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 509 (♂). A. GIRARD, 1881, Journ. Sci. Ac. Lisboa, VIII, No. 31, p. 227; in Capello and Ivens, 1882, 'From Benguela to the Territory of Yacca,' II, p. 294.

Camponotus maculatus ROGER, 1863, 'Verzeich. Formicid.,' p. 2. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 400; 1865, 'Reise Novara, Zool.,' II, Formicidae, p. 27. GERSTÄCKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 355; 1873, in 'v. d. Decken's Reisen in Ostafrika, Gliederthiere,' p. 344 (♂, ♀). EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 364 (♂, ♀, ♂). W. F. KIRBY, 1884, Ann. Mag. Nat. Hist., (5) XIII, p. 404. EMERY, 1890, Bull. Soc. Ent. Italiana, XXII, p. 56; 1892, Ann. Soc. Ent. France, LX, (1891), p. 566. FOREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 349 (♂). EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 119 (♂). MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 195. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 46; 1895, Ann. Mus. Civ. Genova, XXXV, p. 180; 1896, Mem. Accad. Sc. Bologna, (5) V, p. 768 (♂, ♀); 1896, Ann. Mus. Civ. Genova, XXXVII, p. 158 (♂, ♂); 1897, ibid., XXXVIII, p. 604. H. STADELMANN, 1898, 'Deutsch Ost-Afrika,' IV, Hym., p. 36. EMERY, 1899, Ann. Mus. Civ. Genova, XXXIX, p. 501. BINGHAM, 1903, Ann. Mag. Nat. Hist., (7) XII, p. 62 (♂, ♀, ♂). MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 11; 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. 1, p. 389; in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 19 (♂, ♀). FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 31 (♂); 1907, Mitt. Naturh. Mus. Hamburg, XXIV, p. 16 (♂). F. F. KOHL, 1907, Denkschr. Ak. Wiss. Wien, LXXI, p. 282 (♂, ♀). DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, pp. 310, 320, 327, 336, 341, 363, and 369. STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 148 (♂). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 26 (♂, ♀, ♂); 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 266 (♂). STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 392 (♂). FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 371 (♂); 1913, Rev. Zool. Afr., II, p. 340 (♀). REICHENSPERGER, 1913, Zool. Jahrb. Abt. Syst., XXXV, pp. 198 and 215.

Camponotus sylvaticus subsp. *maculatus* FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 64.

Camponotus sylvaticus EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 525 (♂, ♀). MAGRETTI, 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 244 (♂, ♀). BINGHAM, 1903, Ann. Mag. Nat. Hist., (7) XII, p. 62 (♂, ♀).

Camponotus sylvaticus var. *maculatus* MAGRETTI, 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 244 (♂, ♀); 1884, Ann. Mus. Civ. Genova, XXI, p. 536 (♂, ♀). LONGSTAFF, 1911, Ent. Monthly Mag., XLVII, p. 124 (♂).

Camponotus rubripes subsp. *maculatus* FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 149.

Camponotus (Myrmoturba) maculatus FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 249 (♂). SANTSCHI, 1914, Bull. Lab. Zool. Gen. Agrar. Portici, VIII, p. 379 (♂, ♀); 1914, Deutsch. Ent. Zeitschr., p. 288 (♂); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 129 (♂, ♂). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 403 (♂).

Camponotus maculatus subsp.? LAMBORN, 1914, Trans. Ent. Soc. London, (1913), p. 444.

SENEGAMBIA: Dakar (C. Alluaud); Thiès (F. Silvestri). SOUTHERN NIGERIA: Oni Camp east of Lagos (Lamborn). CAMEROON: Duala (v. Rothkirch). FRENCH CONGO: Fort Archambault (Schubotz). BELGIAN CONGO: Kiambi (J. Bequaert); Kasindi (Schubotz). RHODESIA: (G. Arnold); Sheshong Road Station (Dixey and Longstaff). BECHUANALAND: Khakhea (L. Schultze); Artesia (Dixey and Longstaff). GERMAN SOUTHWEST AFRICA: (Lübbert; Kunze); Okahandja (Casper; Dinter). CAPE PROVINCE: (Wilms); Cape Town; Kimberley (E. Simon); King William's Town (R. Godfrey); East London (Dixey and Longstaff). ORANGE FREE STATE: Bloemfontein (E. Simon). TRANSVAAL: Makapan; Hamman's Kraal (E. Simon); Pretoria (Distant); Hebron (E. Simon); Barberton (Rendall); Lydenburg (Wilms); Johannesburg (Dixey and Longstaff; Distant). NATAL: The Bluff, Durban; Sydenham, Durban (Dixey and Longstaff). MOZAMBIQUE: (Peters). ZANZIBAR: (Stuhlmann); Bawi Island near Zanzibar (Stuhlmann). GERMAN EAST AFRICA: Kilimanjaro (Bornemisza); Ulenge Island (H. Prell); Bulko; Uluguru Mts. (Zimmer); Meru (Sjöstedt); Tanga (Alluaud and Jeannel). BRITISH EAST AFRICA: Mombasa (v. d. Decken; Alluaud and Jeannel); Shimoni; Tiwi (Alluaud and Jeannel); Lamu Island; Fundu Island near Pemba; Manda Island; Patta Island (Voeltzkow). SOMALILAND: (C. Keller); Obbia (Bricchetti-Robecchi); Milmil (Pavesi); wells of Laffarugh to wells of Aberio (V. Bottego); Magala Re Umberto (Ruspoli); Lugh; Matagoi to Lugh (V. Bottego). ABYSSINIA: Ganale; Bela; Daua; Ogaden (Ruspoli); Webi (Bricchetti-Robecchi); Sencurar to Amarr Burgi; Lake Bass Narok (V. Bottego). ERITREA: Massaua; Assab (Doria and Beccari); Ghinda (K. Escherich). ANGLO-EGYPTIAN SUDAN: Kaka, White Nile (I. Trägårdh); Assuan; Khor Attar; Gondokoro (F. Werner); Metemma; Kassala; Sebderat (Magretti); Khartum (Longstaff). SOKOTRA: Ras Shoab. SOUTHERN ARABIA: Aden (O. Simony); Gischin (W. Hein).

6. Var. *thomensis* SANTSCHI, 1920, 'Études Maladies Parasites Cacaoyer S. Thomé,' X, p. 3 (♂, ♀, ♂).

Type locality: SAN THOMÉ (de Seabra).

6a. Subsp. *adenensis* EMERY, 1893, Ann. Soc. Ent. France, LXII, p. 257 (♀, ♀); 1895, Ann. Soc. Ent. Belgique, XXXIX, p. 337; 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 604.

Type locality: Aden, southern ARABIA (E. Simon; Doria; C. Alluaud).

ABYSSINIA: Salole (Ruspoli).

6a. Var. *assabensis* EMERY, 1893, Ann. Soc. Ent. France, LXII, p. 258 (♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 768.

Type locality: Assab, ERITREA.

6b. Subsp. *egyptiacus* EMERY, 1915, Bull. Soc. Ent. France, p. 79; 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 22 (♀).

Type locality: EGYPT.

ERITREA.

6c. Subsp. *agricola* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 454 (♀).

Type locality: Mountains of NATAL (R. C. Wroughton).

6c. Var. *tacitus* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 240 (♀).

Type locality: NATAL (Haviland).

6d. Subsp. *atramentarius* FOREL, 1905, Ann. Soc. Ent. Belgique, XLIX, p. 180. LICHTWARDT, 1908, Deutsch. Ent. Zeitschr., p. 338. FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 26 (♀, ♀); 1911, Rev. Zool. Afr., I, p. 280 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 131 (♀). BUCHNER, 1918, Arch. Microsc. Anat., XCI, Abt. 2, p. 78, footnote.

Camponotus maculatus var. *atramentarius* FOREL, 1904, Ann. Mus. Zool. Ac. Sc. St. Pétersbourg, VIII, (1903), p. 379 (♀).

Type locality: ABYSSINIA (Dmitriev).

NATAL: Durban (Wilms). MOZAMBIQUE: Lobombo Borges (Wilms). BELGIAN CONGO: Uele-Bili. UGANDA: M'Bale (C. Alluaud). BENGUELA (C. Wellman).

6d. Var. *cluis* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 67 (♀); 1911, Rev. Zool. Afr., I, p. 280 (♀).

Type locality: BENGUELA (C. Wellman).

East Africa (between the Great Lakes).

6d. Var. *cluisoides* (FOREL).

Camponotus maculatus var. *cluisoides* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 354 (♀).

Type locality: Campo Tembo, Tsavo, BRITISH EAST AFRICA (Bayer).

6e. Subsp. *brutus* (FOREL) MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 150 (♂); 1896, Ent. Tidskr., XVII, p. 249. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 769; 1899, Ann. Soc. Ent. Belgique, XLIII, p. 501 (♀). FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 69 (♀). SJÖSTEDT, 1904, 'I Västafrikas Urskogar,' p. 507. FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 54 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393 (♀, ♀). STRITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 148 (♀). FOREL, 1911, Rev. Zool. Afr., I, p. 280 (♀); 1913, *ibid.*, II, p. 340 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 355 (♀, ♀, ♂); 1914, Rev. Suisse Zool., XXII, p. 267. SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 379 (♀). STRITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 403 (♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 442 (♀, ♀, ♂). BUCHNER, 1918, Arch. Microsc. Anat., XCI, Abt. 2, p. 78, footnote. See p. 236.

Camponotus rubripes subsp. *brutus* FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 155 (♀, ♀).

Camponotus brutus DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 223. L. REH, 1905, Zeitschr. f. Pflanzenkrankh., XV, p. 134 (♀). AULMANN and LA BAUME, 1912, 'Fauna d. Deutsch. Kolon.,' V, 3, p. 62. AULMANN, 1913, *ibid.*, V, 5, p. 92.

Camponotus (Myrmoturba) brutus EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 255.

Type locality: Chinchoxo, PORTUGUESE CONGO (Falkenstein).

SIERRA LEONE: Samlia Falls, River N'Gamie (Mocquers). LIBERIA: FERNANDO PO: (Schultze). CAMEROON: (Sjöstedt; H. Brauns); Victoria (F. Silvestri; H. Winkler); Mundame (Conradt); Nssanakang (Rudatis); Bibundi; Mokundange (Tessmann); Yukaduma (Schultze). SPANISH GUINEA: Alen (Tessmann). FRENCH CONGO: Brazzaville (A. Weiss); Sibange; Sette-Cama (Soyaux; Hupfer). BELGIAN CONGO: (Kohl); Yabena Mabote, Lomami-Kondué (Luja); Lukula (Daniel); Beni (Borgerhoff); Dungu, Mayombe (Deleval); Kiniati; Congo da Lemba (R. Mayné); Mayombe (de Briey); Avakubi; Medje; Faradje; Bafwasende; Stanleyville; Batama; Lukolela; Malela; Isangi; Nouvelle Anvers; Zambi; Poko; Akenge; Niangara (Lang and Chapin); Malela (J. Bequaert); Kimuenza (Schultze); Libenge (Schubotz).

6e. Var. *lycurgus* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 501 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393 (♀). See p. 236.

Type locality: CAMEROON.

IVORY COAST. FRENCH CONGO: Brazzaville (A. Weiss); Ogowe. BELGIAN CONGO: Leopoldville (Lang and Chapin).

6f. Subsp. *cavallus* SANTSCHI, 1911, Rev. Zool. Afr., I, p. 211 (♀, ♀). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 403 (♀). SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 347, fig. 4b (♀).

Type locality: Cucala, BENGUELA (J. Cruchet).

BELGIAN CONGO: Duma (Schubotz).

6g. Subsp. *cognatus* (F. SMITH) EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 566 (♀). ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 1 (♀). MAYR, 1896, Ent. Tidskr., XVII, p. 249. FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 26 (♀, ♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 148 (♀).

Formica cognata F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 35 (♀).

Camponotus rubripes ROGER, 1863, Berlin. Ent. Zeitschr., VII, p. 129 (♀, ♀, ♂); 1863, 'Verzeich. Formicid.,' p. 2. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 401. FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 143.

Camponotus sylvaticus subsp. *cognatus* FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 65.

Camponotus sylvaticus subsp. *cognatus* var. *rubripes* FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 66 (♀).

Camponotus sylvaticus var. *cognatus* MAGRETTI, 1884, Ann. Mus. Civ. Genova, XXI, p. 536 (♀); 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 244 (♀).

Camponotus rubripes subsp. *cognatus* FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 151; 1890, *ibid.*, XXXIII, C. R., p. lv (♀).

Camponotus maculatus var. *cognatus* MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 195. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 36. ZAVATTARI, 1909, 'Il Ruwenzori, Parte Scientif.,' I, p. 215 (♀, ♂).

Camponotus cognatus DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 225. WASMANN, 1899, Deutsch. Ent. Zeitschr., p. 411. BERGROTH, 1915, Wien. Ent. Zeitg., XXXIV, p. 291.

Camponotus maculatus subsp. *carinatus* var. *cognatus* EMERY, 1894, Mem. Accad. Sc. Bologna, (5) IV, p. 112 (♂, ♀); 1895, Ann. Soc. Ent. France, LXIV, p. 46.

Type locality: Port Natal, NATAL.

IVORY COAST: Assinie (C. Alluaud). TOGO: Bismarckburg (Conradt). FERNANDO PO: (Conradt). FRENCH CONGO: Ogowé (Mocquereys). GERMAN SOUTHWEST AFRICA: (Lübbert). CAPE PROVINCE: Kamaggas; Cape Flats, Bergvliet (L. Schultze); Cape Town (E. Simon); Port Elizabeth (H. Brauns). MOZAMBIQUE: Inhambane (Fornasini); Quillimane (Stuhlmann). UGANDA: Sanda; Katende; Mitiana; Butiti (Duke of Abruzzi). ANGLO-EGYPTIAN SUDAN: Suakin; Kor Langhebb (Magretti).

69i. Var. *ballioni* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 27.

Camponotus maculatus subsp. *ballioni* FOREL, 1904, Ann. Soc. Ent. Belgique, XLVIII, p. 176 (♂).

Type locality: Cape of Good Hope.

CAPE PROVINCE: Port Elizabeth (H. Brauns).

692. Var. *boerus* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 27 (♂, ♀).

Type locality: Steinkopf, CAPE PROVINCE (L. Schultze).

CAPE PROVINCE: Kamaggas (L. Schultze).

6A. Subsp. *congolensis* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 499, fig. (♂, ♀). FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 69 (♀); 1909, Ann. Soc. Ent. Belgique, LIII, p. 54 (♂, ♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393 (♂, ♀). FOREL, 1913, Rev. Zool. Afr., II, p. 340 (♂, ♀, ♂). BEQUAERT, *ibid.*, p. 430. FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 250 (♂). BUCHNER, 1918, Arch. Microsc. Anat., XCI, Abt. 2, p. 78, footnote. See p. 235.

Type locality: Banana, BELGIAN CONGO (Busschodts).

WEST AFRICA: (H. Freyschmidt). LIBERIA: Cape Mesurado. FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: Kinshasa (Waelbroeck); Boma Sudi (Rollin); Kisantu (J. Bequaert); Beni (Bergerhoff); Yakuluku; Garamba; Faradjé; Medje; Niangara (Lang and Chapin). RHODESIA: Bindura (G. Coqhill).

6i. Subsp. *desantii* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 280, figs. 14a-c (♂, ♀, ♂).

Type locality: Kouandé, UPPER DAHOMEY (Desanti).

6j. Subsp. *fornasini* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 46 (♂); 1896, Mem. Accad. Sc. Bologna, (5) L, p. 769. FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 288 (♂).

Camponotus (Myrmoturba) fornasini FOREL, 1914, Rev. Suisse Zool., XXII, p. 267. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 255.

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA.

NATAL: (Bates).

76k. Subsp. *guttatus* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 498 (♂, ♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393 (♂). FOREL, 1913, Rev. Suisse Zool., XXI, p. 670 (♂, ♀, ♂). See p. 234.

Type locality: CAMEROON (Conradt).

FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: Zambi (Lang, Chapin, and J. Bequaert).

Type locality: Patta Island, BRITISH EAST AFRICA (Voeltzkow).

GERMAN EAST AFRICA: Bukoba, Victoria Nyanza (Schubotz).

12. *Camponotus (Myrmoturba) werthi* FOREL, 1914, Rev. Suisse Zool., XXII, p. 268; 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 253 (♀).

Camponotus werthi FOREL, 1908, Bull. Soc. Vaudoise Sc. Nat., (5) XLIV, p. 18 (♀). ENDERLEIN, 1909, 'Deutsch. Südpolar-Expedition,' X, 4, pp. 340, 393, and 426 (♀).

Type locality: KERGUELEN (Werth); probably imported from the Cape, since it was found in a house only.

CAPE PROVINCE: Caledon (E. P. Phillips). CROZET ARCHIPELAGO: Possession Island (Werth).

Subgenus 2. *Dinomyrmex* ASHMEAD

Dinomyrmex ASHMEAD, 1905, Canadian Ent., XXXVII, p. 384.

Camponotus subg. *Myrmogigas* FOREL, 1912, Mém. Soc. Ent. Belgique, XX, p. 91.

Camponotus subg. *Dinomyrmex* FOREL, 1914, Rev. Suisse Zool., XXII, p. 259. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 236.

Subgenotype: *Formica gigas* Latreille, 1802.

13. *Camponotus (Dinomyrmex) æquatorialis* ROGER. FOREL, 1914, Rev. Suisse Zool., XXII, p. 268. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256.

Camponotus æquatorialis ROGER, 1863, Berlin. Ent. Zeitschr., VII, p. 135 (♀); 1863, 'Verzeich. Formicid.,' p. 2. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 458. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 220.

Camponotus maculatus ? subsp. *æquatorialis* EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 769.

Camponotus æquatorialis SANTSCHI, 1911, Rev. Zool. Afr., I, p. 211 (♀).

Type locality: Gaboon, FRENCH CONGO.

FRENCH CONGO: (A. Weiss).

13a. Subsp. *kohli* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 349 (♀).

Type locality: Bengamisa, BELGIAN CONGO (Kohl).

14. *Camponotus (Dinomyrmex) brevicollis* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 401, fig. 13 (♀).

Type locality: between Fort de Possel and Fort Crampel, FRENCH CONGO (Schubotz).

15. *Camponotus (Dinomyrmex) cæsar* FOREL, 1914, Rev. Suisse Zool., XXII, p. 268. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256. See p. 243.

Camponotus cæsar FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 162 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 223. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 768. FOREL, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 503, fig. (♂, ♀); 1909, *ibid.*, LIII, p. 66 (♀). EMERY, 1912, Annuar. Mus. Zool. Univ. Napoli, N. S., III, No. 26, p. 2 (♀).

Type locality: ANGOLA (M. Buchner).

BENGUELA: (C. Wellman). BELGIAN CONGO: Faradje (Lang and Chapin); Luapula River.

15a. Subsp. *imperator* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 503, fig. (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 149 (♀); 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 391 (♀). See p. 244.

Type locality: CAMEROON (Conradt).

Camponotus (Myrmoturba) natalensis subsp. *fulvipes* FOREL, 1914, Rev. Suisse Zool., XXII, p. 267.

Type locality: Makapan, TRANSVAAL (E. Simon).

ZULULAND: (I. Trägårdh).

85. Subsp. *diabolus* FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 81 (?). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 771.

Camponotus diabolus DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 228.

Camponotus (Myrmoturba) diabolus FOREL, 1914, Rev. Suisse Zool., XXII, p. 267.

Type locality: Lessouto, BASUTOLAND (P. Berthoud).

9. *Camponotus (Myrmoturba) roubaudi* SANTSCHI. FOREL, 1913, Rev. Zool. Afr., II, p. 340 (?); 1914, Rev. Suisse Zool., XXII, p. 267. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 255.

Camponotus roubaudi SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 365, fig. (5).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss and Roubaud).

BELGIAN CONGO: Beni (Borgerhoff).

91. Var. *agonia* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 282 (?).

Type locality: Lugombe, BELGIAN CONGO (Gérard).

10. *Camponotus (Myrmoturba) sacchii* EMERY. FOREL, 1914, Rev. Suisse Zool., XXII, p. 267. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 255.

Camponotus sacchii EMERY, 1899, Ann. Mus. Civ. Genova, XXXIX, p. 501 (?).

Type locality: Buditu to Dimé, ABYSSINIA (V. Bottego).

11. *Camponotus (Myrmoturba) somalinus* ERN. ANDRÉ. FOREL, 1914, Rev. Suisse Zool., XXII, p. 267. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 132 (?). EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 255.

Camponotus somalinus ERN. ANDRÉ, 1887, Rev. d'Ent. Caen, VI, p. 280 (?). EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 120. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 253. MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 195. EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 180. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 150. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 768; 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 604. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 37. EMERY, 1899, Ann. Mus. Civ. Genova, XXXIX, p. 500; 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 63. MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 11; in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 20. FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 371 (?).

Type locality: SOMALILAND.

EGYPT. SOMALILAND: Magala Re Umberto (Ruspoli); Mogadiscio; Obbia; between Obbia and Berbera (Bricchetti-Robecchi). ABYSSINIA: Sannur to Amarr Burgi; Dimé to Bass Narok; Arussi Galla, Ganale Gudda (V. Bottego). ERITREA: Asmara (Mancini and Ruggeri). BRITISH EAST AFRICA: Taveta; Mombasa (C. Alluaud). GERMAN EAST AFRICA: Tanga (Zimmer); Mbusini, Usegua (Stuhlmann); Daressalaam (H. Brauns); Kahe; Mt. Kilimanjaro (Sjöstedt).

111. Var. *curtior* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 65 (?). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 768.

Type locality: SOUTHERN ABYSSINIA (Ilg).

112. Var. *pattensis* FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 91 (?).

Camponotus somalicus var. *pattensis* STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 392 (?).

Camponotus sexpunctatus var. *liengmei* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 67 (♀).

Camponotus maculatus subsp. *sexguttatus* var. *liengmei* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 46.

Camponotus maculatus subsp. *sexpunctatus* var. *liengmei* EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 769.

Camponotus maculatus var. *liengmei* FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 88 (♂, ♂).

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA (Liengme).

NATAL: Durban (G. Arnold). TRANSVAAL: Pretoria (E. Simon). CAPE PROVINCE: Kimberley (E. Simon); Algoa Bay (H. Brauns). BRITISH EAST AFRICA: Fundu Island, W. Pemba (Voeltzkow). ZANZIBAR: (Voeltzkow).

6r. Var. *hansingi* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 452 (♀); 1911, Sitzb. Bayer. Akad. Wiss., p. 288; 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 250 (♀).

Type locality: Beira, PORTUGUESE EAST AFRICA (H. Hansing).

ORANGE FREE STATE: Reddersburg (H. Brauns). CAPE PROVINCE: Willowmore (G. Arnold).

6s. Var. *importunoides* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 249 (♀).

Type locality: Smithwinkle Bay, CAPE PROVINCE (E. P. Phillips).

6s. Subsp. *manzer* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 452 (♀); 1911, Rev. Zool. Afr., I, p. 280 (♀?). SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 509 (♂, ♂).

Type locality: Victoria Falls, RHODESIA (R. C. Wroughton; G. Arnold).

GERMAN EAST AFRICA: Kigerama and Weranjanje, near Kagera (between the great lakes).

6s. Var. *contaminatus* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 291 (♀).

Type locality: Matoppe Hills, SOUTHERN RHODESIA (G. Arnold).

6t. Subsp. *mathildae* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 266 (♀, ♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 130 (♀).

Type locality: Zoutpansberg, 800 m., TRANSVAAL (Rolle).

BRITISH EAST AFRICA: Kenia Region, 1800-2000 m.; Amboni River, 1800 m. (Alluaud and Jeannel).

6u. Subsp. *melanocnemis* (SANTSCHI) FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 355 (♀); 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 250 (♀). SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 278, fig. 14d (♀). See p. 234.

Camponotus maculatus var. *melanocnemis* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 368 (♀).

Type locality: FRENCH CONGO (H. Pobeguïn).

FRENCH CONGO: Grand Lahou (H. Pobeguïn). BELGIAN CONGO: Ubangi (Augustin); Dungu to Niangara, village of Denge (Hutereau); Faradje; Yakuluku (Lang and Chapin). NATAL: Durban (G. Arnold).

6u. Var. *flavominor* SANTSCHI, 1920, 'Études Maladies Parasites Cacaoyer S. Thomé,' X, p. 4, footnote (♀).

Type locality: Kouandé, UPPER DAHOMEY (Desanti).

6l. Subsp. **hannæ** SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 349, fig. 4a (♀).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

6m. Subsp. **hieroglyphicus** SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 290 (♀, ♀).

Type locality: Caconda, BENGUELA (J. Cruchet).

6n. Subsp. **importunus** FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 288 (♀, ♀, ♂).

Camponotus maculatus subsp. *serpunctatus* var. *importunus* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 27 (♀).

Type locality: CAPE PROVINCE (Krebs).

CAPE PROVINCE: Matjesfontein (Wilms); Algoa Bay; Port Elizabeth (H. Brauns). MOZAMBIQUE: Lobombo Borges (Wilms).

6o. Subsp. **intonsus** EMERY, 1905, Rend. Accad. Sc. Bologna, N. S., IX, p. 29, footnote (♀).

Type locality: CAPE PROVINCE.

TRANSVAAL.

6p. Subsp. **kersteni** (GERSTÄCKER) FOREL, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 309 (♀); 1914, Rev. Suisse Zool., XXII, p. 267.

Camponotus kersteni GERSTÄCKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 355 (♀); 1873, in 'v. d. Decken's Reisen in Ost-Afrika, Gliederthiere,' p. 344 (♀; nec ♂). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 770. H. STADÉLMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 37.

Camponotus kerstenii DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 237.

Camponotus (Myrmoturba) kersteni SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 131 (♂, ♀, ♀, ♂). EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 255.

Type locality: Mt. Kilimanjaro, 8000 ft., GERMAN EAST AFRICA (v. d. Decken).

GERMAN EAST AFRICA: Mt. Kilimanjaro, 2740-3000 m. (C. Alluaud). Dalla Torre's record "Zanzibar" is erroneous.

6q. Subsp. **lacteipennis** (F. SMITH) EMERY, 1896, Mem. Accad. Sc. Bologna, (5) I, p. 768. FOREL, 1914, Rev. Suisse Zool., XXII, p. 267.

Formica lacteipennis F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 34 (♀, ♀, ♂). MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 416.

Camponotus lacteipennis ROGER, 1863, 'Verzeich. Formicid.,' p. 2. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 237.

Camponotus sylvaticus subsp. *lacteipennis* MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 355.

Camponotus (Myrmoturba) lacteipennis EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 255.

Type locality: Port Natal, NATAL.

TRANSVAAL: Makapan; Pretoria (E. Simon).

6r. Subsp. **liengmei** FOREL, 1897, Abhandl. Senckenberg. Naturf. Ges., XXI, p. 188 (♀, ♀); 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 69 (♀, ♀); 1914, Rev. Suisse Zool., XXII, p. 267; 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 249 (♀).¹

¹Santschi (1921) removes this to *Dinomyrmex*, in which case it should be treated as a variety of *C. serpunctatus* Forel, as originally described.

6us. Var. **lohieri** SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 313 (♂); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 379 (♂, ♀); 1915, Ann. Soc. Ent. France, LXXXIV, p. 279 (♂).

Camponotus (Myrmoturba) maculatus var. **lohieri** EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 22 (♂).

Type locality: Jacqueville, IVORY COAST (Lohier).

SENEGAL. FRENCH GUINEA: Konakry (F. Silvestri). UPPER DAHOMEY: Kouandé (Desanti). SAN THOMÉ: (Gravier). ERITREA: Asmara (F. Silvestri).

6us. Var. **schultzei** (FOREL) SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 279 (♀).

Camponotus maculatus var. **schultzei** FOREL, 1912, Mitt. Naturh. Mus. Hamburg, XXIX, p. 179 (♂, ♂).

Camponotus (Myrmoturba) maculatus subsp. **schultzei** SANTSCHI, 1914; 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 130 (♂).

Type locality: Entebbe, UGANDA (Schultze).

UGANDA: Western Unyoro, between Hoima and Butiaba; Mt. Ruwenzori, Ibanda, 1400 m. (C. Alluaud). BRITISH EAST AFRICA: Kisumu, Victoria Nyanza (C. Alluaud).

6us. Var. **semispicatus** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 279 (♂).

Camponotus (Myrmoturba) maculatus subsp. **mathildeae** var. **semispicata** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 130 (♂).

Type locality: Nairobi, Kikuyu, 1700 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

BRITISH EAST AFRICA: Naivasha in the Rift Valley, 1900 m.; Gilgil, 1980 m.; Nakuru, 1280 m.; Kijabé on the Kikuyu Escarpment, 2100 m.; Masai Steppe near Nairobi (Alluaud and Jeannel).

6e. Subsp. **minusculus** VIEHMEYER, 1914, Arch. f. Naturg., LXXIX, Abt. A, Heft 12, (1913), p. 44 (♂, ♀). SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 282, fig. 15a (2).

Type locality: Manow, Langenburg, GERMAN EAST AFRICA.

6w. Subsp. **miserabilis** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 379 (♂); 1915, Ann. Soc. Ent. France, LXXXIV, p. 281, fig. 15b (♂).

Type locality: Kindia, FRENCH GUINEA (F. Silvestri).

6w1. Var. **pessimus** WM. M. WHEELER. See p. 235 (♂).

Type locality: Yakuluku, BELGIAN CONGO (Lang and Chapin).

6z. Subsp. **negus** FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 142 (♂, ♀, ♂).

Type locality: Kounhi, southern ABYSSINIA (de Rothschild).

ABYSSINIA: Abou; Mt. Zyoual near the Colba River; Uomber (de Rothschild).

BRITISH EAST AFRICA: Rendile, Mt. Karoli (de Rothschild).

6z1. Var. **beritchi** FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 348 (♂).

Type locality: George, CAPE PROVINCE (H. Brauns).

6z2. Var. **etiennel** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 453 (♂).

Type locality: Banana, BELGIAN CONGO (Etienne).

6z3. Var. **nefassitensis** FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 266 (♂); 1911, Sitzb. Bayer. Akad. Wiss., p. 290 (♂, ♀, ♂). KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 11 (♂, ♀).

Camponotus (Myrmoturba) maculatus subsp. *negus* var. *nefasitensis* EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 4 (♂).

Type locality: Nefasit, ERITREA (K. Escherich; F. Silvestri).

EGYPT: Port Sudan (Karawaiew).

6y. Subsp. *pictiventris* MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 28

(♀). FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 27 (♀, ♀).

Type locality: Port Elizabeth, CAPE PROVINCE (H. Brauns).

CAPE PROVINCE: Kamaggas (L. Schultze).

6y. Var. *immaculifrons* SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 349 (♂).

Type locality: Willowmore, CAPE PROVINCE (H. Brauns).

6y. Var. *octomaculatus* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 452 (♂).

Type locality: NATAL, 5300 ft. (R. C. Wroughton).

6z. Subsp. *pulvinatus* MAYR, in Sjöstedt, 1907, 'Kilimandjaro, Meru Exped.,' II, 8, p. 20 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 130 (♀).

Type locality: Kibonoto, Mt. Kilimanjaro, GERMAN EAST AFRICA (Sjöstedt).

UGANDA: Western Unyoro, between Hoima and Butiaba; Lake Albert (C. Alluaud).

6a'. Subsp. *radamoides* (FOREL). See p. 1042.

6a'. Var. *cataractæ* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 239 (♀).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

6a'. Var. *diffusus* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), pp. 292 and 293 (♀).

Type locality: Matoppo Hills, SOUTHERN RHODESIA (G. Arnold).

NATAL: (v. Muralt; I. Trägårdh).

6a'. Var. *liocnemis* (EMERY) SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 293 (♀).

Camponotus maculatus subsp. *lacteipennis* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 46 (♀).

Camponotus maculatus subsp. *liocnemis* EMERY, 1905, Rend. Accad. Sc. Bologna, N. S., IX, p. 30, footnote (♀).

Camponotus maculatus subsp. *atramentarius* var. *liocnemis* FOREL, 1907, Mitt. Naturh. Mus. Hamburg, XXIV, p. 16; 1909, Ann. Soc. Ent. Belgique, LIII, p. 54; in Schultze, 1910, 'Forschungsreise in Südafrika,' p. 26 (♀); 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 371 (♀); 1913, Rev. Zool. Afr., II, p. 340 (♀). BEQUAERT, *ibid.*, p. 430.

Camponotus (Myrmoturba) maculatus subsp. *liocnemis* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 379 (♀, ♀).

Camponotus (Myrmoturba) maculatus subsp. *liocnemis* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 38 (♀, ♀, ♂). TRÄGÅRDH, *ibid.*, p. 46.

Type locality: Makapan, TRANSVAAL (E. Simon).

BELGIAN CONGO: Welgelegen (J. Bequaert). ANGOLA: Quifangondo (F. Silvestri). GERMAN SOUTHWEST AFRICA: (Lübbert; Casper). BASUTOLAND. TRANSVAAL: Pretoria (E. Simon). NATAL: Greyminie; Stamford Hill (I. Trägårdh). ZULULAND: (I. Trägårdh). MOZAMBIQUE. GERMAN EAST AFRICA: Buiko.

6a'. Var. **madecassus** (EMERY) SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), pp. 292 and 293 (♀). See p. 1042.

NATAL: (I. Trägårdh).

6b'. Subsp. **schereri** FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 289 (♀, ♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 379 (♀); 1915, Ann. Soc. Ent. France, LXXXIV, p. 280.

Type locality: Nebena, LIBERIA (Scherer).

SOUTHERN NIGERIA: Olokemeji (F. Silvestri). FRENCH GUINEA: Kakoulima (F. Silvestri).

6c'. Subsp. **solon** (FOREL) EMERY, 1896, Mem. Accad. Sc. Bologna, (5) L, p. 769; 1899, Ann. Soc. Ent. Belgique, XLIII, p. 500 (♀, ♀). FOREL, 1909, *ibid.*, LIII, pp. 54 (♀) and 73 (♀, ♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393 (♀, ♀). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 452 (♀). STRITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 149 (♀, ♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 255 (♀); 1914, Rev. Suisse Zool., XXII, p. 267; 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 348 (♀); 1916, Rev. Suisse Zool., XXIV, p. 442 (♀). See p. 235.

Camponotus rubripes subsp. **solon** FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 152 (♀).

Camponotus solon DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 253.

Camponotus (Myrmoturba) solon EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 255.

Type locality: Malange, ANGOLA (M. Buchner).

CAMEROON: Mundame (Conradt). FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: Katanga (Lemaire); Kinshasa (Waelbroeck); Lumbulumbu; St. Gabriel (Kohl); Kimpoko (Büttner); Sankuru (Luja); Nyangwe to Stanleyville (Fauconnet); Boma (Styczinski), Medje; Niangara; Akenge; Bafwabaka (Lang and Chapin).

6c'. Var. **chilon** FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 68 (♀, ♀); 1911, Sitzb. Bayer. Akad. Wiss., p. 288.

Camponotus (Myrmoturba) maculatus subsp. **brutus** var. **chilon** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 379 (♀).

Type locality: Senegal.

SENEGAMBIA: Thiès (F. Silvestri). DAHOMEY: (Martienssen). SOUTHERN NIGERIA: Old Calabar (Bates).

6c'. Var. **jugurtha** Wm. M. WHEELER. See p. 236 (♀).

Type locality: Batama, BELGIAN CONGO (Lang and Chapin).

6d'. Subsp. **thales** FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 453 (♀, ♀, ♂); 1913, *ibid.*, LVII, p. 145 (♀).

Type locality: BASUTOLAND (R. C. Wroughton).

CAPE PROVINCE: Reddersburg (H. Brauns).

6d'. Var. **empedocles** FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 224 (♀).

Type locality: Salisbury, SOUTHERN RHODESIA (G. Arnold).

6e'. Subsp. **thoracicus** (FABRICIUS).

Formica thoracica FABRICIUS, 1804, 'Syst. Piez.', p. 397 (♀). ROGER, 1862, Berlin. Ent. Zeitschr., VI, p. 285.

Type locality: BARBARY (Stub).

6e'. Var. **fellah** EMERY, 1908, Deutsch. Ent. Zeitschr., p. 194 (♀, ♀, ♂).

Camponotus maculatus subsp. *oasium* var. *fellah* EMERY, 1891, 'Explor. Tunisie, Fourmis,' p. 18 (♀).

Camponotus maculatus subsp. *fellah* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 267 (♀).

Type locality: EGYPT.

SYRIA. WESTERN ABYSSINIA: (Ilg).

6c'. Var. *incommodus* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 250 (♀).

Type locality: CAPE PROVINCE.

6f'. Subsp. *trægaordhi* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 38 (♀, ♀, ♂). TRÄGÅRDH, *ibid.*, p. 46.

Type locality: Stamford Hill, NATAL (I. Trägårdh).

NATAL: Durban (I. Trägårdh). ZULULAND: Dukudu (I. Trägårdh).

6f'. Var. *muger* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 250 (♀).

Type locality: Durban, NATAL (H. B. Marley).

6g'. Subsp. *weissi* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 368 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 355 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

BELGIAN CONGO: Boma (Styczinski).

7. *Camponotus (Myrmoturba) maguassa* WM. M. WHEELER. See p. 237 (♀).

Type locality: Avakubi, BELGIAN CONGO (Lang and Chapin).

8. *Camponotus (Myrmoturba) natalensis* (F. SMITH) FOREL, 1914, Rev. Suisse Zool., XXII, p. 267. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 255.

Formica natalensis F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 33 (♀, ♀, ♂).

MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 418.

Camponotus natalensis ROGER, 1863, 'Verzeich. Formicid.,' p. 2. FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 79. MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 355. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 244. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 771. BINGHAM, 1903, Ann. Mag. Nat. Hist., (7) XII, p. 63 (♀). FOREL, in Schultz, 1910, 'Forschungsreise in Südafrika,' IV, p. 29 (♀).

Type locality: Port Natal, NATAL (R. W. Plant).

WEST AFRICA: (Pülleborn). TRANSVAAL: Pretoria (Distant).

8i. Var. *politiceps* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 38 (♀).

Type locality: Pietermaritzburg, NATAL (I. Trägårdh).

8a. Subsp. *corvus* FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 79 (♀, ♀, ♂). EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 51; 1896, Mem. Accad. Sc. Bologna, (5) V, p. 771.

Camponotus corvus DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 227.

Type locality: Valdezia, TRANSVAAL (P. Berthoud).

TRANSVAAL: Pretoria; Makapan (E. Simon).

8a. Var. *fulvipes* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 51 (♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 771.

Camponotus (Myrmoturba) natalensis var. *fulvipes* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 38 (♀, ♀, ♂).

CAMEROON: Lomie (Thesing); Mundame (Conrad). BELGIAN CONGO: Kasongo Forest (Grauer); Isangi (Lang and Chapin).

16. *Camponotus (Dinomyrmex) caffer* EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256.

Camponotus caffer EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 47 (♀), Pl. II, figs. 25, 26; 1896, Mem. Accad. Sc. Bologna, (5) V, p. 768. FOREL, 1904, Ann. Soc. Ent. Belgique, XLVIII, p. 177.

Camponotus maculatus subsp. *caffer* FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 69 (♀, ♀).

Camponotus (Myrmoturba) caffer FOREL, 1914, Rev. Suisse Zool., XXII, p. 267. Type locality: Makapan, TRANSVAAL (E. Simon).

CAPE PROVINCE: Port Elizabeth (H. Brauns); Cape of Good Hope.

17. *Camponotus (Dinomyrmex) immigrans* SANTSCHI, 1921, Ann. Soc. Ent. Belgique, LXI, p. 312 (♀).

Camponotus (Myrmotriz) immigrans SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 436 (♀, ♀, ♂).

Camponotus (? *Myrmotriz*) *immigrans* EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256.

Type locality: Molundu, CAMEROON (Reichensperger).

18. *Camponotus (Dinomyrmex) langi* Wm. M. WHEELER. See p. 241. (♀, ♀, ♂).

Type locality: Faradje, BELGIAN CONGO (Lang and Chapin).

BELGIAN CONGO: Garamba (Lang and Chapin).

19. *Camponotus (Dinomyrmex) longipes* (GERSTÄCKER) FOREL, 1913, Rev. Zool. Afr., II, p. 340 (♀); 1914, Rev. Suisse Zool., XXII, p. 268; 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 249 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 132 (♀, ♀ ?). EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256.

Formica longipes GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 262 (♀); in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 509 (♀), Pl. xxxii, fig. 7.

Camponotus longipes ROGER, 1863, 'Verzeich. Formicid.,' p. 2. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 400. GERSTÄCKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 354; 1873, in 'v. d. Decken's Reisen in Ost-Afrika, Gliederthiere,' p. 343 (♀). FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 164 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 240. EMERY, 1894, Mem. Accad. Sc. Bologna, (5) IV, p. 113 (♂). FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 70. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 768. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 37. FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 91 (♀, ♂); 1907, Mitt. Naturh. Mus. Hamburg, XXIV, p. 16 (♀); 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 266 (♀); 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 371 (♀). STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 391 (♀).

Type locality: Mossambique, PORTUGUESE EAST AFRICA (Peters).

BELGIAN CONGO: Sankisia (J. Bequaert); Boga, west of Mt. Ruwenzori (Schubotz). SOUTHERN RHODESIA: Mondu River (Zeally). PORTUGUESE EAST AFRICA: Inhambane (Fornasini). NYASALAND: Shore of Lake Nyasa (Heyne). GERMAN EAST AFRICA: Himo River, Mt. Kilimanjaro (C. Alluaud); Barikiwa (Schröder); Mafia Island (Voeltzkow); Darassalaam (A. Müller). ZANZIBAR: (Hildebrandt). BRITISH EAST AFRICA: Ndara (v. d. Decken); Moschi to the Uganda Railroad (Zimmer); Nairobi, Kikuyu, 1700 m.; Taveta, 750 m.; Voi, Taita; Mt. Kenia,

Meranga District, Fort Hall; Cheteni (Alluaud and Jeannel); Chake-Chake, Pemba Island; Fundu, W. Pemba (Voeltzkow). UGANDA: Buzubizi (C. Alluaud). WESTERN ABYSSINIA: (Ilg).

20. **Camponotus (Dinomyrmex) massinissa** Wm. M. WHEELER. See p. 244 (♀).

Type locality: Medje, BELGIAN CONGO (Lang and Chapin).

21. **Camponotus (Dinomyrmex) pompeius** FOREL. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256.

Camponotus rubripes subsp. *pompeius* FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 157 (♂).

Camponotus pompeius DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 248. EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 501, fig. (♂, ♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393 (♂, ♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 149 (♂, ♀).

Camponotus maculatus subsp. *pompeius* EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 769.

Camponotus (Myrmogigas) pompeius SANTSCHI, 1914, Deutsch. Ent. Zeitschr., p. 288 (♂).

Camponotus (Myrmoturba) pompeius FOREL, 1914, Rev. Suisse Zool., XXII, p. 267. STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 403 (♂).

Type locality: Gaboon, FRENCH CONGO (Büttner).

Togo: Bismarckburg (Conradt). CAMEROON: Duala (v. Rothkirch); Campo Mountains (Schultze); Mundane (Conradt). FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: Libenge; Duma (Schubotz).

21a. Subsp. *cassius* Wm. M. WHEELER. See p. 239 (♂).

Type locality: Yakuluku, BELGIAN CONGO (Lang and Chapin).

BELGIAN CONGO: Medje (Lang and Chapin).

21b. Subsp. *marius* EMERY, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 502, fig. (♂). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 403 (♂, ♀). See p. 239.

Camponotus maculatus subsp. *cognatus* MAYR, 1896, Ent. Tidskr., XVII, p. 249 (nec F. Smith).

Type locality: CAMEROON (Conradt; Sjöstedt).

CAMEROON: Molundu; Yukaduma (Schultze). FRENCH CONGO: Gaboon. BELGIAN CONGO: Duma (Schubotz); Kimuenza (Schultze); Medje; Akenge; Niapu (Lang and Chapin).

22. **Camponotus (Dinomyrmex) sexpunctatus** FOREL, 1914, Rev. Suisse Zool., XXII, p. 268. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 133 (♂, ♀). EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256.

Camponotus sexpunctatus FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 66 (♂, ♀).

Camponotus maculatus subsp. *sexpunctatus* EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 769. ALLUAUD and JEANNEL, 1914, Arch. Zool. Gén. Exp., LIII, p. 375.

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA (P. Berthoud).

BRITISH EAST AFRICA: Kibwezi, Wa-Kamba (C. Alluaud). ZANZIBAR: Haitajwa Cave (Alluaud and Jeannel).

23. *Camponotus (Dinomyrmex) varus* FOREL, 1914, Rev. Suisse Zool., XXII, p. 268. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256.

Camponotus varus FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 455 (♀).

Type locality: MOSSAMEDES (Baum and Van der Kellen).

24. *Camponotus (Dinomyrmex) wellmani* FOREL, 1913, Rev. Zool. Afr., II, p. 340 (♀); 1914, Rev. Suisse Zool., XXII, p. 268. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256.

Camponotus wellmani FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 67 (♀, ♀); 1911, Rev. Zool. Afr., I, p. 281 (♀). SANTSCHI, *ibid.*, p. 211 (♀). FOREL, 1912, Mitt. Naturh. Mus. Hamburg, XXIX, p. 180 (♀).

Camponotus (Dinomyrmex) wellmanni FOREL, 1916, Rev. Suisse Zool., XXIV, p. 441 (♀).

Type locality: BENGUELA (C. Wellman).

BENGUELA: CUCALA (J. Cruchet). BELGIAN CONGO: Beni (Murtula); Kapema; Kipaila (S. Neave); Bamayanga. GERMAN EAST AFRICA: Kifumbiro to Karagwe, between the great lakes.

24. Var. *rufipartis* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 441 (♀, ♀, ♂). See p. 245.

Type locality: BELGIAN CONGO (Kohl).

BELGIAN CONGO: Stanleyville; Faradje; Ngayu; Niangara (Lang and Chapin).

Subgenus 3. *Myrmosericus* FOREL

Camponotus subg. *Myrmosericus* FOREL, 1912, Mém. Soc. Ent. Belgique, XX, p. 91; 1914, Rev. Suisse Zool., XXII, p. 259. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 236.

Subgenotype: *Formica rufoglaucia* Jerdon, 1851.

25. *Camponotus (Myrmosericus) angusticeps* EMERY. FOREL, 1914, Rev. Suisse Zool., XXII, p. 268. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256.

Camponotus angusticeps EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 355 (♀), Pl. XVII, fig. 1. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 221. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 770.

Type locality: Cape of Good Hope (L. Péringuey).

26. *Camponotus (Myrmosericus) druryi* FOREL, 1914, Rev. Suisse Zool., XXII, p. 268. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256.

Camponotus druryi FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 160 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 229. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 769. H. STADELMANN, 1898, Deutsch-Ost-Afrika, IV, Hym., p. 37.

Type locality: ZANZIBAR (Hildebrandt).

27. *Camponotus (Myrmosericus) eugeniae* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 146 (♀); 1914, Rev. Suisse Zool., XXII, p. 268. SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 41 (♀). TRÄGÅRDH, *ibid.*, p. 46. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256.

Camponotus micans subsp. *eugeniae* FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 78 (♀).

Camponotus eugeniae FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 174 (♀, ♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 230. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 50; 1896, Mem. Accad. Sc. Bologna, (5) V, p. 770.

Camponotus rufoglaucus var. *eugeniae* MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 195. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 36.

Type locality: Valdezia, TRANSVAAL (P. Berthoud).

SOUTHERN RHODESIA: Springvale (G. Arnold). TRANSVAAL: Makapan (E. Simon). CAPE PROVINCE: Kimberley (E. Simon). BASUTOLAND: Lessouto (P. Berthoud). ZULULAND: Junction of the Umfolosi Rivers (I. Trägårdh). GERMAN EAST AFRICA: Kihengo (Stuhlmann).

27a. Subsp. *amplior* FOREL, 1913, Rev. Zool. Afr., II, p. 341 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 355 (♀).

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

BELGIAN CONGO: Old Kasongo (J. Bequaert).

28. *Camponotus (Myrmosericus) petersi* EMERY. FOREL, 1914, Rev. Suisse Zool., XXII, p. 268; 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 254 (♀). EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256.

Camponotus petersi EMERY, in litt., FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 70 (without description). EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 50 (♀), Pl. II, fig. 28; 1896, Mem. Accad. Sc. Bologna, (5) V, p. 770. FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 30 ((♀)); 1910, Ann. Soc. Ent. Belgique, LIV, p. 451.

Type locality: Vrijburg, CAPE PROVINCE (E. Simon).

RHODESIA: Bulawayo (G. Arnold). TRANSVAAL: Pretoria (E. Simon); Lydenburg (Wilms). NATAL: (Haviland). PORTUGUESE EAST AFRICA: Delagoa (Liengme).

28i. Var. *janus* FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 288 (♀).

Type locality: Bothaville, ORANGE FREE STATE (H. Brauns).

29. *Camponotus (Myrmosericus) rufoglaucus* (JERDON) FOREL, 1912, Mém. Soc. Ent. Belgique, XX, p. 91; 1914, Rev. Suisse Zool., XXII, p. 268. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256.

Formica rufoglauca JERDON, 1851, Madras Journ. Litt. Sc., XVII, p. 124; 1854, Ann. Mag. Nat. Hist., (2) XIII, p. 107 (♀).

Camponotus rufoglaucus EMERY, 1895, Ann. Soc. Ent. France, LXIV, pp. 48, footnote and 50, footnote, fig. (♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 770.

Type locality: Carnatic, INDIA (Jerdon).

29a. Subsp. *cinctellus* (GERSTÄCKER) FOREL, 1897, Abhandl. Senckenberg. Naturf. Ges., XXI, p. 188 (♀); 1901, Mitth. Schweiz. Ent. Ges., X, p. 311; in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 564; in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 89; 1909, Ann. Soc. Ent. Belgique, LIII, p. 67 (♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 25 (♀). KARAWALEW, 1911, Rev. Russe Ent., XI, p. 12 (♀). FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 371 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 356 (♀). SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 381 (♀, ♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 133 (♀); 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 41. TRÄGÅRDH, ibid., p. 46. See p. 246.

Formica cinctella GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 262 (♀, ♀, ♂); in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 510 (♀, ♀, ♂). MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 413.

Camponotus cinctellus ROGER, 1863, 'Verzeich. Formicid.,' p. 2. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 224.

Camponotus venustus FRAUENFELD, 1867, Verh. Zool. Bot. Ges. Wien, XVII, p. 441 (♀). MAYR, 1878, *ibid.*, XXVIII, p. 662, footnote (♀).

Camponotus rufoglaucus var. *cinctellus* MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 195. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 48, footnote (♀); 1895, Ann. Mus. Civ. Genova, XXXV, p. 180; 1896, *ibid.*, XXXVII, p. 158 (♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 770. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 36.

Type locality: Mozambique, PORTUGUESE EAST AFRICA (Peters).

BELGIAN CONGO: Zambi (J. Bequaert); Congo da Lemba; Mondombe; Mandungu; Eala (R. Mayné); Leopoldville (J. Maes). BENGUELA: (C. Wellman). MOSSAMEDES: between the Cubango and Cuito Rivers (Baum and Van der Kellen). TRANSVAAL: Pretoria (F. Silvestri). NATAL: (v. Schuckardt); Stamford Hill (I. Trägårdh). PORTUGUESE EAST AFRICA: Delagoa (Liengme); Tete (Peters); Lobombo Borges (Wilms). East Africa: common (Voeltzkow). GERMAN EAST AFRICA: Sachsenwald near Daressalaam (H. Prell); Uluguru Mts.; Amani (Zimmer); Tanga (Zimmer; C. Alluaud). ZANZIBAR: (Stuhlmann; Voeltzkow); Mwera River (C. Alluaud). BRITISH EAST AFRICA: Gazi; Likoni; Tiwi; Mombasa; Buru, Wa-Taita; Fort Hall, Kikuyu, 1330 m. (Alluaud and Jeannel). SOMALILAND: Matagoi to Lugh (V. Bottego). ABYSSINIA: Boran Galla, Middle Canale (V. Bottego). ANGLO-EGYPTIAN SUDAN: Khartum (Kafawaiew).

29a. Var. *intuens* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 67 (♀); 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 271 (♀); 1913, Rev. Zool. Afr., II, p. 340 (♀).

Type locality: BENGUELA (C. Wellman).

BELGIAN CONGO: Kwesi to Kilo (Bayer). ERITREA: Nefasit (K. Escherich).

29a. Var. *rufigenis* FOREL, 1913, Rev. Zool. Afr., II, p. 341 (♀). BEQUAERT, *ibid.*, p. 430. See p. 246.

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

BELGIAN CONGO: Stanleyville; Medje; Foko; Akenge; Niangara; Garanba; Faradje (Lang and Chapin).

29a. Var. *ustithorax* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 271 (♀); 1911, Bull. Soc. Vaudoise Sc. Nat., (5) LXVII, p. 371 (♀). STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 392 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 356 (♀); 1913, Rev. Zool. Afr., II, p. 340 (♀); 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 350 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 133 (♀).

Type locality: Ghinda, ERITREA (K. Escherich).

BELGIAN CONGO: Beni (Borgerhoff); St. Gabriel (Kohl); Yambata; Mandungu; Mondombe (R. Mayné); Karemi (Bayer); Kwidjwi Island, Lake Kivu (Schubotz). GERMAN EAST AFRICA: Buiko; Monga (H. Prell); Amani; Ulenge (Zimmer). BRITISH EAST AFRICA: Kavirondo Bay, Victoria Nyanza, 1112 m. (Alluaud and Jeannel). UGANDA: Western Unyoro, between Hoima and Butiaba; Ibanda, Mt. Ruwenzori, 1400 m. (C. Alluaud); Chacansengula to Kasengui (Bayer).

29b. Subsp. *controversus* SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 509.

Camponotus rufoglaucus subsp. *flavopilosus* VIEHMAYER, 1914, Arch. f. Naturg., LXXIX, Abt. A, Heft 12, (1913), p. 47 (♀) (nec *C. fulvopilosus* var. *flavopilosus* Emery, 1895).

Type locality: Manow, Langenburg, GERMAN EAST AFRICA.

BELGIAN CONGO: Lungube, Katanga (Gérard).

29c. Subsp. *cosmicus* (F. SMITH) EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 49; 1896, Mem. Accad. Sc. Bologna, (5) V, p. 770. MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.', II, 8, p. 20. STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11', I, p. 401 (♀).

Formica cosmica F. SMITH, 1858, 'Cat. Hym. Brit. Mus.', VI, p. 34 (♀). MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 413.

Camponotus cosmicus ROGER, 1863, 'Verzeich. Formicid.', p. 2. MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 355. DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 227. G. K. MARSHALL, 1902, Trans. Ent. Soc. London, pp. 535 and 569 (♀), Pl. XIX, figs. 55 and 56. BINGHAM, 1903, Ann. Mag. Nat. Hist., (7) XII, p. 63 (♀). DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, pp. 331 and 375.

Type locality: Port Natal, NATAL (R. W. Plant).

TRANSVAAL: Waterval Onder (Ross); Pretoria; Hamman's Kraal (E. Simon). CAPE PROVINCE: Hebron; De Aar (E. Simon); East London (Dixey and Longstaff). NATAL: Estcourt (Dixey and Longstaff). GERMAN EAST AFRICA: Ngare na Nyuki, Meru (Sjöstedt). BELGIAN CONGO: Duma (Schubotz). RHODESIA: Salisbury (Marshall).

29d. Subsp. *flavomarginatus* (MAYR) EMERY, 1895, Ann. Soc. Ent. France, LXIV, pp. 48, footnote (♀) and 50, footnote, fig.; 1896, Mem. Accad. Sc. Bologna, (5) V, p. 770; 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 604; 1899, *ibid.*, XXXIX, p. 501. FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 144 (♀, ♀, ♂). REICHENSPERGER, 1913, Zool. Jahrb. Abt. Syst., XXXV, p. 215. See p. 247.

Camponotus flavomarginatus MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 664 (♀); 1863, *ibid.*, XIII, p. 399. ROGER, 1863, 'Verzeich. Formicid.', p. 2. EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 365 (♀).

Camponotus micans var. *flavomarginatus* FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 174 (♀, ♀).

Camponotus micans subsp. *albisectus* EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 120 (♀).

Camponotus albisectus DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 221.

Camponotus rufoglaucus subsp. *cinctellus* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 70.

Camponotus rufoglaucus subsp. *cinctellus* var. *flavomarginatus* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 271 (♀).

Camponotus (*Myrmosericus*) *albisectus* FOREL, 1914, Rev. Suisse Zool., XXII, p. 268.

Camponotus (*Myrmosericus*) *rufoglaucus* subsp. *cinctellus* var. *flavomarginatus* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afđ., III, p. 41 (♀). FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 254 (♀).

Type locality: Akwapim Mountains, GOLD COAST.

ABYSSINIA: Harar; Tehercher; confluent Akaki; Karssa; Mt. Zyoual; Kounhi; Barko; Bourka (de Rothschild); Bass Narok (V. Bottego); western Abyssinia (Ilg). ERITREA: Keren (Beccari). SOMALILAND: Ganale (Ruspoli); Mogadiscio (Bricchetti-Robecchi). NATAL: Stamford Hill (I. Trägårdh); Durban (C. B. Cooper). PORTUGUESE EAST AFRICA: Delagoa Bay (Liengme). TRANSVAAL:

Valdezia (P. Berthoud). ANGOLA: (M. Buchner). BELGIAN CONGO: Thysville (J. Bequaert); Akenge; Vankerekhovenville; Garamba; Faradje (Lang and Chapin).

29d. Var. *contrarius* SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 509 (♀).

Type locality: Victoria Falls, RHODESIA (G. Arnold).

29d. Var. *paucipubens* SANTSCHI, 1911, Rev. Zool. Afr., I, p. 212 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 381 (♀).

Camponotus rufoglaucus subsp. *cinctellus* var. *paucipubens* FOREL, 1913, Rev. Zool. Afr., II, p. 341 (♀). BEQUAERT, *ibid.*, p. 430. STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 401 (♀).

Type locality: Cucala, BENGUELA (J. Cruchet).

ANGOLA: Quifangondo (F. Silvestri). BELGIAN CONGO: Welgelegen (J. Bequaert). FRENCH CONGO: Fort Crampel (Schubotz).

29e. Subsp. *syphax* Wm. M. WHEELER. See p. 246 (♀).

Type locality: Zambi, BELGIAN CONGO (Lang, Chapin, and J. Bequaert).

BELGIAN CONGO: Boma (Lang, Chapin, and J. Bequaert).

29f. Subsp. *vestitus* (F. SMITH) EMERY, 1894, Mem. Accad. Sc. Bologna, (5) IV, p. 113 (♀). FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 70. EMERY, 1895, Ann. Soc. Ent. France, LXIV, pp. 48, 49, footnote (♀), and 50, footnote, fig.; 1896, Mem. Accad. Sc. Bologna, (5) V, p. 770. FOREL, 1907, Mitt. Naturh. Mus. Hamburg, XXIV, p. 16 (♀). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 41 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 381 (♀).

Formica vestita F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 32 (♀). MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 422.

Camponotus rufoglaucus subsp. *cinctellus* var. *vestita* FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 89; in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 25 (♀, ♀); 1910, Ann. Soc. Ent. Belgique, LIV, p. 457 (♀, ♀); 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 272 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 355 (♀); 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 254 (♀).

Type locality: Port Natal, NATAL.

CAPE PROVINCE: Cape Town (E. Simon; Bergius; Wilms); Mafeking (G. Arnold). TRANSVAAL: Pretoria (F. Silvestri); Lydenburg (Wilms). NATAL: (Haviland). ZULULAND: Junction of the Umfolosi Rivers (I. Trägårdh). PORTUGUESE EAST AFRICA: Delagoa (Liengine); Mozambique (Fornasini). GERMAN SOUTHWEST AFRICA: Lübbert; Okahandja (Peters; Casper); Grootfontein (v. Erffer). BELGIAN CONGO: Elisabethville (Leprieux); Karemi (Bayer); valley of the Lubumbashi (Buttgenbach). WESTERN ABYSSINIA: (Ilg). East Africa: (Voeltzkow).

29f. Var. *pectitus* SANTSCHI, 1911, Rev. Zool. Afr., I, p. 212 (♀, ♀).

Camponotus (Myrmosericus) rufoglaucus subsp. *cinctellus* var. *pectita* FOREL, 1913, Deutsch. Ent. Zeitschr., Beih., p. 224 (♀).

Type locality: Cucala, BENGUELA (J. Cruchet).

FRENCH CONGO: Moyen Chari (Decorse). SOUTHERN RHODESIA: Bembesi (G. Arnold).

29g. Subsp. *zanzibaricus* FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 287 (♀).

Type locality: ZANZIBAR.

29h. Subsp. *zimmermanni* (FOREL) EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 49, footnote; 1896, Mem. Accad. Sc. Bologna, (5) V, p. 770; 1896, Ann. Mus. Civ. Genova, XXXVII, p. 158 (♀); 1897, *ibid.*, XXXVIII, p. 604 (♀).

Camponotus zimmermanni FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 66 (♀); 1904, Ann. Mus. Zool. Ac. Sc. St. Pétersbourg, VIII, (1903), p. 381 (♀).

Camponotus (Myrmosericus) zimmermanni EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 256.

Type locality: SOUTHERN ABYSSINIA (Ilg).

BRITISH EAST AFRICA: Maddo Wells (V. Bottego). SOMALILAND: Salmoreto (Ruspoli). ABYSSINIA: Harar (Dmitriev); Webi (Ruspoli).

29i. Subsp. *zulu* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 49, footnote, and p. 50 (♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 770; 1895, Ann. Mus. Civ. Genova, XXXV, p. 180. SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 41 (♀).

Type locality: Isipingo, NATAL (Weitzacker).

NATAL: Verulam (Weitzacker); Amanzimtoti (Trägårdh). ABYSSINIA: Arussi Galla, Ganale Gudda (V. Bottego).

30. *Camponotus (Myrmosericus) valdezianus* FOREL, 1914, Rev. Suisse Zool., XXII, p. 268.

Camponotus valdezianus FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 90 (♀, ♀). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 256. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 770. MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.', II, 8, p. 20.

Type locality: Valdezia, TRANSVAAL (P. Berthoud).

GERMAN EAST AFRICA: Kilimanjaro (Sjöstedt).

Subgenus 4. *Orthonotomyrmex* ASHMEAD

Orthonotomyrmex ASHMEAD, 1906, Proc. Ent. Soc. Washington, VIII, p. 31.

Orthonotus ASHMEAD, 1905, Canadian Ent., XXXVII, p. 384 (nec Westwood, 1829).

Camponotus subg. *Myrmentoma* FOREL, 1912, Mém. Soc. Ent. Belgique, XX, p. 92.

Camponotus subg. *Orthonotomyrmex* FOREL, 1914, Rev. Suisse Zool., XXII, p. 264. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 244.

Subgenotype: *Formica sericea* Fabricius, 1798 (Ashmead, 1905); *Formica lateralis* Olivier, 1791 (Forel, 1914).

31. *Camponotus (Orthonotomyrmex) acutisquamis* MAYR. FOREL, 1914, Rev. Suisse Zool., XXII, p. 273. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 257.

Camponotus acutisquamis MAYR, 1902, Verh. Zool. Bot. Ges. Wien, LII, p. 296 (♀).

Camponotus chrysurus subsp. *acutisquamis* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 269; 1913, Rev. Suisse Zool., XXI, p. 670 (♀, ♀).

Camponotus (Orthonotomyrmex) chrysurus subsp. *acutisquamis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 357 (♀); 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 256 (♀); 1916, Rev. Suisse Zool., XXIV, p. 444 (♀, ♂).

Type locality: Bonjongo, CAMEROON (R. Buchholz).

CAMEROON: (Conradt). BELGIAN CONGO: (Kohl); Mayombe (de Briey).

32. *Camponotus (Orthonotomyrmex) arminius* (FOREL) EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 257.

Camponotus barbarossa subsp. *arminius* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 457 (♂).¹

Type locality: Delagoa, PORTUGUESE EAST AFRICA (Liengme).

32i. Var. *bicontractus* (FOREL).

Camponotus barbarossa subsp. *arminius* var. *bicontractus* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 256 (♂).

Type locality: Durban, NATAL (G. Arnold).

33. *Camponotus (Orthonotomyrmex) braunsi* MAYR. FOREL, 1914, Rev. Suisse Zool., XXII, p. 273; 1913, Ann. Soc. Ent. Belgique, LVII, p. 357 (♂). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 135 (♀). EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 257.²

Camponotus braunsi MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 151 (♂). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774. MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 28, footnote (♂, ♀).

Camponotus (Myrmentoma) braunsi FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 146 (♂).

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA (H. Brauns).

RHODESIA: (G. Arnold). BELGIAN CONGO: Congo da Lemba (R. Mayné).

GERMAN EAST AFRICA: New Moschi, Mt. Kilimanjaro, 800 m. (Alluaud and Jeannel); Waboniland.

33a. Subsp. *erythromelus* EMERY, 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 605; 1898, Rend. Accad. Sc. Bologna, N. S., II, p. 227; 1899, Ann. Mus. Civ. Genova, XXXIX, p. 501 (♂).

Camponotus erythromelus EMERY, 1896, *ibid.*, XXXVII, p. 158, fig. (♂).

Type locality: Lugh, SOMALILAND (V. Bottego).

SOMALILAND: Matagoi to Lugh (V. Bottego); Magala Re Umberto (Ruspoli).

ABYSSINIA: Ganale River; Ogaden (Ruspoli); Sancurar to Amarr Burgi (V. Bottego).

33a₁. Var. *transitorius* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 135 (♂).

Type locality: River Amboni, Mt. Kenia, BRITISH EAST AFRICA (C. Alluaud).

34. *Camponotus (Orthonotomyrmex) chrysurus* GERSTÄCKER. FOREL, 1913, Rev. Zool. Afr., II, p. 346 (♀); 1914, Rev. Suisse Zool., XXII, p. 273; 1914, Ann. Soc. Ent. Belgique, LVII, p. 357 (♂, ♀); 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 350 (♂, ♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 135 (♂, ♀). EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 257.³

Camponotus chrysurus GERSTÄCKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 355 (♂); 1873, in 'v. d. Decken's Reisen in Ost Afrika, Gliederthiere,' p. 345 (♂), Pl. XIV, fig. 9. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 224. EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 183 (♂); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 37. FOREL, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 308 (♂); 1909, *ibid.*, LIII, p. 67 (♂). BEQUAERT, 1913, Rev. Zool. Afr., II, p. 431.

Type locality: Ndaja, BRITISH EAST AFRICA (v. d. Decken).

¹*Camponotus barbarossa* "Emery, in litt." Forel, 1914, Rev. Suisse Zool., XXII, p. 273, has never been described. Santschi (1921) makes *C. arminius* the type of his subgenus *Myrmopelta*.

²Placed by Santschi (1921) in his subgenus *Myrmioleptis*.

³Placed by Santschi (1921) in his subgenus *Myrmopelta*.

BELGIAN CONGO: Elisabethville; Sankisia (J. Bequaert); St. Gabriel (Kohl); Congo da Lemba; Mondembe (R. Mayné); Mayombe (de Briey). BENGUELA: (C. Wellman). GERMAN EAST AFRICA: New Moschi, Mt. Kilimanjaro, 800 m. (Alluaud and Jeannel). ABYSSINIA: Arussi Galla, Ganale Gudda (V. Bottego). BRITISH EAST AFRICA: Nairobi, Kikuyu, 1700 m. (Alluaud and Jeannel); Tsavo (Bayer).

34. Var. *apellis* (FOREL) SANTSCHI, 1911, Rev. Zool. Afr., I, p. 213.

Camponotus chrysurus subsp. *acutisquamis* var. *apellis* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 269 (♂); 1911, Rev. Zool. Afr., I, p. 281 (♀). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 397 (♀).

Type locality: BENGUELA (C. Wellman).

BENGUELA: Cucala (J. Cruchet). BELGIAN CONGO: Banana (Etienne); Duma (Schubotz).

34a. Subsp. *kohlbrunneri* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 268 (♂).

Type locality: WESTERN ABYSSINIA (Ilg).

35. *Camponotus (Orthonotomyrmex) dofeini* FOREL. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 257.

Camponotus dofeini FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 291 (♀).

Camponotus (Myrmosphincta) dofeini FOREL, 1914, Rev. Suisse Zool., XXII, p. 273.

Type locality: Bothaville, ORANGE FREE STATE (H. Brauns).

36. *Camponotus (Orthonotomyrmex) epinotalis* SANTSCHI, 1916, Ann. Soc. Ent. France, LXXXIV, (1915), p. 508 (♀). EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 257.¹

Type locality: Victoria Falls, RHODESIA (G. Arnold).

37. *Camponotus (Orthonotomyrmex) erinaceus* GERSTAECKER. FOREL, 1914, Rev. Suisse Zool., XXII, p. 273. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 135 (♀). EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 257.²

Camponotus erinaceus GERSTAECKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 355 (♀); 1873, in 'v. d. Decken's Reisen in Ostafrika, Gliederthiere,' p. 345 (♀), Pl. XIV, fig. 10. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 229. EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 183 (♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 38. FOREL, 1899, Ann. Soc. Ent. Belgique, XLIII, p. 306 (♀); 1907, Ann. Mus. Nat. Hungarici, V, p. 31 (♀). MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 21 (♀, ♀). FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 371 (♀).

Type locality: Arusha, GERMAN EAST AFRICA (v. d. Decken).

GERMAN EAST AFRICA: Kilimanjaro (Bornemisza); Kibosho, Mt. Kilimanjaro (C. Alluaud); Kibonoto, Kilimanjaro, 1300-1900 m. (Sjöstedt); Moschi (H. Prell). BRITISH EAST AFRICA: Monga (H. Prell); Nairobi, 1700 m. (Alluaud and Jeannel). ABYSSINIA: Arussi Galla, Ganale Gudda (V. Bottego).

38. *Camponotus (Orthonotomyrmex) mayri* FOREL, 1914, Rev. Suisse Zool., XXII, p. 273. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus mayri FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 92 (♀), Pl. I, fig. 1; 1886, Ann. Soc. Ent. Belgique, XXX, p. 191 (♀). DALLA TORRE,

¹Santschi (1921) makes this the type of his subgenus *Myrmisolepis*.

²Placed by Santschi (1921) in his subgenus *Myrmisolepis*.

1893, 'Cat. Hym.,' VII, p. 242. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 70. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 55 (♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 37. FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 91 (♀). MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 21. SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393 (♀). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 451 (♀). STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exped. (1907-08),' III, p. 392 (♀).

Camponotus (Myrmentoma) mayri FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 146 (♀).

Type locality: Valdezia, TRANSVAAL (P. Berthoud).

FRENCH CONGO: Brazzaville (A. Weiss). PORTUGUESE CONGO: Chinchoxo. BELGIAN CONGO: Kasindi (Schubotz). SOUTHERN RHODESIA: Springvale (G. Arnold). TRANSVAAL: Makapan; Pretoria; Hamman's Kraal (E. Simon). NATAL: Durban. PORTUGUESE EAST AFRICA: Delagoa (Liengme). GERMAN EAST AFRICA: Kibonoto, Kilimanjaro (Sjöstedt). ZANZIBAR: (Hildebrandt). BRITISH EAST AFRICA: Chake-Chake, Pemba Island (Voeltzkow); Mombasa (Hildebrandt).

38i. Var. *cubangensis* FOREL, 1895, Mitth. Schweiz. Ent. Ges., X, p. 309 (♀, ♀).

Camponotus mayri var. *kubangensis* FOREL, in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 563 (♀).

Type locality: between the Cubango and Cuito Rivers, MOSSAMEDES (Baum and Van der Kellen).

38a. Subsp. *ledieui* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 444 (♀).

Type locality: Shilouvane, TRANSVAAL (Junod).

38b. Subsp. *sankisianus* FOREL, 1913, Rev. Zool. Afr., II, p. 347 (♀). SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 313.

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

FRENCH CONGO: Mandouga (A. Weiss).

39. *Camponotus (Orthonotomyrmex) scalaris* FOREL, EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.¹

Camponotus scalaris FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 308 (♀).

Camponotus (Myrmosphincta) scalaris FOREL, 1914, Rev. Suisse Zool., XXII, p. 273.

Type locality: NATAL, 1800 in. (Haviland).

40. *Camponotus (Orthonotomyrmex) sericeus* (FABRICIUS) FOREL, 1914, Rev. Suisse Zool., XXII, p. 273. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 134 (♀, ♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 381 (♀). EMERY, 1915, *ibid.*, X, p. 4 (♀). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 397 (♀, ♂). EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258. See pp. 249 and 1050.

Formica sericea FABRICIUS, 1798, 'Suppl. Ent. Syst.,' p. 279 (♀). LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 117 (♀), Pl. III, fig. 17. COQUEBERT, 1804, 'Illustr. Iconogr. Ins.,' III, p. 99 (♀), Pl. XXII, fig. 12. LEPELETIER, 1836, 'Hist. Nat. Ins. Hym.,' I, p. 217 (♀). F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 31.

¹Placed by Santschi (1921) in his subgenus *Myrmisolepis*.

Formica aurulenta LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 114 (♀), Pl. III, fig. 9. F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 31. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 412.

Lasius sericeus FABRICIUS, 1804, 'Syst. Piez.,' p. 416 (♀).

Formica cinerascens JERDON, 1851, Madras Journ. Litt. Sc., XVIII, p. 123 (♀, ♂) (nec Fabricius); 1854, Ann. Mag. Nat. Hist., (2) XIII, p. 106 (♀, ♂).

Formica obtusa F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 30 (♀).

Camponotus sericeus MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 675 (♀), Pl. XIX, fig. 4; 1863, *ibid.*, XIII, p. 401. ROGER, 1863, 'Verzeich. Formicid.,' pp. 2 and 44 (♀). MAYR, 1866, Verh. Zool. Bot. Ges. Wien, XVI, p. 886 (♀). EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 364 (♀, ♀). FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 94 (♀, ♀). EMERY, 1881, Ann. Mus. Civ. Genova, XVI, p. 526. ERN. ANDRÉ, 1882, 'Spec. Hym. Europ. Algérie,' II, p. 149 (♀). MAGRETTI, 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 244 (♀); 1884, Ann. Mus. Civ. Genova, XXI, p. 537 (♀, ♀). FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 192 (♀, ♀, ♂); 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 19 (♀). EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 120; 1892, Ann. Soc. Ent. France, LX, (1891), p. 566 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 251. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 70. EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 184; 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774; 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 605; 1899, *ibid.*, XXXIX, p. 501. G. K. MARSHALL, 1902, Trans. Ent. Soc. London, pp. 535 and 569 (♀), Pl. XIX, fig. 54. MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.,' p. 11. BINGHAM, 1903, Ann. Mag. Nat. Hist., (7) XII, p. 63 (♀). MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 21; 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. 1, p. 389. FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 91 (♀). F. F. KOHL, 1907, Denkschr. Ak. Wiss. Wien, LXXI, p. 283 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 149 (♀). FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 268 (♀). LONGSTAFF, 1911, Ent. Monthly Mag., XLVII, p. 124 (♀). KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 11 (♀). LAMBORN, 1914, Trans. Ent. Soc. London, (1913), Proc., p. cxxiv.

Camponotus (Myrmecotoma) sericeus FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 146 (♀).

Type locality: Senegal (Bosc).

INDIA, CEYLON, ARABIA, EGYPT. SENEGAMBIA: Dakar (C. Alluaud; F. Silvestri); Cape Verde. TOGO: Misahöhe (Smend). SOUTHERN NIGERIA: Moor Plantation near Ibadan (Lamborn). FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: Medje; Poko; Faradje (Lang and Chapin); Duma (Schubotz); Kabare (J. Bequaert). ANGLO-EGYPTIAN SUDAN: Khartum (I. Trägårdh; Karawaiew); Kerrer (Longstaff); Suakin; Kor Langhebb; Kassala (Magretti). ERITREA: Keren; between Massaua and Ain (Beccari); Asmara; Ghinda (F. Silvestri); Assab (Doria and Beccari); Kor Lebka (Magretti). SOUTHERN ARABIA: Makalla (O. Simony). ABYSSINIA: (Ilg); Arussi Galla, Ganale Gudda; Dimé to Bass Narok; Bass Narok (V. Bottego); Coromma; Etoke; Bela (Ruspoli). SOMALILAND: (C. Keller); Obbia (Bricchetti-Robecchi). UGANDA: Gondokoro (F. Werner); Western Nyoro, between Hoima and Butiaba (C. Alluaud). EAST AFRICA: common (Voeltzkow). BRITISH EAST AFRICA: Mt. Kenia, Meranga District, Fort Hall; Kisumu, Victoria Nyanza; Tehania River, 1520 m.; Likoni (Alluaud and Jeannel).

GERMAN EAST AFRICA: Kiboroto, Mt. Kilimanjaro, 1000-1200 m. (Sjöstedt); Himo River, Mt. Kilimanjaro (C. Alluaud). SOUTHERN RHODESIA: Salisbury (Marshall); Khami River (G. Arnold).

40. Var. *sulgeri* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 313 (♀).

Camponotus (Orthonotomyrmex) sericeus subsp. *sulgeri* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 397 (♂, ♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

BELGIAN CONGO: Duma (Schubotz).

41. *Camponotus (Orthonotomyrmex) vividus* (F. SMITH) SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 264. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258. See p. 247.¹

Formica vivida F. SMITH, 1853, 'Cat. Hym. Brit. Mus.,' VI, p. 31 (♀). MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 422.

Formica laboriosa F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 32 (♀). MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 416.

Camponotus vividus ROGER, 1863, 'Verzeich. Formicid.,' p. 2. MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 354. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 257. EMERY, 1898, Rend. Accad. Sc. Bologna, N. S., II, p. 227.

Camponotus laboriosus ROGER, 1863, 'Verzeich. Formicid.,' p. 2. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 237. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 771.

Colobopsis vivida MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 354.

Camponotus meinerti FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 189 (♂, ♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 242. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 70. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 1 (♂, ♀). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 151. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 775. FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 74; 1907, *ibid.*, XXIV, p. 16 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXXVIII, (1909), p. 393 (♂, ♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 149 (♂, ♀). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 451 (♂); 1911, Rev. Zool. Afr., I, p. 282 (♂).

Camponotus (Colobopsis) vividus EMERY, 1889, Ann. Mus. Civ. Genova, XXVII, p. 517.

Camponotus (Orthonotomyrmex) meinerti SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 381 (♂). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 357 (♂); 1913, Rev. Zool. Afr., II, p. 346 (♂, ♂); 1914, Rev. Suisse Zool., XXII, p. 273; 1916, *ibid.*, XXIV, p. 444 (♂, ♀). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 396 (♂, ♀).

Type locality: SIERRA LEONE (D. F. Morgan).

SENEGAMBIA: Dakar (F. Silvestri). FRENCH GUINEA: Los Islands (H. Brauns); Konakry (F. Silvestri). SIERRA LEONE: (H. Brauns); Samlia Falls, River N'Gamie (Mocquerys). LIBERIA: (Kieselbach); Monrovia (Duke). GOLD COAST: Akra; Addah. TOGO: Bismarckburg (Conradt). SOUTHERN NIGERIA: Olokemeji (F. Silvestri). CAMEROON: Mundame (Conradt); Bibundi (Tessmann). FERNANDO PO: (Conradt). SPANISH GUINEA: Alen (Tessmann). FRENCH CONGO: Loango (H. Brauns); Gaboon (Büttner); Ogowe (Mocquerys); Brazzaville; Mandouga (A.

¹Placed by Santschi (1921) in his subgenus *Myrmopelta*.

Weiss); Fort Archambault; Fort de Possel to Fort Crampel; Mongumba (Schubotz). BELGIAN CONGO: Boma (H. Brauns); Kondué (Luja); Leopoldville (Dubois); Eala (R. Mayné); Mayombe (de Briey); Duma (Schubotz); Lukolela; Malela (Lang and Chapin). ANGOLA: Malange (M. Buchner). PORTUGUESE EAST AFRICA: Delagoa (Liengme); Ibo.

41. Var. *semidepilis* Wm. M. WHEELER. See p. 248 (♀).

Type locality: Medje, BELGIAN CONGO (Lang and Chapin).

BELGIAN CONGO: Leopoldville (Lang and Chapin).

41a. Subsp. *cato* (FOREL). See p. 248.

Camponotus (*Orthonotomyrmex*) *meinerti* subsp. *cato* FOREL, 1913, *Rev. Zool. Afr.*, II, p. 346 (♀).

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

BELGIAN CONGO: Stanleyville; Garamba; Medje; Avakubi; Akenge; Bengamisa; Niangara; Thysville (Lang and Chapin).

41b. Subsp. *reginæ* (FOREL).

Camponotus reginæ FOREL, 1901, *Mith. Schweiz. Ent. Ges.*, X, p. 307 (♀); in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 562 (♀); 1909, *Ann. Soc. Ent. Belgique*, LIII, p. 66 (♀).

Camponotus meinerti var. *reginæ* FOREL, 1911, *Rev. Zool. Afr.*, I, p. 282 (♀); 1911, *Sitzb. Bayer. Akad. Wiss.*, p. 291 (♀).

Camponotus (*Orthonotomyrmex*) *meinerti* subsp. *reginæ* FOREL, 1913, *Rev. Suisse Zool.*, XXI, p. 670 (♀); 1913, *Rev. Zool. Afr.*, II, p. 346 (♀).

Camponotus meinerti subsp. *reginæ* BEQUAERT, *ibid.*, p. 431.

Camponotus (*Myrmotoma*) *meinerti* subsp. *reginæ* FOREL, 1913, *Ann. Soc. Ent. Belgique*, LVII, p. 146 (♀).

Type locality: between the Cubango and Cuito Rivers, MOSSAMEDES (Baum and Van der Kellen).

LIBERIA. CAMEROON: (Conradt); Victoria (Fickendey).^{*} BELGIAN CONGO: Shinsenda (J. Bequaert); Kondué (Luja). BENGUELA: (C. Wellman). SOUTHERN RHODESIA: Kandahar Island, Zambesi River (G. Arnold).

Subgenus 5. *Myrmotrema* FOREL

Camponotus subg. *Myrmotrema* FOREL, 1912, *Mém. Soc. Ent. Belgique*, XX, p. 91; 1914, *Rev. Suisse Zool.*, XXII, p. 262. EMERY, 1920, *Rev. Zool. Afr.*, VIII, 2, p. 245.

Subgenotype: *Camponotus foraminosus* Forel, 1879.

42. *Camponotus* (*Myrmotrema*) *aurofasciatus* SANTSCHI, 1915, *Ann. Soc. Ent. France*, LXXXIV, pp. 267 and 276 (♀); 1920, 'Études Maladies Parasites Cacaoyer S. Thomé,' p. 4 (♀).

Type locality: SAN THOMÉ.

43. *Camponotus* (*Myrmotrema*) *bayeri* FOREL, 1914, *Rev. Suisse Zool.*, XXII, p. 270. SANTSCHI, 1915, *Ann. Soc. Ent. France*, LXXXIV, p. 271. EMERY, 1920, *Rev. Zool. Afr.*, VIII, 2, p. 258. See p. 252.

Camponotus bayeri FOREL, 1913, *Ann. Soc. Ent. Belgique*, LVII, p. 356 (♀).

Type locality: Karemi, BELGIAN CONGO (Bayer).

BELGIAN CONGO: Paradje (Lang and Chapin).

44. *Camponotus* (*Myrmotrema*) *bituberculatus* ERN. ANDRÉ. FOREL, 1914, *Rev. Suisse Zool.*, XXII, p. 270. SANTSCHI, 1914, *Boll. Lab. Zool. Gen. Agrar.*

Portici, VIII, p. 381 (♂); 1915, Ann. Soc. Ent. France, LXXXIV, p. 271. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus bituberculatus ERN. ANDRÉ, 1889, Rev. d'Ent. Caen, VIII, p. 271 (♂); 1890, *ibid.*, IX, p. 311 (♀). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 223. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 775.

Type locality: Dakar, SENEGAMBIA (Mocquersys).

SIERRA LEONE: (Mocquersys). FRENCH GUINEA: Kakoulima (F. Silvestri).

45. *Camponotus (Myrmotrema) bottegoi* EMERY. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 136 (♂, ♀); 1915, Ann. Soc. Ent. France, LXXXIV, p. 270. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus bottegoi EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 181 (♂, ♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 775.

Camponotus (Myrmotremba) bottegoi FOREL, 1914, Rev. Suisse Zool., XXII, p. 267.

Type locality: Lower Ganana, SOMALILAND (V. Bottego).

ABYSSINIA: Boran Galla, Middle Ganale (V. Bottego). SOMALILAND: Obbia (Bricchetti-Robecchi). BRITISH EAST AFRICA: Taveta; Voi (Alluaud and Jeannel). GERMAN EAST AFRICA: New Moschi, Mt. Kilimanjaro, 800 m. (Alluaud and Jeannel). ZANZIBAR: Mvera River (C. Alluaud).

46. *Camponotus (Myrmotrema) carbo* EMERY. FOREL, 1914, Rev. Suisse Zool., XXII, p. 270. SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 271. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus carbo EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 364, fig. (♂). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 223. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 775; 1898, Rend. Accad. Sc. Bologna, N. S., II, p. 227.

Type locality: Sciotele, ERITREA (Beccari).

46i. Var. *occidentalis* MAYR, 1902, Verh. Zool. Bot. Ges. Wien, LII, p. 300 (♂, ♀). SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 271.

Type locality: Akta, GOLD COAST (R. Buchholz).

46a. Subsp. *honorus* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 268 (♂). *Camponotus (Myrmotrema) foraminosus* subsp. *honorus* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 268.

Type locality: WESTERN ABYSSINIA (Ilg).

46b. Subsp. *osiris* FOREL, 1911, Rev. Zool. Afr., I, p. 281 (♂).

Camponotus (Myrmotrema) olivieri subsp. *osiris* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 270.

Type locality: Congo da Lemba, BELGIAN CONGO (R. Mayné).

46c. Subsp. *puberulus* EMERY, 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 604 (♂); 1898, Rend. Accad. Sc. Bologna, N. S., II, p. 227. FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 268 (♂); 1911, Rev. Zool. Afr., I, p. 280; 1913, *ibid.*, II, p. 341 (♂). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 136 (♂).

Camponotus carbo var. *puberula* EMERY, 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 63.

Camponotus (Myrmotrema) puberulus SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 271 and 278 (♂).

Type locality: Coromma, ABYSSINIA (Ruspoli).

BELGIAN CONGO: Congo da Lemba (R. Mayné); Kwesi to Kilo (Bayer). UGANDA: Unyoro Province, near Lake Albert (C. Alluaud). WESTERN ABYSSINIA: (Ilg). ERITREA: Keren, between Saganeiti and Adi Ugri (Mancini and Ruggeri).

47. **Camponotus (Myrmotrema) compressiscapus** ERN. ANDRÉ. SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 271. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus compressiscapus ERN. ANDRÉ, 1889, Rev. d'Ent. Caen, VIII, p. 218 (♂); 1890, *ibid.*, IX, p. 312 (♂). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 225. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 775.

Camponotus (Myrmosericus) compressiscapus FOREL, 1914, Rev. Suisse Zool., XXII, p. 268.

Type locality: SIERRA LEONE (Mocquerys).

48. **Camponotus (Myrmotrema) confluens** FOREL. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus (Myrmamblys) confluens FOREL, 1913, Rev. Zool. Afr., II, p. 342 (♂); 1914, Rev. Suisse Zool., XXII, p. 271.

Camponotus confluens BEQUAERT, 1913, Rev. Zool. Afr., II, p. 431.

Type locality: Welgelegen, BELGIAN CONGO (J. Bequaert).

BELGIAN CONGO: Sankisia (J. Bequaert).

48s. Var. **bequaerti** FOREL, 1913, Rev. Zool. Afr., II, p. 343 (♂, ♀). BEQUAERT, *ibid.*, p. 431.

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

48s. Var. **trematogaster** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 266 (♂).

Type locality: Lake Nyanza, Katanga, BELGIAN CONGO (Gérard).

49. **Camponotus (Myrmotrema) crucheti** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 271.

Camponotus perrisii subsp. *crucheti* SANTSCHI, 1911, Rev. Zool. Afr., I, p. 215 (♂, ♀, ♂).

Type locality: Cucala, BENGUELA (J. Cruchet).

50. **Camponotus (Myrmotrema) diplopunctatus** EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 22, fig. 12 (♂, ♀). EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Type locality: Nefasit, ERITREA (F. Silvestri).

ERITREA: Mayabal (F. Silvestri).

51. **Camponotus (Myrmotrema) foraminosus** FOREL, 1914, Rev. Suisse Zool., XXII, p. 270; 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 350; 1916, Rev. Suisse Zool., XXIV, p. 443 (♂, ♀). SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 268 and 272. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258. See pp. 249 and 1050.

Camponotus niveosetosus EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 364 (♂, ♀) (*nec* Mayr). MAGRETTI, 1884, *ibid.*, XXI, p. 537; 1884, Bull. Soc. Ent. Italiana, XV, (1883), p. 244 (♂, ♀) (*nec* Mayr).

Camponotus foraminosus FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 87 (♂); 1886, Ann. Soc. Ent. Belgique, XXX, p. 174 (♂). EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 358. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 231. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774. SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393 (♂). H. KOHL, 1909, Natur u. Offenbarung, LV, p. 102. STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 149 (♂, ♀). FOREL, 1911, Sitzb. Bayer. Akad. Wiss., p. 291 (♂); 1913, Rev. Suisse Zool., XXI, p. 671 (♂).

Type locality: Cape Verde, SENEGAMBIA.

SOUTHERN NIGERIA: Old Calabar (Bates). CAMEROON: (Conradt). FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: Kimpoko (Büttner); St. Gabriel (Kohl); Faradje; Avakubi; Stanleyville; Bengamisa (Lang and Chapin). ANGLO-EGYPTIAN SUDAN: Sebderat (Magretti). ERITREA: Keren (Beccari); Kor Lebka (Magretti). CAPE PROVINCE: Cape of Good Hope (L. Péringuey).

51a. Subsp. *auropubens* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 67 (♂). EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 182 (♂); 1896, *ibid.*, XXXVII, p. 159 (♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774; 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 604 (♂, ♀). FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 90. SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Af., III, p. 41 (♂). See p. 250.

Camponotus grandidieri subsp. *auropubens* MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 150 (♂).

Camponotus (Myrmotrema) auropubens SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 267, 272, and 275.

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA (Liengme).

SOMALILAND: Magala Re Umberto (Ruspoli); Lugh; Lower Ganale (V. Bottego). ABYSSINIA: Boran Galla, Middle Ganale (V. Bottego); Daa (Ruspoli). NATAL: Stamford Hill (I. Trägårdh). BELGIAN CONGO: Stanleyville (Lang and Chapin).

51a₁. Var. *absolon* (SANTSCHI):

Camponotus (Myrmotrema) auropubens var. *absolon* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 267 and 274 (♂, ♀).

Type locality: NATAL (I. Trägårdh).

51a₂. Var. *argentopubens* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 66 (♀).

Camponotus (Myrmotrema) auropubens subsp. *argentopubens* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 268.

Type locality: BENGUELA (C. Wellman).

51a₃. Var. *jacob* (SANTSCHI).

Camponotus (Myrmotrema) auropubens var. *jacob* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 267 and 275 (♂).

Type locality: Pongué Valley, Guengera, MOZAMBIQUE (G. Vasse).

RHODESIA: Bulawayo (G. Arnold).

51b. Subsp. *benguensis* SANTSCHI, 1911, Rev. Zool. Afr., I, p. 213 (♂, ♀, ♂); 1915, Ann. Soc. Ent. France, LXXXIV, p. 269.

Type locality: Cucala, BENGUELA (J. Cruchet).

51c. Subsp. *chrysogaster* EMERY, 1895, Ann. Mus. Civ. Genova, XXXV, p. 182 (♂); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774; 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 604. FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 31; in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 90 (♀, ♂). MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 21 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 136 (♂); 1915, Ann. Soc. Ent. France, LXXXIV, p. 268.

Camponotus chrysogaster EMERY, 1899, Ann. Mus. Civ. Genova, XXXIX, p. 501 (♀).

Type locality: Arussi Galla, Ganale Gudda, ABYSSINIA (V. Bottego).

ABYSSINIA: Buditu to Dimé (V. Bottego). SOMALILAND: Magala Re Umberto (Ruspoli). BRITISH EAST AFRICA: Blue Post Hotel, Kikuyu, 1520 m. (Alluaud and

Jeannel); Fundu Island, W. Pemba (Voeltzkow). GERMAN EAST AFRICA: New Moschi, Mt. Kilimanjaro, 800 m. (Alluaud and Jeannel); Kibonoto, Kilimanjaro; Ngare-na-Nyuki, Meru (Sjöstedt); Mto-ya-Kifaru (Katona). UGANDA: region of Lake Albert, southern Unyoro (C. Alluaud).

51c. Var. **annobonensis** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 268 and 277 (♀).

Camponotus (Myrmotrema) foraminosus var. *annobonensis* SANTSCHI, 1920, 'Études Maladies Parasites Cacaoyer S. Thomé,' X, p. 4.

Camponotus (Myrmotrema) perrisi var. *insularis* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. (1910-11),' I, p. 399, fig. 12 (♀, ♂).

Type locality: ANNOBON (Reichensperger; Schultze).

51d. Subsp. **cuitensis** FOREL, 1901, Mitth. Schweiz. Ent. Ges., X, p. 309 (♀); in Baum, 1903, 'Kunene-Sambesi Expedition,' p. 563 (♀).

Camponotus (Myrmotrema) robbechii subsp. *troglodytes* var. *cuitensis* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 271.

Type locality: Between the Cubango and Cuito Rivers, MOSSAMEDES (Baum and Van der Kellen).

51e. Subsp. **delagoensis** FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 68 (♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774.

Camponotus (Myrmotrema) olivieri var. *delagoensis* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 270.

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA (P. Berthoud; Liengme).

51f. Subsp. **flavus** STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 397, fig. 11 (♀).

Type locality: Fort Crampel, FRENCH CONGO (Schubotz).

51g. Subsp. **haereticus** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 382 (♀). See p. 250.

Camponotus (Myrmotrema) haereticus SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 269.

Type locality: Olokemeji, SOUTHERN NIGERIA (F. Silvestri).

GOLD COAST: Aburi (F. Silvestri). BELGIAN CONGO: Lukolela (Lang and Chapin).

51h. Subsp. **lemma** FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 176 (♀, ♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774. H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 36. FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 357 (♀).

Camponotus lemma DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 238.

Camponotus (Myrmotrema) olivieri var. *lemma* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 270.

Type locality: Malange, ANGOLA (M. Buchner).

BELGIAN CONGO: Lukula (Daniel); Banza Masola; Mondombe; Ganda Sundi; Congo da Lemba (R. Mayné). ZANZIBAR: (Hildebrandt).

51h. Var. **infelix** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 254 (♀).

Camponotus (Myrmotrema) olivieri subsp. *infelix* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 270.

Type locality: Durban, NATAL (G. Arnold).

51h. Var. **moshianus** FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 31 (♀).
Camponotus (Myrmotrema) olivieri var. *moshiana* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 270.

Type locality: Moschi, GERMAN EAST AFRICA (Katona).

51i. Subsp. **orthodoxus** SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 381 (♀).

Camponotus (Myrmotrema) orthodoxus SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 267.

Type locality: Kindia, FRENCH GUINEA (F. Silvestri).

51j. Subsp. **ruspolii** FOREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 349 (♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774. FOREL, 1914, Rev. Suisse Zool., XXII, p. 270. See p. 1050.

Camponotus ruspolii DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 250.

Camponotus (Myrmotrema) grandidieri subsp. *ruspolii* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 269.

Camponotus (Myrmotrema) ruspolii EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Type locality: SOMALILAND (C. Keller).

51j. Var. **rollei** FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 270 (♀); 1913, Rev. Zool. Afr., II, p. 341 (♀). BEQUAERT, *ibid.*, p. 430.

Camponotus (Myrmotrema) grandidieri subsp. *ruspolii* var. *rollei* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 269.

Type locality: Zoutpansberg, 800 m., northern TRANSVAAL (Rolle).

BELGIAN CONGO: Sankisia (J. Bequaert).

52. **Camponotus (Myrmotrema) galla** FOREL. SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 267. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus galla FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 68 (♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 775; 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 604; 1899, *ibid.*, XXXIX, p. 501 (♀). MAYR, in Jägerskiöld, 1903, 'Exped., IX, Formicid.', p. 11; 1907, Sitzb. Ak. Wiss. Wien, math. naturw. Kl., CXVI, Abt. I, p. 389. FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 372.

Camponotus foraminosus subsp. *latinotus* FOREL, 1907, Rev. d'Ent. Caen, XXVI, p. 144 (♀).

Camponotus erinaceus subsp. *galla* FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 271. KARAWAIEW, 1911, Rev. Russe Ent., XI, p. 12 (♀). EMERY, 1915, Boll. Lab. Zool. Gen. Agrar. Portici, X, p. 4 (♀).

Camponotus (Orthonotomyrmex) galla FOREL, 1914, Rev. Suisse Zool., XXII, p. 273.

Type locality: SOUTHERN ABYSSINIA (Ilg).

ERITREA: Nefasit (K. Escherich); Mayabal (F. Silvestri). ABYSSINIA: Hauacio (Ruspoli); Upper Aouache-Endessa (de Rothschild); Bass Narok (V. Bottego). ANGLO-EGYPTIAN SUDAN: Kaka, White Nile (I. Trägårdh); Renk (F. Werner); Khartum; Port Sudan (Karawaiew). BRITISH EAST AFRICA: Lake Rudolf (de Rothschild).

53. **Camponotus (Myrmotrema) grandidieri** FOREL, 1914, Rev. Suisse Zool., XXII, p. 270. SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 268 and 272. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258. See p. 1050.

Camponotus grandidieri FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, C. R., p. ciii (♀). DISTANT, 1892, 'A Naturalist in the Transvaal,' p. 211. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 70. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 150. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 775. MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 21.

Camponotus foraminosus subsp. *grandidieri* FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 89 (♀, ♂, ♂); 1913, Ann. Soc. Ent. Belgique, LVII, p. 145 (♀, ♂).

Camponotus foraminosus var. *grandidieri* (?) MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 195 (♀). H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 37.

GERMAN EAST AFRICA: Ngare na Nyuki, Meru (Sjöstedt). BRITISH EAST AFRICA: Fundu Island, W. Pemba (Voeltzkow). RHODESIA: Bulawayo (G. Arnold). PORTUGUESE EAST AFRICA: Quilimane (Stuhlmann); Mozambique Island (H. Brauns); Delagoa (Liengme). TRANSVAAL: Pretoria (Distant).

53. Var. *atrabilis* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 269 and 273 (♀). See p. 1051.

MOZAMBIQUE: Gorongosa (G. Vasse).

53a. Subsp. *mendax* (EMERY) SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 269 and 274 (♀).

Camponotus foraminosus subsp. *mendax* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 54 (♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774. FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 30 (♀, ♀).

Type locality: Hamman's Kraal, TRANSVAAL (E. Simon).

NATAL. CAPE PROVINCE: (Krebs). BECHUANALAND: Khakhea to Kang (L. Schultze). GERMAN SOUTHWEST AFRICA: (Lübbert).

54. *Camponotus (Myrmotrema) ilgi* FOREL, 1914, Rev. Suisse Zool., XXII, p. 270. SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 270. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus ilgi FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 64 (♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 775. SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 314 (♀).

Type locality: SOUTHERN ABYSSINIA (Ilg).

SENEGAMBIA: St. Louis (Claveau).

55. *Camponotus (Myrmotrema) micipsa* Wm. M. WHEELER. See p. 252 (♀).

Type locality: Between Leopoldville and Yumbi, BELGIAN CONGO (Lang and Chapin).

56. *Camponotus (Myrmotrema) olivieri* (FOREL) SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 270 and 272. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus foraminosus subsp. *olivieri* FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 175 (♀, ♀). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 150. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774. SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393. FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 356 (♀).

Camponotus olivieri DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 246.

Type locality: Malange, ANGOLA (M. Buchner).

GOLD COAST: Akra. FRENCH CONGO: Gaboon (H. Brauns); Brazzaville (A. Weiss). BELGIAN CONGO: Congo da Lemba (R. Mayné).

56i. Var. **sorptus** (FOREL) SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 270. See p. 252.

Camponotus foraminosus subsp. *delagoensis* var. *sorpta* FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 62 (♀, ♂).

Type locality: Lower CONGO, in stomach of *Manis temmincki* (Solon).

BELGIAN CONGO: Kwamouth; Leopoldville; Lukolela; Stanleyville (Lang and Chapin).

56j. Var. **tenuipilis** (EMERY) SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 270.

Camponotus foraminosus subsp. *olivieri* var. *tenuipilis* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 54 (♀).

Type locality: Pretoria, TRANSVAAL (E. Simon).

56a. Subsp. **concordia** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 269 and 272 (♀).

Type locality: Cucala, BENGUELA (J. Cruchet).

56b. Subsp. **tauricollis** (FOREL) SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 270.

Camponotus foraminosus subsp. *tauricollis* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 68 (♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774.

Type locality: Delagoa Bay, PORTUGUESE EAST AFRICA (Liengme).

56b. Var. **pax** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 270 and 273 (♀).

Type locality: CAMEROON.

57. **Camponotus (Myrmotrema) perrisi** FOREL, 1914, Rev. Suisse Zool., XXII, p. 270. SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 272. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus foraminosus subsp. *perrisi* FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 177 (♀, ♂). H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 36.

Camponotus perrisi DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 247.

Camponotus perrisi EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 775.

Camponotus foraminosus subsp. *perrini* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393 (♀).

Type locality: Malange, ANGOLA (M. Buchner).

FRENCH CONGO: (A. Weiss). BRITISH EAST AFRICA: Kitui (Hildebrandt).

57i. Var. **densipunctatus** STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 399 (♀, ♂).

Type locality: Duma, BELGIAN CONGO (Schubotz).

FRENCH CONGO: Fort Crampel; Fort de Possel to Fort Crampel (Schubotz).

57a. Subsp. **jucundus** SANTSCHI, 1911, Rev. Zool. Afr., I, p. 216 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 314; 1915, Ann. Soc. Ent. France, LXXXIV, p. 272. See p. 251.

Camponotus (Myrmotrema) perrisi subsp. *jucundus* STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 399 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

FRENCH CONGO: Mandouga (A. Weiss). BELGIAN CONGO: Ubangi; Banzyville (Augustin); Garamba; Faradje; Niangara; Kwamouth (Lang and Chapin); Kimuenza (Schultze).

57a₁. Var. **grandior** (FOREL) SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 272. See p. 252.

Camponotus (*Myrmotrema*) *foraminosus* subsp. *perrisi* var. *grandior* FOREL, 1913, Rev. Zool. Afr., II, p. 342 (♀).

Camponotus foraminosus subsp. *perrisi* var. *grandis* BEQUAERT, 1913, *ibid.*, II, p. 431.

Type locality: Welgelegen, BELGIAN CONGO (J. Bequaert).

BELGIAN CONGO: Yakuluku; Garamba (Lang and Chapin).

57b. Subsp. **nigeriensis** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 272 and 277 (♀, ♀, ♂).

Camponotus (*Myrmotrema*) *bayeri* subsp. *nigeriensis* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 383 (♀).

Type locality: Ibadan, SOUTHERN NIGERIA (F. Silvestri).

SOUTHERN NIGERIA: Lagos (F. Silvestri). BELGIAN CONGO: Kitempuka; Katakai (Gérard).

58. **Camponotus** (*Myrmotrema*) **postoculatus** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 254 (♀).

Type locality: Durban, NATAL (G. Arnold).

59. **Camponotus** (*Myrmotrema*) **rhamses** SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 268 and 275 (♀).

Type locality: Lake No, Upper EGYPT (Reichensperger).

60. **Camponotus** (*Myrmotrema*) **robecchii** EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus robecchii EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 120, fig. (♀). DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 250. EMERY, 1896, Mem. Accad. Sc. Bologna, (V) 5, p. 774; 1901, Bull. Soc. Ent. Italiana, XXXIII, p. 63 (♀).

Camponotus foraminosus subsp. *robecchii* FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 90 (♀).

Camponotus (*Myrmotrema*) *foraminosus* subsp. *robecchii* FOREL, 1914, Rev. Suisse Zool., XXII, p. 270.

Camponotus (*Myrmotrema*) *robecchii* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, pp. 271 and 272.

Type locality: Obbia, SOMALILAND (Brichetti-Robecchi).

BRITISH EAST AFRICA: Chake-Chake, Pemba (Voeltzkow). ERITREA: Massaua (Belli).

61. **Camponotus** (*Myrmotrema*) **troglodytes** FOREL, 1914, Rev. Suisse Zool., XXII, p. 270; 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 254 (♀). SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 40 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 137 (♀). EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus troglodytes FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 69 (♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 775. FOREL, 1904, Ann. Mus. Zool. Ac. Sc. St. Pétersbourg, VIII, (1903), p. 377 (♀); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 29 (♀).

Camponotus (*Myrmotrema*) *robecchii* subsp. *troglodytes* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 271.

Type locality: Delagca Bay, PORTUGUESE EAST AFRICA (Liengme).

GERMAN SOUTHWEST AFRICA: (Lübbert). BECHUANALAND: Khakhea to Kang (L. Schultze); Gröotfontein (v. Erffer). TRANSVAAL: Lydenburg (Wilms). CAPE

PROVINCE: Cape Town (Wilms). ZULULAND: Dukudu (I. Trägårdh). NATAL: Durban (G. Arnold). BRITISH EAST AFRICA: Ramisi River (Alluaud and Jeannel). ABYSSINIA: Southern Danakil; Harar (Kachovskij).

61. Var. **abyssinicus** SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 314 (♀).

Camponotus (Myrmotrema) robbeckii subsp. *troglodytes* var. *abyssinica* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 271.

Type locality: Diré Dawa, ABYSSINIA (J. Roger).

61a. Var. **rhodesianus** FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 145 (♂).

Camponotus (Myrmotrema) robbeckii subsp. *rhodesiana* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 271.

Type locality: Redbank, SOUTHERN RHODESIA (G. Arnold).

SOUTHERN RHODESIA: Sebakwe (G. Arnold).

Subgenus 6. **Myrmopiromis** WHEELER

Camponotus subg. *Myrmopiromis* WHEELER, 1921, Psyche, XXVIII, p. 17.

Camponotus subg. *Myrmepomis* EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 245 (not of Forel, 1912).

Subgenotype: *Formica fulvopilosa* De Geer, 1778.

62. **Camponotus (Myrmopiromis) conradti** FOREL.

Camponotus (Camponotus) conradti FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) I, p. 256 (♀).

Camponotus (Myrmepomis) conradti EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Type locality: CAMEROON (Conradt).

63. **Camponotus (Myrmopiromis) fulvopilosus** (DE GEER).

Formica fulvopilosa DEGEER, 1778, 'Mém. Hist. Ins.,' VII, p. 613, Pl. XLV, figs. 13, 14. GÖTZE, in DeGeer, 1783, 'Abh. Gesch. Insekt.,' VII, p. 218 (♂), Pl. XLV, figs. 13, 14. RETZIUS, 1783, 'Gen. et Spec. Insect.,' p. 75. LEPELETIER, 1836, 'Hist. Nat. Ins. Hym.,' I, p. 213. F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 32.

Formica pilosa OLIVIER, 1791, 'Encycl. Méthod. Insect.,' VI, p. 498.

Formica rufiventris FABRICIUS, 1804, 'Syst. Piez.,' p. 409 (♀).

Camponotus fulvopilosus MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 668 (♂); 1863, *ibid.*, XIII, p. 399. ROGER, 1863, 'Verzeich. Formicid.,' p. 2. MAYR, 1865, 'Reise Novara, Zool.,' II, Formicide, p. 36. FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 108 (♂); 1886, Ann. Soc. Ent. Belgique, XXX, p. 192 (♀). EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 356 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 232. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 54. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 150. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774; 1900, Bull. Soc. Ent. Italiana, XXXII, p. 117, fig. 12 (♀). BINGHAM, 1903, Ann. Mag. Nat. Hist., (7) XII, p. 63 (♀). DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, pp. 349 and 363. FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 25 (♂, ♂).

Camponotus (Myrmepomis) fulvopilosus FOREL, 1914, Rev. Suisse Zool., XXXII, p. 273. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Type locality: Cape of Good Hope.

CAPE PROVINCE: Kimberley (E. Simon); Tulbagh. BASUTOLAND: Lessouto (P. Berthoud). GERMAN SOUTHWEST AFRICA: (Lübbert); Okahandja (Casper; Peters); Grootfontein (v. Erffer); Windhoek (Kunze and L. Schultze); Bethanien (Schenck); Salem (L. Schultze). PORTUGUESE CONGO: Chinchoxo (Falkenstein). BENGUELA. RHODESIA: Victoria Falls; Tsessebe Station (Dixey and Longstaff).

63. Var. *detritoides* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 25 (♀).

Type locality: Glatkop, Little Namaland, CAPE PROVINCE (L. Schultze).

63. Var. *flavopilosus* EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 54 (♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774. FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 25 (♀, ♂); 1911, Sitzb. Bayer. Akad. Wiss., p. 291 (♀). EMERY, 1912, Annuar. Mus. Zool. Univ. Napoli, N. S., III, No. 26, p. 2 (♀).

Type locality: De Aar, CAPE PROVINCE (E. Simon).

GERMAN SOUTHWEST AFRICA: (G. v. Bayern); Salem; Kubub (L. Schultze). CAPE PROVINCE: Namaland (L. Péringuey). BECHUANALAND: Kooa (L. Schultze). BELGIAN CONGO: Luapula River.

63a. Subsp. *brevisetosus* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 25 (♀, ♀).

Type locality: NATAL (Haviland).

CAPE PROVINCE: Cape Town (Wilms).

63b. Subsp. *detritus* (EMERY) FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 26 (♀).

Camponotus detritus EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 357 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 228. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774.

Camponotus (Myrmepomis) detritus EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Type locality: Damaraland, GERMAN SOUTHWEST AFRICA.

GERMAN SOUTHWEST AFRICA: Rooibank near the Walfish Bay (L. Schultze).

63c. Subsp. *storeatus* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 26 (♀).

Type locality: Willowmore, CAPE PROVINCE (H. Brauns).

64. *Camponotus (Myrmopiromis) maynei* FOREL.

Camponotus (Orthonotomyrmex) maynei FOREL, 1916, Rev. Suisse Zool., XXIV, p. 445 (♀, ♀).

Camponotus (Myrmepomis) maynei EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Type locality: BELGIAN CONGO (R. Mayné).

65. *Camponotus (Myrmopiromis) niveosetosus* MAYR. See p. 1052.

Camponotus niveosetosus MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 665 (♂); 1863, *ibid.*, XIII, p. 400. ROGER, 1863, 'Verzeich. Formicid.,' p. 2. MAYR, 1865, 'Reise Novara, Zool.,' II, Formicidae, p. 35 (♀, ♀), Pl. I, fig. 3; 1866, Verh. Zool. Bot. Ges. Wien, XVI, p. 885 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 245. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 55; 1896, Mem. Accad. Sc. Bologna, (5) V, p. 775. FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 89 (♂); in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 30 (♀).

Camponotus (Myrmoturba) niveosetosus FOREL, 1914, Rev. Suisse Zool., XXII, p. 267.

Camponotus (Myrmepomis) niveosetosus EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Type locality: Cape of Good Hope (Novara Expedition).

CAPE PROVINCE: Cape Town (E. Simon). BECHUANALAND: Lehututu (L. Schultze). GERMAN SOUTHWEST AFRICA: (L. Schultze). BRITISH EAST AFRICA: Fundu Island, W. Pemba (Voeltzkow).

65a. Subsp. *irredux* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 30 (§); 1910, Ann. Soc. Ent. Belgique, LIV, p. 451.

Camponotus (Myrmoturba) niveosetosus var. *irredux* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 41 (§).

Type locality: NATAL (R. C. Wroughton).

NATAL: (Haviland); Pietermaritzburg (I. Trägårdh).

Subgenus 7. *Myrmorhachis* FOREL

Camponotus subg. *Myrmorhachis* FOREL, 1912, Mém. Soc. Ent. Belgique, XX, p.

92.

Camponotus subg. *Myrmacantha* EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 246.

Subgenotype: *Camponotus polyrhachioides* Emery, 1898.

66. *Camponotus (Myrmorhachis) aberrans* MAYR. FOREL, 1914, Rev. Suisse Zool., XXII, p. 274. SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 264 (§).

Camponotus aberrans MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 152 (♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774. SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393 (§).

Camponotus (Myrmacantha) aberrans EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Type locality: GOLD COAST.

FRENCH CONGO: Brazzaville (A. Weiss). BELGIAN CONGO: Kitempuka (Gérard).

67. *Camponotus (Myrmorhachis) polyrhachioides* EMERY. FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 357 (§); 1914, Rev. Suisse Zool., XXII, p. 274. See p. 254.

Polyrhachis paradoxa ERN. ANDRÉ, 1892, Rev. d'Ent. Caen, XI, p. 46 (§). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 266. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 153. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 1 (§). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 778.

Camponotus polyrhachioides EMERY, 1898, Rend. Accad. Sc. Bologna, N. S., II, p. 227; 1900, Bull. Soc. Ent. Italiana, XXXII, p. 117, fig. 11a (§). FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 74 (§). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393 (§).

Camponotus paradoxus STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 149 (§, ♀) (nec *Camponotus paradoxus* Mayr, 1866).

Camponotus (Myrmacantha) polyrhachioides EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Type locality: Gaboon, FRENCH CONGO.

SIERRA LEONE. LIBERIA: Grand Bassa (H. Brauns). CAMEROON: Mundame (Conradt); Longji (Paschen). FRENCH CONGO: Ogowe (Mocquerys); Sette-Cama (Hupfer); Brazzaville (A. Weiss). BELGIAN CONGO: Mayombe; Kiniati (R. Mayné); Lie; Lukolela (Lang and Chapin).

Subgenus 8. **Myrmopsamma** FOREL

Camponotus subg. *Myrmopsamma* FOREL, 1914, Rev. Suisse Zool., XXII, p. 261. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 246.

Subgenotype: *Camponotus mystaceus* Emery, 1886.

68. ***Camponotus (Myrmopsamma) cuneiscapus*** FOREL, 1914, Rev. Suisse Zool., XXII, p. 270. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus cuneiscapus FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 29 (♀), Pl. 1, fig. 2.

Type locality: Steinkopf, CAPE PROVINCE (L. Schultze).

69. ***Camponotus (Myrmopsamma) mystaceus*** EMERY. FOREL, 1914, Rev. Suisse Zool., XXII, p. 270. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus mystaceus EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 356 (♀, ♂), Pl. XVII, fig. 3. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 244. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 768. FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 27 (♀).

Type locality: Cape of Good Hope (L. Péringuey).

GERMAN SOUTHWEST AFRICA: Okahandja (Casper); Windhoek (L. Schultze). BECHUANALAND: Severelela to Khakhea; Kang to Lehututu; Koa to Sekgoma (L. Schultze).

60. Var. ***exsanguis*** FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 28 (♀, ♀, ♂).

Type locality: Prince of Wales Bay, South of Lüderitzbucht, GERMAN SOUTHWEST AFRICA (L. Schultze).

CAPE PROVINCE: Steinkopf (L. Schultze).

69. Var. ***kamæ*** FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 27 (♀).

Type locality: Rooibank, near Walfish Bay, GERMAN SOUTHWEST AFRICA (L. Schultze).

70. ***Camponotus (Myrmopsamma) simulans*** FOREL, 1914, Rev. Suisse Zool., XXII, p. 270. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Camponotus simulans FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 28 (♀), Pl. 1, fig. 4.

Type locality: Prince of Wales Bay, south of Lüderitzbucht, GERMAN SOUTHWEST AFRICA (L. Schultze).

Subgenus 9. **Myrmamblys** FOREL

Camponotus subg. *Myrmamblys* FOREL, 1912, Mém. Soc. Ent. Belgique, XX, p. 90.

Camponotus subg. *Myrmotemnus* EMERY, 1920, (in part), Rev. Zool. Afr., VIII, p. 246.

Subgenotype: *Camponotus reticulatus* Roger, 1863.

71. ***Camponotus (Myrmamblys) bedoti*** EMERY.

Camponotus bedoti EMERY, 1893, Rev. Suisse Zool., I, p. 196 (♀), Pl. VIII, fig. 2.

Type locality: Batjan, MOLECCAS (Bedot).

MALAYSIA.

71. Var. **klugi** EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 51 (♀).
Camponotus reticulatus subsp. *yerburyi* var. *klugi* EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 772.
 Type locality: CAPE TOWN, CAPE PROVINCE (E. Simon).
 72. **Camponotus (Myrmamblys) bertolonii** EMERY.
Camponotus bertolonii EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 51 (♀, ♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 771. MAYR, 1901, Ann. Naturh. Hofmus. Wien, XVI, p. 28 (♀, ♀, ♂).
Camponotus (Myrmoturba) bertolonii FOREL, 1914, Rev. Suisse Zool., XXII, p. 267.
Camponotus (Myrmotenus) bertolonii EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.
 Type locality: CAPE TOWN, CAPE PROVINCE (E. Simon).
 CAPE PROVINCE: Port Elizabeth (H. Brauns).
 73. **Camponotus (Myrmamblys) brookei** FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 259 (♀).
 Type locality: Stella Bush, Durban, NATAL (Marley).
 74. **Camponotus (Myrmamblys) chapini** Wm. M. WHEELER. See p. 254 (♀).
 Type locality: Garamba, BELGIAN CONGO (Lang and Chapin).
 BELGIAN CONGO: Medje; Faradje (Lang and Chapin).
 75. **Camponotus (Myrmamblys) emarginatus** EMERY.
Camponotus emarginatus EMERY, 1886, Bull. Soc. Ent. Italiana, XVIII, p. 358 (♀, ♀, ♂), Pl. XVII, fig. 2. DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 229. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 771.
Camponotus (Myrmophyma) emarginatus FOREL, 1914, Rev. Suisse Zool., XXII, p. 269.
Camponotus (Myrmotenus) emarginatus EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.
 Type locality: Cape of Good Hope, CAPE PROVINCE (L. Péringuey).
 76. **Camponotus (Myrmamblys) ferrerii** FOREL, 1914, Rev. Suisse Zool., XXII, p. 271.
Camponotus ferrerii FOREL, 1913, *ibid.*, XXI, p. 671 (♀).
 Type locality: CAMEROON (Conradt).
 76a. Subsp. **akka** FOREL, 1916, Rev. Suisse Zool., XXIV, p. 446 (♀, ♀).
 Type locality: BELGIAN CONGO (Kohl).
 77. **Camponotus (Myrmamblys) limbiventris** SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 134 (♀).
Camponotus limbiventris SANTSCHI, 1910, Ann. Soc. Ent. France, LXXIX, p. 366, fig. (♀).
Camponotus (?Myrmotenus) limbiventris EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.
 Type locality: Kibosho, Mt. Kilimanjaro, GERMAN EAST AFRICA (C. Alluaud).
 BRITISH EAST AFRICA: Amboni River, Mt. Kenia, 1800 m. (Alluaud and Jeannel).
 78. **Camponotus (Myrmamblys) nasutus** EMERY.
Camponotus nasutus EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 53 (♀), Pl. II, fig. 27; 1896, Mem. Accad. Sc. Bologna, (5) V, p. 771.
Camponotus (Myrmophyma) nasutus FOREL, 1914, Rev. Suisse Zool., XXII, p. 269.

Camponotus (Myrmetennus) nasutus EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Type locality: Pretoria, TRANSVAAL (E. Simon).

78. Var. *quinque-dentatus* FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 451 (♀).

Type locality: NATAL (Haviland).

79. *Camponotus (Myrmamblys) orinobates* SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 348, fig. 4c (♀).

Camponotus (Myrmamblys) bertolonii SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 134 (♀) (nec Emery).

Type locality: eastern slope of Mt. Kenia, BRITISH EAST AFRICA (Alluaud and Jeannel).

BRITISH EAST AFRICA: River Amboni, 1800 m.; Kijabe, Kikuyu Escarpment, 2100 m. (Alluaud and Jeannel).

80. *Camponotus (Myrmamblys) schoutedeni* FOREL.

Camponotus schoutedeni FOREL, 1911, Rev. Zool. Afr., I, p. 281 (♀); 1912, Ent. Mitt. Deutsch. Ent. Mus. Berlin, I, p. 83; 1913, Ann. Soc. Ent. Belgique, LVII, pp. 146 (♀) and 357 (♀).

Camponotus (Myrmosaga) schoutedeni FOREL, 1913, Rev. Zool. Afr., II, p. 347 (♀); 1914, Rev. Suisse Zool., XXII, p. 269. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 137 (♀).

Type locality: Congo da Lemba, BELGIAN CONGO (R. Mayné).

BELGIAN CONGO: Banza Masola; Yandumba; Ganda Sundi; Mandungu; Yambata; Kiniati (R. Mayné); Beni (Bergerhoff). BENGUELA: (C. Wellman). RHODESIA: Redbank (G. Arnold). GERMAN EAST AFRICA: New Moschi, Mt. Kilimanjaro, 800 m. (Alluaud and Jeannel). BRITISH EAST AFRICA: Likoni; Shimoni; Kikuyu Escarpment (Alluaud and Jeannel).

81. *Camponotus (Myrmamblys) simus* EMERY. FOREL, 1914, Rev. Suisse Zool., XXII, p. 272.

Camponotus simus EMERY, 1908, Ann. Soc. Ent. Belgique, LII, p. 188, fig. 2 (♀, ♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393 (♀).

Camponotus (Myrmetennus) simus EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Type locality: Sankuru, BELGIAN CONGO (Luija).

FRENCH CONGO: Brazzaville (A. Weiss).

81a. Subsp. *manidis* FOREL, 1911, Rev. Zool. Afr., I, p. 282 (♀).

Camponotus manidis FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 62 (♀).

Type locality: Lower CONGO, in stomach of *Manis temmincki* (Solon).

82. *Camponotus (Myrmamblys) viri* SANTSCHI, 1915, Ann. Soc. Ent. France, LXXXIV, p. 265, fig. 13 (♀).

Camponotus (Myrmetennus) viri EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 258.

Type locality: Mombasa, BRITISH EAST AFRICA.

Subgenus 10. *Colobopsis* MAYR

Colobopsis MAYR, 1861, 'Europ. Formicid.,' p. 38.

Camponotus subg. *Colobopsis* EMERY, 1889, Ann. Mus. Civ. Genova, XXVII, p. 517. FOREL, 1912, Mém. Soc. Ent. Belgique, XX, p. 90; 1914, Rev. Suisse Zool., XXII, p. 263. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 247.

Subgenotype: *Formica truncata* Spinola, 1808.

83. **Camponotus (Colobopsis) bifossus** SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 293 (2).

Type locality: Caledon, CAPE PROVINCE (L. Péringuey).

Species Incertæ Sedis

84. **Camponotus æquitas** SANTSCHI.¹

Camponotus (Myrmamblys) æquitas SANTSCHI, 1920, Ann. Soc. Ent. France, LXXXVIII, (1919), p. 387, fig. 16 (♀, ♂).

Type locality: Bulawayo, RHODESIA (G. Arnold).

85. **Camponotus berthoudi** FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 108 (♀), Pl. I, fig. 6. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 222. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774.

Camponotus (Myrmamblys) berthoudi FOREL, 1914, Rev. Suisse Zool., XXII, p. 271.

Type locality: Valdezia, TRANSVAAL (P. Berthoud).

86. **Camponotus buchholzi** MAYR, 1902, Verh. Zool. Bot. Ges. Wien, LII, p. 297 (♀, ♂).

Camponotus (Myrmamblys) buchholzi FOREL, 1914, Rev. Suisse Zool., XXII, p. 271.

Type locality: Victoria, CAMEROON (R. Buchholz).

86. Var. **furvus** SANTSCHI, 1911, Rev. Zool. Afr., I, p. 213 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

87. **Camponotus dewitzi** FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 187 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 228. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 775.

Camponotus (Myrmosaga) dewitzi FOREL, 1914, Rev. Suisse Zool., XXII, p. 269.

Type locality: Chinchoxo, PORTUGUESE CONGO (Falkenstein).

88. **Camponotus favorabilis** SANTSCHI.

Camponotus (Myrmosaga) favorabilis SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 349 (♀).

Type locality: Nyamandloru, RHODESIA (G. Arnold).

89. **Camponotus jeanneli** SANTSCHI.

Camponotus (Myrmosaga) jeanneli SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 137, fig. 30 (♀).

Type locality: Mombasa, BRITISH EAST AFRICA (Alluaud and Jeannel).

BRITISH EAST AFRICA: Likoni; Ramisi; Tchania River, Kikuyu, 1520 m. (Alluaud and Jeannel).

90. **Camponotus lilianæ** FOREL. BEQUAERT, 1913, Rev. Zool. Afr., II, p. 431.²

Camponotus (Myrmamblys) lilianæ FOREL, 1913, ibid., II, p. 344 (♀); 1914, Rev. Suisse Zool., XXII, p. 271. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 134 (♀).

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

BRITISH EAST AFRICA: Mbuyuni, Pori, 1150 m.; Taveta, 750 m. (Alluaud and Jeannel).

¹Placed by Santschi (1921) in the subgenus *Myrmosphincta*.

²Placed by Santschi (1921) in the subgenus *Myrmamblys*.

90a. Subsp. **cornutus** FOREL, 1913, Rev. Zool. Afr., II, p. 345 (♀).

Type locality: Elisabethville, BELGIAN CONGO (J. Bequaert).

91. **Camponotus orinodromus** SANTSCHI.

Camponotus (Myrmamblys) orinodromus SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 347; fig. 4e (♀).

Type locality: Mt. Matroosberg, CAPE PROVINCE (R. W. Tucker).

92. **Camponotus orites** SANTSCHI.

Camponotus (Myrmamblys) orites SANTSCHI, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 346, fig. 4d (♀).

Type locality: Mt. Matroosberg, CAPE PROVINCE (R. W. Tucker).

93. **Camponotus ostiarius** FOREL.

Camponotus (Myrmamblys) ostiarius FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 258 (♀).

Type locality: Durban, NATAL (G. Arnold).

94. **Camponotus yvonnæ** FOREL.¹

Camponotus (Myrmamblys) yvonnæ FOREL, 1919, Bull. Soc. Vaudoise Sc. Nat., (5) LII, p. 480 (♀).

Type locality: Stanleyville, BELGIAN CONGO (Kohl).

Phasmomyrmex STITZ

Phasmomyrmex STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 146. EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 252.

Camponotus subg. *Phasmomyrmex* FOREL, 1914, Rev. Suisse Zool., XXII, p. 264.

Genotype: *Camponotus buchneri* Forel, 1886 (= *Phasmomyrmex sericeus* Stitz, 1910).

1. **Phasmomyrmex buchneri** (FOREL) EMERY, 1920, Rev. Zool. Afr., VIII, 2, p. 260. See p. 256.

Camponotus buchneri FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 183 (♀). ERN. ANDRÉ, 1892, Rev. d'Ent. Caen, XI, p. 45 (♀); DALLA TORRE, 1893, 'Cat. Hym.', VII, p. 223. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 151 (♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 774; 1900, Bull. Soc. Ent. Italiana, XXXII, p. 116, fig. 10 (♀). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 421 (♀); 1913, Rev. Suisse Zool., XXI, p. 671 (♀).

Phasmomyrmex sericeus STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 146, fig. 11 (♀).

Camponotus (Phasmomyrmex) buchneri FOREL, 1914, Rev. Suisse Zool., XXII, p. 273.

Type locality: Malange, ANGOLA (M. Buchner).

CAMEROON: Mundame (Conradt); Lomie (Thesing). FRENCH CONGO: Ligny (H. Brauns). BELGIAN CONGO: Lukolela; Avakubi; Medje (Lang and Chapin); Lubutu (J. Bequaert).

Polyrhachis F. SMITH²

Polyrhachis F. SMITH, 1858, Journ. Proc. Linn. Soc. London, Zool., II, p. 58. ROGER, 1861, Berlin. Ent. Zeitschr., V, p. 174.

¹Placed by Santschi (1921) in his subgenus *Myrmopelta*.

²I have been unable to find the description of *Polyrhachis setulosus* "Smith" recorded from Angola (Welwitsch) by Radoszkowsky, 1881, Jorn. Sci. As. Lisboa, VIII, No. 31, p. 197.

Hoplostomyrma GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 262.

Formica FABRICIUS, LATREILLE, DRURY, etc.

Genotype: *Formica bhamata* Drury, 1773.

Subgenus 1. *Cyrtomyrma* FOREL

Polyrhachis subg. *Cyrtomyrma* FOREL, 1915, Arkiv f. Zool., IX, No. 16, p. 107.

Subgenotype: *Formica rastellata* Latreille, 1802.

1. *Polyrhachis (Cyrtomyrma) alexisi* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 455, fig. 7 (♀).

Type locality: BELGIAN CONGO (Kohl).

2. *Polyrhachis (Cyrtomyrma) kohli* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 454 (♂, ♀).

Type locality: BELGIAN CONGO (Kohl).

Subgenus 2. *Myrma* BILLBERG

Myrma BILLBERG, 1820, 'Enumer. Insect.,' p. 104. WHEELER, 1911, Science, N. S., XXXIII, p. 859.

Hoplostomyrma GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 262; in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 507.

Subgenotype: *Formica militaris* FABRICIUS, 1781.

3. *Polyrhachis (Myrma) senescens* STITZ.

Polyrhachis senescens STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 151 (♀).

Type locality: CAMEROON (v. Knobloch).

4. *Polyrhachis (Myrma) aërope* Wm. M. WHEELER. See p. 265 (♀).

Type locality: NIANGARA, BELGIAN CONGO (Lang and Chapin).

5. *Polyrhachis (Myrma) alluaudi* EMERY.

Polyrhachis alluaudi EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 567 (♂), Pl. xv, figs. 9 and 10 (nest). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 258. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777.

Type locality: ASSINIE, IVORY COAST (C. Alluaud).

51. Var. *anteplana* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 448 (♀). See p. 266.

Type locality: BELGIAN CONGO (Kohl).

BELGIAN CONGO: Stanleyville (Lang and Chapin).

6. *Polyrhachis (Myrma) arnoldi* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 263 (♀).

Type locality: DURBAN, NATAL (G. Arnold).

7. *Polyrhachis (Myrma) atalanta* Wm. M. WHEELER. See p. 263 (♀).

Type locality: Stanleyville, BELGIAN CONGO (Lang and Chapin).

8. *Polyrhachis (Myrma) bequaerti* Wm. M. WHEELER. See p. 267 (♀).

Type locality: UTIASIKI, between Lubutu and Kirundu, BELGIAN CONGO (J. Bequaert).

9. *Polyrhachis (Myrma) concava* ERN. ANDRÉ. FOREL, 1916, Rev. Suisse Zool., XXIV, p. 448 (♂, ♀). See p. 265.

Polyrhachis concava ERN. ANDRÉ, 1889, Rev. d'Ent. Caen, VIII, p. 218 (♀); 1892, *ibid.*, XI, p. 45 (♂, ♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 260. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 154 (♀). EMERY, 1896, Mem. Accad. Sc.

Bologna, (5) V, p. 777. FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 61 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 149 (♀). FOREL, 1911, Rev. Zool. Afr., I, p. 282 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 359 (♀).

Type locality: SIERRA LEONE (Mocquerys).

CAMEROON: (H. BRAUNS); Mundame (Conradt); Bibundi (Tessmann); Yaunde (Zenker). FRENCH CONGO: Gaboon. BELGIAN CONGO: (Kohl); Congo da Lemba; Mondombe (R. Mayné); Lower Congo, in stomach of *Manis temmincki* (Solon); Stanleyville; Akenge (Lang and Chapin).

10. *Polyrhachis (Myrma) consimilis* F. SMITH.

Polyrhachis consimilis F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 73 (♀), Pl. IV, figs. 30 and 31. ROGER, 1863, 'Verzeich. Formicid.,' p. 6. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 444; 1886, *ibid.*, XXXVI, p. 357. ERN. ANDRÉ, 1887, Rev. d'Ent. Caen, VI, p. 286 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 260. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 779.

Type locality: SIERRA LEONE (D. F. Morgan).

11. *Polyrhachis (Myrma) cornuta* STITZ.

Polyrhachis cornuta STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 150 (♀).

Type locality: Kimpoko, BELGIAN CONGO (Büttner).

12. *Polyrhachis (Myrma) cubaënsis* MAYR.

Polyrhachis cubaënsis MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 686 (♀); 1863, *ibid.*, XIII, p. 444. ROGER, 1863, 'Verzeich. Formicid.,' p. 9. EMERY, 1889, Bull. Soc. Ent. Italiana, XXI, p. 69. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 261. MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 197. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777.

Type locality: Port Natal, NATAL (described as from Cuba by Mayr).

12a. Var. *striolato-rugosa* MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 196 (♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777.

Polyrhachis cubaënsis var. *striolato-rugosa* H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 38.

Type locality: ZANZIBAR (Stuhlmann).

12a. Subsp. *durbanensis* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 262 (♀, ♀).

Type locality: Durban, NATAL (C. B. Cooper).

12b. Subsp. *gallicola* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 71 (♀, ♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777. FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 92 (♀).

Type locality: Delagoa, PORTUGUESE EAST AFRICA (Liengme).

BRITISH EAST AFRICA: Chake Chake, Pemba Island (Voeltzkow).

12c. Subsp. *gerstaeckeri* (FOREL)-EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777.

Polyrhachis gerstaeckeri FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 197 (♀). ERN. ANDRÉ, 1887, Rev. d'Ent. Caen, VI, p. 287. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 262. MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 196.

Polyrhachis cubaënsis var. *gerstaeckeri* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 71.

Polyrhachis gerstaeckeri H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 38.

Type locality: ZANZIBAR (Hildebrandt).

PORTUGUESE EAST AFRICA: Delagoa Bay (Liengme).

12d. Subsp. *wilmsi* FOREL, in Schultze, 1910, 'Forschungsreise in Südafrika,' IV, p. 30 (♀).

Type locality: Lobombo Borges, MOZAMBIQUE (Wilms).

13. *Polyrhachis (Myrma) curta* ERN. ANDRÉ.

Polyrhachis curta ERN. ANDRÉ, 1890, Rev. d'Ent. Caen, IX, p. 312 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 261. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777.

Type locality: SIERRA LEONE (Mocquerys).

14. *Polyrhachis (Myrma) decemdentata* ERN. ANDRÉ. FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 351 (♀, ♀, ♂); 1916, Rev. Suisse Zool., XXIV, p. 450 (♀). See p. 267.

Polyrhachis decemdentata ERN. ANDRÉ, 1889, Rev. d'Ent. Caen, VIII, p. 219 (♀). EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 566 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 261. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 1 (♀, ♀). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 153 (♀, ♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777. MAYR, 1902, Verh. Zool. Bot. Ges. Wien, LII, p. 300 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 149 (♀). FOREL, 1911, Rev. Zool. Afr., I, p. 282 (♀); 1913, Rev. Suisse Zool., XXI, p. 673 (♀).

Type locality: SIERRA LEONE (Mocquerys).

FRENCH GUINEA: Los Islands (H. Brauns). SIERRA LEONE: Sherbro Island (H. Brauns). IVORY COAST: Assinie (C. Alluaud). CAMEROON: (Conradt); Bibundi (Tessmann). FERNANDO PO: (R. Buchholz). FRENCH CONGO: Ogowe (Mocquerys). BELGIAN CONGO: Congo da Lemba (R. Mayné); St. Gabriel (Kohl); Stanleyville (Lang and Chapin); Malela (J. Bequaert).

14. Var. *fernandensis* FOREL, 1901, Ann. Soc. Ent. Belgique, XLV, p. 377 (♀). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 396 (♀, ♀).

Type locality: FERNANDO PO (Conradt).

FRENCH CONGO: Brazzaville (A. Weiss).

14. Var. *flavipes* STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 149 (♀, ♀).

Type locality: Victoria, CAMEROON (Faber).

15. *Polyrhachis (Myrma) fissa* MAYR.

Polyrhachis fissa MAYR, 1902, Verh. Zool. Bot. Ges. Wien, LII, p. 301 (♀, ♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 149 (♀, ♀).

Type locality: Victoria, CAMEROON (R. Buchholz).

CAMEROON: Bibundi (Tessmann); Mundame (Conradt). SPANISH GUINEA: Nkolentanga (Tessmann).

16. *Polyrhachis (Myrma) gagates* F. SMITH. SANTSCHI, 1914 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 139 (♀). See p. 262.

Polyrhachis gagates F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 71 (♀). Pl. IV, fig. 14. ROGER, 1863, 'Verzeich. Formicid.,' p. 6. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 444. GERSTÄCKER, 1871, Arch. f. Naturg., XXXVII, 1, p. 354; 1873, in 'v. d. Decken's Reisen in Ost Afrika, Gliederthiere,' p. 342 (♀). FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 117 (♀); 1886, Ann. Soc. Ent. Belgique, XXX, p. 194 (♀, ♀). MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 357. ERN. ANDRÉ, 1887, Rev. d'Ent. Caen, VI, p. 287 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 262. MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 197. WASMANN, 1894, 'Verzeichn. Myrmecoph. Termitoph. Arthrop.,' p. 184. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 72. MAYR, 1895, Ann.

Naturh. Hofmus. Wien, X, p. 154. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777. MAYR, in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 22. DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, p. 375. FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 38 (♀); in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 92 (♀); 1909, Ann. Soc. Ent. Belgique, LIII, p. 54. ZAVATTARI, 1909, 'Il Ruwenzori, Parte Scientif.,' I, p. 215 (♀). STITZ, 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08),' III, p. 392 (♀). FOREL, 1913, Rev. Zool. Afr., II, p. 349 (♀).

Polyrachis gagates H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 38.

Type locality: Port Natal, NATAL.

BELGIAN CONGO: Zambézi (Lang and Chapin); Boma (H. Brauns); Beni (Borgerhoff); Kasindi (Schubotz). ANGOLA: Malange (M. Buchner). UGANDA: Katende (Duke of Abruzzi). BRITISH EAST AFRICA: Mombasa (v. d. Decken); Kitui (Hildebrandt); Chake Chake, Pemba Island (Voeltzkow). GERMAN EAST AFRICA: Tanga (H. Brauns; Alluaud and Jeannel); Mt. Kilimanjaro (Bornemisza); Kibonoto, Mt. Kilimanjaro, 1000-1200 m.; Mombo, Usambara (Sjöstedt); Rosako, Usaramo (Stuhlmann). ZANZIBAR: (Hildebrandt). PORTUGUESE EAST AFRICA: Delagoa (Liengme). TRANSVAAL: Valdezia (P. Berthoud). CAPE PROVINCE: East London (Dixey and Longstaff).

16a. Subsp. *indefinita* FOREL, 1913, Rev. Zool. Afr., II, p. 349 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 147 (♀).

Polyrachis gagates var. *indefinita* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 42 (♀). TRÄGÅRDH, *ibid.*, p. 47.

Polyrachis (Myrma) gagates var. *indefinita* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 139 (♀).

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

NATAL: Amanzimtoti (I. Trägårdh). SOUTHERN RHODESIA: Sebakwe (G. Arnold). BRITISH EAST AFRICA: Voi, Wa-Taita, 600 m. (Alluaud and Jeannel).

17. *Polyrachis (Myrma) gamai* SANTSCHI, 1917, Ann. Soc. Ent. France, LXXXV, (1916), p. 295 (♀).

Type locality: Durban, NATAL (Marley).

18. *Polyrachis (Myrma) laboriosa* F. SMITH. SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 384 (♀). FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 351 (♀); 1916, Rev. Suisse Zool., XXIV, p. 447 (♀, ♀). See p. 258.

Polyrachis laboriosus F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 72 (♀), Pl. iv, figs. 21 and 22. ROGER, 1863, 'Verzeich. Formicid.,' p. 6. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 445. RADOSZKOWSKY, 1881, Journ. Sci. Ac. Lisboa, VIII, No. 31, p. 197. MAYR, 1886, Verh. Zool. Bot. Ges. Wien, XXXVI, p. 357. FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 195 (♀). ERN. ANDRÉ, 1887, Rev. d'Ent. Caen, VI, p. 286 (♀). EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 566 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 264. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 72. MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 153 (♀); 1896, Ent. Tidskr., XVII, pp. 250 (♀) and 255, Pl. iv, fig. 3 (nest). SJÖSTEDT, 1904, 'I Västafrikas Urskogar,' p. 501, fig. 3 (nest). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 393 (♀, ♀, ♂). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 151 (♀, ♀). FOREL, 1911, Rev. Zool. Afr., I, p. 282 (♀). WASMANN, 1911, 1^{er} Congr. Intern. Entom. Bruxelles (1910), II, Mém., p. 231, Pl. xvi, fig. 33 (nest); 1913, Ann. Rept. Smiths. Inst. for 1912, p. 472, Pl. x, fig. 33 (nest). FOREL, 1913, Rev. Zool. Afr., II, p. 349 (♀).

Polyrachis laboriosa M. L. SYKES, 1900, Trans. Manchester Microsc. Soc., XX, (1899), p. 88 (nest).

Type locality: SIERRA LEONE (D. F. Morgan).

IVORY COAST: Assinie (C. Alluaud). GOLD COAST: (Ganger). TOGO: Bismarekburg (Conradt). SOUTHERN NIGERIA: Lagos (F. Silvestri). CAMEROON: (Sjöstedt; H. Brauns); Bipindi (Zenker); Bibundi (Tessmann). SPANISH GUINEA: Alen (Tessmann). FRENCH CONGO: Cape Lopez (H. Brauns); Brazzaville (A. Weiss). PORTUGUESE CONGO: Chinchoxo (Falkenstein). BELGIAN CONGO: Congo da Lemba (R. Mayné); Kondué (Luja); St. Gabriel; Tshopo River near Stanleyville (Kohl); Stanleyville; Bafwasende; Niangara (Lang and Chapin). ANGOLA: (Welwitsch).

19. *Polyrachis (Myrma) lanuginosa* SANTSCHI.

Polyrachis lanuginosa SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 394, fig. 17 (♀).

Type locality: Mindouli, FRENCH CONGO (A. Weiss).

20. *Polyrachis (Myrma) lauta* SANTSCHI.

Polyrachis lauta SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 397, fig. 19 (♀). FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 450 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

20a. Var. *localis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 359 (♀).

Type locality: Congo da Lemba, BELGIAN CONGO (R. Mayné).

21. *Polyrachis (Myrma) maynei* FOREL.

Polyrachis maynei FOREL, 1911, Rev. Zool. Afr., I, p. 282 (♀).

Type locality: Congo da Lemba, BELGIAN CONGO (R. Mayné).

22. *Polyrachis (Myrma) militaris* (FABRICIUS). See p. 260.

Formica militaris FABRICIUS, 1781, 'Spec. Insect.', I, p. 493 (♀); 1787, 'Mantissa Insect.', I, p. 310 (♀). GMELIN, in Linnæus, 1790, 'Syst. Nat.', Ed. 13, I, 5, p. 2802. OLIVIER, 1791, 'Encycl. Méthod. Insect.', VI, p. 499. FABRICIUS, 1793, 'Ent. Syst.', II, p. 362 (♀). LATREILLE, 1802, 'Hist. Nat. Fourmis,' p. 124 (♀), Pl. iv, fig. 22. FABRICIUS, 1804, 'Syst. Piez.', p. 414 (♀).

Polyrachis militaris F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 72 (♂), Pl. III, fig. 5 and Pl. IV, fig. 36. ROGER, 1863, 'Verzeich. Formicid.,' p. 6. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 445; 1866, *ibid.*, XVI, p. 886 (♀). FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 119 (♀, ♀, ♂); 1886, Ann. Soc. Ent. Belgique, XXX, p. 194 (♀, ♀). ERN. ANDRÉ, 1887, Rev. d'Ent. Caen, VI, p. 287 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 265. H. STADELMANN, 1893, Mitth. Deutsch. Schutzgeb., VI, p. 217. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 72. EMERY, 1894, Mem. Accad. Sc. Bologna, (5) IV, p. 113 (♀); ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 1 (♀, ♀). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 154 (♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777. MAYR, 1896, Ent. Tidsskr., XVII, p. 252 (♀). FOREL, 1897, Abhandl. Senckenberg. Naturf. Ges., XXI, p. 188 (♀). SjöSTEDT, 1904, 'I Västafrikas Urskogar,' p. 507. FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 38 (♀); 1907, Mitt. Naturh. Mus. Hamburg, XXIV, p. 16 (♀); 1909, Ann. Soc. Ent. Belgique, LIII, p. 73 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 150 (♀, ♀). FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 372; 1913, Rev. Suisse Zool., XXI, p. 672 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 357 (♀).

Polyrachis militaris H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 38. M. L. SYKES, 1900, Trans. Manchester Microsc. Soc., XX, (1899), p. 91.

Type locality: Tropical Africa.

SIERRA LEONE: Samlia Falls, N'Gamie River (Mocquerys). TOGO: Bismarckburg (Büttner). SOUTHERN NIGERIA: Old Calabar (H. Brauns). CAMEROON: Bibundi (Tessmann); Longji (Paschen); Yaunde (Scheunemann); Mundame (Conradt). FERNANDO PO: (Conradt). FRENCH CONGO: Ogowe (Mocquerys). PORTUGUESE CONGO: Chinchoxo (Falkenstein). BELGIAN CONGO: Mayombe (de Brier); Congo da Lemba (R. Mayné); Sankuru (Luja); Leopoldville; Lukolela; Lie; Stanleyville; Avakubi; Lubila; Panga; Medje; Ngayu; Boyulu (Lang and Chapin). PORTUGUESE EAST AFRICA: Delagoa (Liengme); Mozambique (Fornasini). GERMAN EAST AFRICA: Tanga (Zimmer); Barikiwa (Ch. Schröder).

22. Var. *calabarica* FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 38 (♂). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 400 (♂); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 385 (♂).

Type locality: Old Calabar, SOUTHERN NIGERIA.

FRENCH GUINEA: Kakoulima (F. Silvestri). FRENCH CONGO: Brazzaville (A. Weiss).

22. Var. *ssibangensis* FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 38 (♂).

Polyrhachis militaris var. *sibangensis* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 400 (♂, ♀).

Type locality: Sibange, FRENCH CONGO (Soyaux).

FRENCH CONGO: Brazzaville (A. Weiss).

22a. Subsp. *bruta* SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 166 (♀).

Type locality: BELGIAN CONGO.

22b. Subsp. *cupreopubescens* FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., (2) XVI, p. 120 (♀). ERN. ANDRÉ, 1887, Rev. d'Ent. Caen, VI, p. 287 (♂). EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 566 (♂). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 154. ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 1 (♂). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777. FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 77 (♂); 1907, Ann. Mus. Nat. Hungarici, V, p. 38 (♂). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 400 (♂). FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 54 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 150 (♂, ♀); 1911, 'Wiss. Ergebn. Deutsch. Zentr. Afr. Exp. (1907-08)', III, p. 392 (♂). FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 372 (♂); 1912, Mitt. Naturh. Mus. Hamburg, XXIX, p. 180; 1913, Rev. Zool. Afr., II, p. 347 (♂). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 140 (♂). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 404 (♂). See p. 260.

Polyrhachis cupreopubescens DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 261.

Type locality: Africa; exact locality unknown.

SIERRA LEONE: Samlia Falls, River N'Gamie (Mocquerys). IVORY COAST: Assinie (C. Alluaud). SOUTHERN NIGERIA: Old Calabar (Duke). CAMEROON: Bibundi (Tessmann); Molundu (Schultze). SPANISH GUINEA: Eloby Island (H. Brauns). FRENCH CONGO: Sibange (Soyaux); Ogowe (Mocquerys); Brazzaville (A. Weiss). BELGIAN CONGO: Katanga (Lemaire); Kwesi to Kilo (Bayer); Avakubi (Schubotz); Medje; Avakubi (Lang and Chapin); Duma; Libenge (Schubotz). UGANDA: Entebbe (Schultze). BRITISH EAST AFRICA: Taveta (Alluaud and Jeannel). GERMAN EAST AFRICA: Mt. Kilimanjaro (Bornemisza); Arusha-chini (Katona); Amani; Bulko (H. Prell; Zimmer); Usambara.

22b₁. Var. *dido* Wm. M. WHEELER, new name. See p. 261.

Polyrhachis militaris subsp. *cupreopubescens* var. *argentatus* STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 150 (♂) (nec *Polyrhachis argentatus* F. Smith, 1858).

Type locality: Bibundi, CAMEROON (Zenker; Tessmann).

LIBERIA: Mt. Coffee (R. P. Currie). CAMEROON: Longji (Paschen).

22b₂. Var. *epinotalis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 357 (♀).

Type locality: Elisabethville, BELGIAN CONGO (Leplae).

22b₁. Var. *nkomoënsis* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 447 (♀, ♂). See p. 261.

Type locality: BELGIAN CONGO (Kohl).

BELGIAN CONGO: Akenge (Lang and Chapin).

22b₄. Var. *sankisiana* FOREL, 1913, Rev. Zool. Afr., II, p. 348 (♀).

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

22b₅. Var. *transversaria* FOREL, 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 77 (♀). See p. 261.

Type locality: LIBERIA (Hadler).

22c. Subsp. *striativentris* EMERY, 1892, Ann. Soc. Ent. France, LX, (1891), p. 566 (♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777. SANTSCHI, 1912, Ann. Soc. Ent. Belgique, LVI, p. 166 (♀).

Polyrhachis striativentris DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 270.

Type locality: Assinie, IVORY COAST (C. Alluaud).

IVORY COAST: (Lohier).

23. *Polyrhachis (Myrma) monista* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 384, fig. 34 (♀). FOREL, 1916, Rev. Suisse Zool., XXIV, p. 452 (♀, ♀).

Polyrhachis monista SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 398, fig. 20 (♀).

Type locality: FRENCH CONGO.

GOLD COAST: Aburi (F. Silvestri). BELGIAN CONGO: Kilongalonga (Kohl).

24. *Polyrhachis (Myrma) natalensis* SANTSCHI.

Polyrhachis natalensis SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 41 (♀).

Type locality: Stamford Hill, NATAL (I. Trägårdh).

25. *Polyrhachis (Myrma) nigrita* MAYR. See p. 267.

Polyrhachis nigrita MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 153 (♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777.

Type locality: Chama, GOLD COAST (H. Brauns).

BELGIAN CONGO: Akenge (Lang and Chapin).

26. *Polyrhachis (Myrma) otleti* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 449 (♀, ♀, ♂).

Type locality: St. Gabriel, BELGIAN CONGO (Kohl).

27. *Polyrhachis (Myrma) phidias* FOREL.

Polyrhachis phidias FOREL, 1910, Ann. Soc. Ent. Belgique, LIV, p. 450 (♀).

Type locality: Equatorial Africa, exact locality unknown.

28. *Polyrhachis (Myrma) revoili* ERN. ANDRÉ. FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 262 (♀); 1916, Rev. Suisse Zool., XXIV, p. 452 (♀, ♀, ♂). See p. 267.

Polyrhachis revoili ERN. ANDRÉ, 1886, Rev. d'Ent. Caen, VI, pp. 285 and 287 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 268. FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 71 (♂). ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 1 (♂, ♀). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 153. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 778. FOREL, 1913, Rev. Zool. Afr., II, p. 349 (♂, ♀).

Type locality: SOMALILAND (Revoil).

FRENCH CONGO: Ogowé (Mocquers). BELGIAN CONGO: (Kohl); Kondué (Luja); Malela (J. Bequaert). PORTUGUESE EAST AFRICA: Delagoa (Liengme). NATAL: Durban (C. B. Cooper); Port Natal (H. Brauns).

28₁. Var. *conduens* FOREL, 1915, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 351 (♂).

Type locality: Kondué, BELGIAN CONGO (Luja).

28₂. Var. *donisthorpei* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 453 (♂); 1917, Bull. Soc. Vaudoise Sc. Nat., (5) LI, p. 252 (♂).

Type locality: NORTHERN RHODESIA.

28₃. Subsp. *volkarti* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 453 (♀).

Type locality: BELGIAN CONGO (Kohl).

29. *Polyrhachis (Myrma) rufipalpis* SANTSCHI.

Polyrhachis rufipalpis SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 396 (♂).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

29₁. Subsp. *mayumbensis* FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 358 (♂).

Type locality: Kiniati, Mayombe, BELGIAN CONGO (R. Mayné).

30. *Polyrhachis (Myrma) schistacea* (GERSTÄCKER) SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' pp. 140 and 142 (♂).

Hoplomyrmus schistaceus GERSTÄCKER, 1858, Monatsb. Ak. Wiss. Berlin, p. 262 (♂).

Hoplomyrmus schistazeus GERSTÄCKER, in Peters, 1862, 'Reise n. Mossambique, Zool.,' V, p. 508 (♂), Pl. xxxii, fig. 6.

Polyrhachis schistazeus MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 446. GERSTÄCKER, 1871, Arch. f. Naturg., XXXVII, I, p. 354; 1873, in 'v. d. Decken's Reisen in Ost-Afrika, Gliederthiere,' p. 342 (♂).

Polyrhachis militaris subsp. *schistacea* ERN. ANDRÉ, 1887, Rev. d'Ent. Caen, VI, p. 288 (♂). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777.

Polyrhachis schistacea ROGER, 1863, 'Verzeich. Formicid.,' p. 6. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 268. BINGHAM, 1903, Ann. Mag. Nat. Hist., (7) XII, p. 63 (♂). DIXEY and LONGSTAFF, 1907, Trans. Ent. Soc. London, pp. 344 and 355. FOREL, 1907, Ann. Mus. Nat. Hungarici, V, p. 38 (♀); 1907, Mitth. Naturh. Mus. Hamburg, XXIV, p. 16 (♀); in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 92 (♂, ♀); 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 372 (♂); 1913, Rev. Zool. Afr., II, p. 348 (♂).

Polyrhachis militaris var. *schistazea* H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 38.

Type locality: Mozambique, PORTUGUESE EAST AFRICA (Peters).

PORTUGUESE CONGO: Chirchoxo. BELGIAN CONGO: Beni (Borgerhoff). BRITISH EAST AFRICA: Victoria Nyanza (Zimmer); Tiwi to Gazi (Alluaud and Jeannel); Mombasa (v. d. Decken); Fundu Island; Chake Chake, Pemba Island

(Voeltzkow). GERMAN EAST AFRICA: Mt. Kilimanjaro (Bornemisza); Barikiwa (Ch. Schröder); Tanga (Zimmer). ZANZIBAR: Mwera River; Bububu (Alluaud and Jeannel). NYASALAND: Fort Johnston (Rendall). RHODESIA: Matopopo Hills; Victoria Falls (Dixey and Longstaff). TRANSVAAL: Pretoria (Distant).

30. Var. *divina* FOREL, 1913, Rev. Zool. Afr., II, p. 348 (♀). SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' pp. 141 and 142 (♀). See p. 261.

Type locality: Pemba Island, BRITISH EAST AFRICA (Voeltzkow).

BELGIAN CONGO: Zambi; Boma; Thysville; Poko (Lang and Chapin). BRITISH EAST AFRICA: Mombasa; Bura, Wa-Taita, 1050 m. (Alluaud and Jeannel). GERMAN EAST AFRICA: Darfessalaam. ZANZIBAR.

30₁. Var. *gagatoides* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 314; 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 142 (♀).

Type locality: FRENCH CONGO (A. Weiss).

30₂. Subsp. *atrociliata* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 314 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 385 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' pp. 141 and 142 (♀). STITZ, 1916, 'Ergebn. Zweit. Deutsch. Zentr. Afr. Exp. 1910-11,' I, p. 404 (♀).

Type locality: Banzyville, BELGIAN CONGO (Augustin).

FRENCH GUINEA: Konakry (F. Silvestri). UGANDA: Ibanda, Mt. Ruwenzori, 1400 m. (C. Alluaud). BELGIAN CONGO: Duma (Schubotz). FRENCH CONGO: Fort Crampel (Schubotz).

30₃. Var. *benguensis* SANTSCHI, 1913, Ann. Soc. Ent. Belgique, LVII, p. 314 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' pp. 141 and 142 (♀). See p. 261.

Type locality: BENGUELA (J. Cruchet).

BELGIAN CONGO: Garamba; Yakuluku (Lang and Chapin).

30₄. Subsp. *congolensis* (SANTSCHI) FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 357 (♀).

Polyrhachis schistacea var. *congolensis* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 399 (♀).

Polyrhachis gagates subsp. *congolensis* FOREL, 1913, Rev. Zool. Afr., II, p. 282 (♀).

Polyrhachis (*Myrma*) *gagates* var. *congolensis* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 139 (♀).

Type locality: Mandouga, FRENCH CONGO (A. Weiss).

FRENCH CONGO: Comba Ibre (A. Weiss). BELGIAN CONGO: Congo da Lemba (R. Mayné). BRITISH EAST AFRICA: Bura, Wa-Taita, 1050 m.; Voi (Alluaud and Jeannel).

30₅. Subsp. *fracta* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' pp. 141 and 142 (♀).

Type locality: Fort Hall, 1330 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

30₆. Var. *subplana* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 142 (♀).

Type locality: Gazi, BRITISH EAST AFRICA (Alluaud and Jeannel).

30₇. Subsp. *medusa* FOREL, 1897, Abhandl. Senckenberg. Naturf. Ges., XXI, p. 206 (♀, ♀, ♂); 1901, Mitt. Naturh. Mus. Hamburg, XVIII, p. 78.

Polyrhachis militaris subsp. *schistacea* var. *medusa* EMERY, 1898, Rend. Accad. Sc. Bologna, N. S., II, p. 228.

Polyrhachis medusa FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 92 (♀, ♀).

Polyrhachis (Myrma) medusa SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 140 (♀).

Type locality: ZANZIBAR (Voeltzkow).

GERMAN EAST AFRICA: Mafia Island (Voeltzkow); Lewa Mambaa (Stuhlmann). BRITISH EAST AFRICA: Ramisi River; Voi, Taita (Alluaud and Jeannel); Chake Chake, Pemba Island (Voeltzkow).

30e. Subsp. *nigriseta* (SANTSCHI) FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 357.

Polyrhachis nigriseta SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 399 (♀). FOREL, 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 272 (♀).

Type locality: Mindouli, FRENCH CONGO (A. Weiss).

WESTERN AFRICA: (Ilg).

30e. Var. *clariseta* (SANTSCHI) FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, p. 357 (♀).

Polyrhachis nigriseta var. *clariseta* SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 400 (♀).

Polyrhachis (Myrma) nigriseta var. *clariseta* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidae,' p. 140 (♀).

Type locality: Mandouga, FRENCH CONGO (A. Weiss).

BELGIAN CONGO: Congo da Lemba (R. Mayné). BRITISH EAST AFRICA: Voi, Taita (Alluaud and Jeannel).

30f. Subsp. *rugulosa* (MAYR) FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 92 (♀); 1910, Ann. Soc. Ent. Belgique, LIV, p. 450 (♀). STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 151 (♀). FOREL, 1913, Ann. Soc. Ent. Belgique, LVII, pp. 147 (♀) and 357 (♀); 1913, Rev. Zool. Afr., II, p. 349 (♀); 1913, Deutsch. Ent. Zeitschr., Beih., p. 225 (♀).

Polyrhachis rugulosus MAYR, 1862, Verh. Zool. Bot. Ges. Wien, XII, p. 685 (♀), Pl. I, fig. 7. ROGER, 1863, 'Verzeich. Formicid.,' p. 9. EMERY, 1889, Bull. Soc. Ent. Italiana, XXI, p. 69, footnote. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 268. EMERY, 1894, Mem. Accad. Sc. Bologna, (5) IV, p. 113 (♀). FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 72. EMERY, 1895, Ann. Soc. Ent. France, LXIV, p. 55 (♀); 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777; 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 605 (♀, ♀, ♂). SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 400. EMERY, 1912, Annuar. Mus. Zool. Univ. Napoli, N. S., III, No. 26, p. 2.

Polyrhachis carinatus F. SMITH, 1858, Journ. Proc. Linn. Soc. London, Zool., II, p. 59; 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 71 (♀), Pl. iv, figs. 48 and 49 (nec *Formica carinata* Fabricius).

Polyrhachis militaris subsp. *cafrorum* FOREL, 1879, Bull. Soc. Vaudoise Sc. Nat., XVI, p. 120 (♀, ♀, ♂); 1886, Ann. Soc. Ent. Belgique, XXX, p. 194 (♀). ERN. ANDRÉ, 1887, Rev. d'Ent. Caen, VI, p. 287 (♀).

Polyrhachis cafrorum FOREL, 1892, Mitth. Schweiz. Ent. Ges., VIII, p. 350 (♀). EMERY, 1892, Ann. Mus. Civ. Genova, XXXII, p. 122. DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 260.

Polyrhachis militaris var. *rugulosa* MAYR, 1893, Jahrb. Hamburg. Wiss. Anst., X, 2, p. 197; 1895, Ann. Naturh. Hofmus. Wien, X, p. 154; in Sjöstedt, 1907, 'Exped. Kilimandjaro, Meru, etc.,' II, 8, p. 22.

Polyrhachis militaris var. *rugulosa* H. STADELMANN, 1898, 'Deutsch-Ost-Afrika,' IV, Hym., p. 38.

Polyrhachis schistacea var. *rugulosa* FOREL, 1911, Bull. Soc. Vaudoise Sc. Nat., (5) XLVII, p. 372 (♀).

Polyrhachis (Myrma) schistacea var. *rugulosa* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' pp. 141 and 142 (♀); 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 385 (♀).

Type locality: Port Natal, NATAL (not Brazil as originally given by Mayr).

TOGO: Bismarckburg (Conradt). CAMEROON: Etombé (Adametz). FRENCH CONGO: Loango (H. Brauns); Brazzaville; Comba Ibire; Mandouga (A. Weiss). PORTUGUESE CONGO: Chinchoxo (Falkenstein). BELGIAN CONGO: Kwezi to Kilo (Bayer); Kimpoko (Büttner); Old Kasongo (J. Bequaert); Congo da Lemba (R. Mayné); Boma (H. Brauns; Styczinski); Luapula River. ANGOLA: Quifangondo (F. Silvestri). RHODESIA: (G. Arnold). TRANSVAAL: Makapan; Hamman's Kraal (E. Simon); Valdezia (P. Berthoud). CAPE PROVINCE: Kimberley (E. Simon). PORTUGUESE EAST AFRICA: Delagoa (Liengme; A. Müller); Mozambique (Fornasini); Quilimane (Stuhlmann). GERMAN EAST AFRICA: Darassalaam (H. Brauns; A. Müller); Kibonoto, Kilimanjaro, 1000-3000 m.; Meru (Sjöstedt); Mafia Island (Voeltzkow). ZANZIBAR: (Stuhlmann). BRITISH EAST AFRICA: Kitui (Hildebrandt); Bura (H. Brauns); Mbuyuni, Pori, 1110 m. (Alluaud and Jeannel). SOMALILAND: (C. Keller); Magala Re Umberto (Ruspoli). ABYSSINIA: Webi (Bricchetti-Robecchi).

30f. Var. *divinoides* FOREL, 1913, Rev. Zool. Afr., II, pp. 348 and 349 (♀). BEQUAERT, *ibid.*, p. 431. See p. 261.

Polyrhachis (Myrma) schistacea var. *divinoides* SANTSCHI, 1914, Boll. Lab. Zool. Gen. Agrar. Portici, VIII, p. 385 (♀); 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 142 (♀).

Type locality: Sankisia, BELGIAN CONGO (J. Bequaert).

BELGIAN CONGO: Boma (F. Silvestri); Banana (Lang and Chapin). TRANSVAAL.

30g. Subsp. *schlüteri* FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 72.

Polyrhachis militaris subsp. *schlüteri* FOREL, 1886, Ann. Soc. Ent. Belgique, XXX, p. 195 (♀). ERN. ANDRÉ, 1886, Rev. d'Ent. Caen, VI, p. 288 (♀). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 154 (♀). EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777. STITZ, 1910, Mitt. Zool. Mus. Berlin, V, p. 151 (♀).

Polyrhachis schlüteri DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 268.

Polyrhachis schistacea subsp. *schlüteri* SANTSCHI, 1914, Med. Göteborgs Mus. Zool. Afd., III, p. 42 (♀).

Polyrhachis (Myrma) schlüteri SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 143 (♀).

Type locality: GERMAN EAST AFRICA (Schlüter).

TOGO: Bismarckburg (Conradt). NATAL: Stamford Hill (I. Trägårdh); Port Natal (H. Brauns). ZULULAND: Dukudu (I. Trägårdh). PORTUGUESE EAST AFRICA: Delagoa (P. Berthoud; Liengme). BRITISH EAST AFRICA: Voi (C. Alluaud).

30h. Var. *indigenus* (FOREL).

Polyrhachis (Myrma) schlüteri var. *indigenus* FOREL, 1914, Bull. Soc. Vaudoise Sc. Nat., (5) L, p. 261 (♀).

Type locality: Durban, NATAL (G. Arnold).

30gs. Var. *plebeia* (SANTSCHI).

Polyrhachis (Myrma) schluteri var. *plebeia* SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 143 (♀).

Type locality: Taveta, 750 m., BRITISH EAST AFRICA (Alluaud and Jeannel).

31. *Polyrhachis (Myrma) schoutedeni* SANTSCHI, 1919, Rev. Zool. Afr., VI, p. 249 (♀).

Type locality: Dolo, BELGIAN CONGO (F. Chaltin).

32. *Polyrhachis (Myrma) spinicola* FOREL.

Polyrhachis spinicola FOREL, 1894, Mitth. Schweiz. Ent. Ges., IX, p. 70 (♀, ♀).

EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777.

Type locality: Delagoa, PORTUGUESE EAST AFRICA (Junod).

PORTUGUESE EAST AFRICA: Muculla (A. Müller).

33. *Polyrhachis (Myrma) spitteleri* FOREL, 1916, Rev. Suisse Zool., XXIV, p. 450, fig. 6 (♀).

Type locality: BELGIAN CONGO (Kohl).

34. *Polyrhachis (Myrma) sulcata* ERN. ANDRÉ.

Polyrhachis sulcata ERN. ANDRÉ, 1895, Rev. d'Ent. Caen, XIV, p. 1 (♀). MAYR, 1895, Ann. Naturh. Hofmus. Wien, X, p. 154 (♀); EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777.

Type locality: Ogowé, FRENCH CONGO (Mocquerys).

CAMEROON: (H. Brauns). BELGIAN CONGO.

35. *Polyrhachis (Myrma) viscosa* F. SMITH. SANTSCHI, 1914, 'Voy. Alluaud et Jeannel Afr. Orient., Formicidæ,' p. 143 (♀, ♀). See p. 267.

Polyrhachis viscosus F. SMITH, 1858, 'Cat. Hym. Brit. Mus.,' VI, p. 71 (♀), Pl. IV, fig. 41. ROGER, 1863, 'Verzeich. Formicid.,' p. 6. MAYR, 1863, Verh. Zool. Bot. Ges. Wien, XIII, p. 447; 1886, *ibid.*, XXXVI, p. 357. ERN. ANDRÉ, 1887, Rev. d'Ent. Caen, VI, p. 287 (♀). DALLA TORRE, 1893, 'Cat. Hym.,' VII, p. 271. EMERY, 1896, Mem. Accad. Sc. Bologna, (5) V, p. 777; 1897, Ann. Mus. Civ. Genova, XXXVIII, p. 605 (♀, ♀); 1899, *ibid.*, XXXIX, p. 501 (♀). FOREL, in Voeltzkow, 1907, 'Reise in Ostafrika,' II, p. 92 (♀); 1910, Zool. Jahrb. Abt. Syst., XXIX, p. 272 (♀); 1913, Ann. Soc. Ent. Belgique, LVII, p. 143 (♀, ♀).

Polyrhachis antinorii EMERY, 1877, Ann. Mus. Civ. Genova, IX, p. 365, fig. (♀). ERN. ANDRÉ, 1887, Rev. d'Ent. Caen, VI, p. 287 (♀).

Type locality: Port Natal, NATAL.

SOUTHERN RHODESIA: Redbank (G. Arnold). GERMAN EAST AFRICA: Mafia Island (Voeltzkow). BRITISH EAST AFRICA: Voi, Taita; Bura, 1050 m. (Alluaud and Jeannel). SOMALILAND: Magala Re Umberto (Ruspoli). ABYSSINIA: Dimé to Bass Narok (V. Bottego); Coromma; Ganale (Ruspoli). ERITREA: Sciotel; Keren (Beccari); Ghinda (K. Escherich). BELGIAN CONGO: Ngayu (Lang and Chapin).

36. *Polyrhachis (Myrma) weissii* SANTSCHI.

Polyrhachis weissii SANTSCHI, 1910, Ann. Soc. Ent. France, LXXVIII, (1909), p. 395, fig. 18 (♀).

Type locality: Brazzaville, FRENCH CONGO (A. Weiss).

37. *Polyrhachis (Myrma) wellmani* FOREL.

Polyrhachis wellmani FOREL, 1909, Ann. Soc. Ent. Belgique, LIII, p. 68 (♀).

Type locality: BENGUELA (C. Wellman).